

DOE/ID-12082(22)  
INL/RPT-23-74740, Rev 1  
October 2023



2022

# IDAHO NATIONAL LABORATORY ANNUAL SITE ENVIRONMENTAL REPORT

# ERRATA SHEET

**Issue Date:** August 6, 2024

“2022 Idaho National Laboratory Annual Site Environmental Report”  
(INL/RPT-23-74740, Rev 1)

Location:	Reads:	Description of Changes/Corrections:
Chapter 2, page 2-25, second paragraph	To that end, the INL Site and EJP have worked diligently to incorporate Indigenous and Traditional Ecological Knowledge (ITEK) into laboratory policies, procedures, and practices. ITEK is a repository of natural and ecological knowledge refined through thousands of years of tribal stewardship. ITEK-informed decision making is a federal priority and is recognized as one of the many important bodies of knowledge that contributes to the scientific, technical, social, and economic advancements of the U.S. and our understanding of the natural world. ITEK-informed science and decision making will be essential as the nation navigates climate change and energy transition. These global challenges demand disparate knowledge and solutions to inform and work cohesively with the scientific process toward a sustainable future.	The paragraph was incorrectly included in the report when it was published in 2023. The error was identified during preparation of the calendar year 2023 report and the paragraph has been removed from this version.



DOE/ID-12082(23)  
INL/RPT-23-74740, Rev 1

# **2022**

## **SITE ENVIRONMENTAL REPORT**

### **Idaho National Laboratory**

## **October 2023**

**Prepared by**

**Battelle Energy Alliance  
For the U.S. Department of Energy  
Under Contract No. DE-AC07-05ID14517**

**Idaho National Laboratory  
Idaho Falls, ID 83415**

**DISCLAIMER**

This information was prepared as an account of work sponsored by an agency of the U.S. Government. Neither the U.S. Government nor any agency thereof, nor any of their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness, of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. References herein to any specific commercial product, process, or service by trade name, trade mark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the U.S. Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the U.S. Government or any agency thereof.

# Acknowledgements:



The following people have provided primary authorship and review of this report:

- Amy Forman, Blane Teckmeyer, Brian Donovan, Bryan Bybee, Colby Kramer, Elizabeth Cook, Nicholas Holmer, Jason Daley, Jeff Sondrup, Jeremy Shive, Kevin Claver, Kira Overin, Kristin Kaser, Peggy Scherbinske, Rajkumar Devasirvatham, Samuel Williams, and Tom Rackow with Battelle Energy Alliance, LLC (BEA).
- Christopher Campbell, Danielle Millward, Eric Traub, Jennifer Thomas, Kristina Alberico, Sarah Thompson, Sue Vilord, and Vanessa Morgan with the Idaho Environmental Coalition, LLC (IEC).
- Betsy Holmes, Charles Ljungberg, Jason Anderson, Jimmy Laner, Nicole Hernandez, Nicole Badrov, Shelby Goodwin, Steve Wahnschaffe, Tauna Butler, Chase Hartzell, Guy Backstrom, Daphne Larsen, Nick Balsmeier, Ty Sanders, Tommy Thompson, Doug Herzog, Danielle Miller, Doug Pruitt, Chris Harvey, Chauntel Simmons, and Trent Neville with the U.S. Department of Energy, Idaho Operations Office (DOE-ID).
- Jason Rich with the National Oceanic and Atmospheric Administration, Air Resources Laboratory, Special Operations and Research Division.
- Brian Twining, Kerri Treinen, and Allison Trcka with the U.S. Geological Survey.
- Maps provided by Dan Mahnami with IEC and by Jeremy Shive and Kurt Edwards with BEA.
- Technical editing of this report was provided by Brande Hendricks, David Livingston, Nikki Peterson, and Rachel Hansen with BEA. Additional technical editing was performed by Emily Slike and Jim Nelson with IEC.
- Publishing layout was executed by Brande Hendricks with BEA and Lauren Perttula with Red Inc.

The primary authors would like to thank all of those who provided data for the completion of this document. We wish to thank the following people for their assistance:

- Anne Dustin, Caitlin Nate, David Twamley, Dawn Davis, Jacob Yedica, Jennifer Jackson, Kim Scully, Kris Murray, Lisbeth Mitchell, Maryl Fisher, Morris Hall, Paul Apolinar Velasquez, Rob Black, Sarah Baccus, and Scott Lee with BEA.
- William H. Clark with the Orma J. Smith Museum of Natural History, College of Idaho.
- Vincent A. Cobb with the Department of Biology, Middle Tennessee State University.
- Matthew J. Germino, Toby Maxwell, and Marie-Anne DeGraff with U.S. Geological Survey Forest and Rangeland Ecosystem Science Center and Boise State University.
- Ken Aho with Idaho State University.
- Jennifer S. Forbey with Boise State University.

We would like to thank Kristin Kaser with BEA for contributing photographs that were used in this report.



*Horned lark*

# To Our Readers:



The Idaho National Laboratory (INL) Site Environmental Report for Calendar Year 2022 is an overview of environmental activities conducted on and in the vicinity of the INL Site from January 1 through December 31, 2022. This report includes the following components:

- Effluent monitoring and environmental surveillance of air, water, soil, vegetation, biota, and agricultural products for radioactivity. The results are compared with historical data, background measurements, and applicable standards and requirements to verify that the INL Site does not adversely impact the environment or the health of humans or biota.
- A summary of environmental management systems in place to protect air, water, land, and other natural and cultural resources potentially impacted by INL Site operations.
- Ecological monitoring and other scientific research conducted onsite that may be of interest to the reader.

The report addresses three general levels of reader interest:


- The first level is a brief summary with a take-home conclusion. This is presented in the chapter highlights text box at the beginning of each chapter. There are no tables, figures, or graphs in the highlights. This section is intended to highlight general findings for an audience with a limited scientific background.
- The second level is a more in-depth discussion with figures, summary tables, and summary graphs accompanying the text. The chapters of the annual report represent this level, which requires some familiarity with scientific data and graphs. A person with some scientific background can read and understand this report after reading the section entitled, “Helpful Information.”
- The third level includes links to supplemental and technical reports and websites that support the annual report. This level is directed toward scientists who would like to see original data and more in-depth discussions of the methods used and results.

The links to these reports may be found in the Publications tab of the webpage at <https://inl.gov/environmental-publications>.

The INL contractor is responsible for contributing to and producing the INL Annual Site Environmental Report. Environmental monitoring within the INL Site boundaries is primarily the responsibility of INL and Idaho Cleanup Project (ICP) contractors.

Major contributors to the annual INL Site Environmental Report include the INL contractor (BEA); ICP contractor (Idaho Environmental Coalition, LLC); U.S. Department of Energy–Idaho Operations Office; National Oceanic and Atmospheric Administration, Air Resources Laboratory, Special Operations and Research Division; and U.S. Geological Survey. Links to their websites are as follows:

- INL (<https://www.inl.gov/>)
- ICP (<https://idaho-evnvironmental.com>)
- U.S. Department of Energy–Idaho Operations (<https://www.id.energy.gov/>)
- Special Operations and Research Division of National Oceanic and Atmospheric Administration’s Air Resources Laboratory (<https://www.noaa.inl.gov>)
- U.S. Geological Survey (<https://www.usgs.gov/centers/idaho-water-science-center>).



*Northwest indian paintbrush and bumblebee*



# Executive Summary:



## Introduction

In operation since 1949, the INL Site is a U.S. Department of Energy (DOE) reservation located in the southeastern Idaho desert, approximately 25 miles west of Idaho Falls (Figure ES-1). At 890 square miles (569,135 acres), the INL Site is roughly 85% of the size of Rhode Island. It was established in 1949 as the National Reactor Testing Station, and for many years, it was the site of the largest concentration of nuclear reactors in the world. Fifty-two nuclear reactors were built, including the Experimental Breeder Reactor-I, which in 1951 produced the first usable amounts of electricity generated by nuclear power. Researchers pioneered many of the world's first nuclear reactor prototypes and advanced safety systems at the INL Site. During the 1970s, the laboratory's mission broadened into other areas such as biotechnology, energy and materials research, and conservation and renewable energy.

Today, INL is a science-based, applied engineering national laboratory dedicated to supporting DOE's nuclear and energy research, science, and national defense missions.



**Figure ES-1. Regional location of the INL Site.**



INL's mission is to discover, demonstrate, and secure innovative nuclear energy solutions, other clean energy options, and critical infrastructure with a vision to change the world's energy future and secure the nation's critical infrastructure.

To mitigate environmental impacts and clear the way for the facilities required for the new nuclear energy research mission, the ICP has been charged with the environmental cleanup of the legacy wastes generated from World War II-era conventional weapons testing, government-owned reactors, and spent fuel reprocessing. The overarching aim of the project is to reduce risks to workers and production facilities, the public, and the environment and to protect the Snake River Plain Aquifer.

## PURPOSE OF THE INL SITE ENVIRONMENTAL REPORT

The INL Site's operations, as well as the ongoing cleanup mission involve a commitment to environmental stewardship and full compliance with environmental protection laws. As part of this commitment, the INL Site Environmental Report is prepared annually to inform the public, regulators, stakeholders, and other interested parties of the INL Site's environmental performance during the year. This report is published for U.S. Department of Energy, Idaho Operations Office (DOE-ID) in compliance with DOE O 231.1B, "Environment, Safety and Health Reporting." The purpose of the report is to provide the following:

- Present the INL Site, mission, and programs
- Report compliance status with applicable federal, state, and local regulations
- Describe the INL Site environmental programs and activities
- Summarize results of environmental monitoring
- Discuss potential radiation doses to the public residing in the vicinity of the INL Site
- Report on ecological monitoring and research conducted by contractors and affiliated agencies and by independent researchers through the Idaho National Environmental Research Park
- Describe quality assurance methods used to ensure confidence in monitoring data
- Provide supplemental technical data and reports that support the INL Site Environmental Report (<https://inl.gov/environmental-publications>).

## MAJOR INL SITE PROGRAMS AND FACILITIES

INL is a combination of all operating contractors and the U.S. Department of Energy, Idaho Operations Office (DOE-ID), and includes the Idaho Falls campus and the research and industrial complexes termed the "INL Site" that is located 50 miles west of Idaho Falls, Idaho. For the purpose of this report, INL consists of those facilities operated by Battelle Energy Alliance, LLC (INL contractor), or by the Idaho Environmental Coalition, LLC (Idaho Cleanup Project [ICP] contractor). INL and ICP contractors are referred to by their noted acronyms and include all facilities under their individual responsibilities.

The INL Site consists of several primary facilities situated on an expanse of otherwise undeveloped terrain. Buildings and structures at the INL Site are clustered within these facilities, which are typically less than a few square miles in size and separated from each other by miles of undeveloped land. In addition, DOE-ID owns or leases laboratories and administrative offices in the city of Idaho Falls, Idaho, some 25 miles east of the INL Site border. About 30% of employees work in administrative, scientific support, and non-nuclear laboratory programs at offices in Idaho Falls, Idaho.

The major facilities at the INL Site are the Advanced Test Reactor (ATR) Complex, Central Facilities Area (CFA), Critical Infrastructure Test Range Complex, Idaho Nuclear Technology and Engineering Center (INTEC), Materials and Fuels Complex (MFC), Naval Reactors Facility (NRF), Radioactive Waste Management Complex (RWMC), and Test Area North (TAN), which includes the Specific Manufacturing Capability (SMC). The Research and Education Campus is located in Idaho Falls, Idaho. The locations of major facilities are shown in Figure ES-2, and their missions are outlined in Table ES-1.




Figure ES-2. INL Site facilities.



**Table ES-1. Major INL Site areas and missions.**

MAJOR INL SITE AREA <sup>a</sup>	OPERATED BY	MISSION
Advanced Test Reactor Complex	INL	Research and development of nuclear reactor technologies. Home of the ATR, a DOE Nuclear Science User Facility and the world's most advanced nuclear test reactor. The ATR provides unique irradiation capabilities for nuclear technology research and development.
Central Facilities Area	INL	INL support for the operation of other INL Site facilities and management responsibility for the balance of the INL outside of the facility boundaries.
Critical Infrastructure Test Range Complex	INL	Supports National and Homeland Security missions of the laboratory, including program and project testing (i.e., critical infrastructure resilience and nonproliferation testing and demonstration).
Idaho Nuclear Technology and Engineering Center	ICP	Dry and wet storage of spent nuclear fuel; management of high-level waste calcine and sodium-bearing liquid waste; and operation of the Idaho Comprehensive Environmental Response, Compensation and Liability Act Disposal Facility, including a landfill, evaporation ponds, and a staging and treatment facility.
Materials and Fuels Complex	INL	Research and development of nuclear fuels. Pyro-processing, which uses electricity to separate waste products in the recycling of nuclear fuel, is also researched here. Nuclear batteries for use on the nation's space missions are made at MFC.
Radioactive Waste Management Complex	ICP	Environmental remediation and waste treatment, storage, and disposal for wastes generated at the INL Site and other DOE sites. The Advanced Mixed Waste Treatment Project characterizes, treats, and packages transuranic waste for shipment out of Idaho to permanent disposal facilities. Location of the Integrated Waste Treatment Unit, a first-of-a-kind, 53,000-square-foot facility that will treat 900,000 gallons of liquid radioactive and hazardous waste that has been stored in underground storage tanks.
Research and Education Campus	INL	Located in Idaho Falls, Idaho, the Research and Education Campus is home to DOE's Radiological and Environmental Sciences Laboratory, INL administration, the INL Research Center, the Center for Advanced Energy Studies, and other energy and security research programs. Research is conducted at INL Reach Center in robotics, genetics, biology, chemistry, metallurgy, computational science, and hydropower. Center for Advanced Energy Studies is a research and education partnership between Boise State University, INL, Idaho State University, and University of Idaho to conduct energy research and address the looming nuclear energy work-force shortage.
Test Area North/Specific Manufacturing Capability	INL	Several historic nuclear research and development projects were conducted at TAN. Major cleanup and demolition of the facility was completed in 2008, and the current mission is the manufacture of tank armor for the U.S. Army's battle tanks at the SMC for the U.S. Department of Defense.

a. NRF is also located onsite. It is operated for Naval Reactors by Fluor Marine Propulsion, LLC. The Naval Nuclear Propulsion Program is exempt from DOE requirements and is therefore not addressed in this report.



## ENVIRONMENTAL PROTECTION PROGRAMS

Directives, orders, guides, and manuals are DOE's primary means of establishing policies, requirements, responsibilities, and procedures for DOE offices and contractors. Among these are a series of orders directing each DOE site to implement sound stewardship practices that are protective of the public and the environment. These orders require the implementation of an environmental management system (EMS), a Site Sustainability Plan, a radioactive waste management program, and programs addressing radiation protection of the public and the environment. The INL and ICP contractors have each established and implemented an EMS and have contributed to the INL Site Sustainability Plan, as required by DOE and executive orders. Each EMS integrates environmental protection, environmental compliance, pollution prevention, and waste minimization into work planning and execution throughout all work areas. The INL Site Sustainability Plan contains strategies and activities that will lead to continual greenhouse gas reductions as well as energy, water, and transportation fuels efficiency at the INL Site. Plan requirements are integrated into each INL Site contractor's Integrated Safety Management System and EMS.

## ENVIRONMENTAL RESTORATION

Environmental restoration at the INL Site is conducted under the Federal Facility Agreement and Consent Order (FFA/CO) among DOE, the state of Idaho, and the U.S. Environmental Protection Agency (EPA). The FFA/CO governs the INL Site's environmental remediation activities. It specifies actions that must be completed to safely clean up sites at INL in compliance with the Comprehensive Environmental Response, Compensation, and Liability Act and with the corrective action requirements of the Resource Conservation and Recovery Act. The INL Site is divided into ten Waste Area Groups (WAGs) as a result of the FFA/CO, and each WAG is divided into smaller cleanup areas called operable units. Since the FFA/CO was signed in 1991, the INL Site has cleaned up sites containing asbestos, acids and bases, radionuclides, unexploded ordnance and explosive residues, polychlorinated biphenyls, heavy metals, and other hazardous materials.

Comprehensive remedial investigation/feasibility studies have been conducted at all WAGs and closeout activities have been completed at six WAGs. In 2022, all institutional controls and operational and maintenance requirements were maintained, and active remediation continued on WAGs 1, 3, and 7.

## RADIATION DOSE TO THE PUBLIC AND BIOTA FROM INL SITE RELEASES

Humans, plants, and animals potentially receive radiation doses from various INL Site operations. DOE sets dose limits for the public and biota to ensure that exposure to radiation from site operations are not a health concern. Potential radiological doses to the public from INL Site operations were calculated to determine compliance with pertinent regulations and limits (Table ES-2). The calculated dose to the maximally exposed individual in 2022 from the air pathway was 0.018 mrem (0.18  $\mu$ Sv), which is well below the 10-mrem standard established by the Clean Air Act. The maximally exposed individual is a hypothetical member of the public who could receive the maximum possible dose from INL Site releases as determined by the air dispersion model. This person is assumed to live at a location east of INL's east entrance and south of Highway 20. For comparison, the dose from natural background radiation was estimated in 2022 to be 384 mrem (3,840  $\mu$ Sv) to an individual living on the Snake River Plain.

The maximum potential population dose to the approximately 349,242 people residing within an 80 km (50 mi) radius of any INL Site facility was calculated as 0.019 person-rem (0.00019 person-Sv), below that expected from exposure to background radiation (134,109 person-rem or 1,341 person-Sv). The 50 mi population dose calculated for 2022 is lower than that calculated for 2021 (0.028 person-rem or 0.00028 person-Sv).

The maximum potential individual dose from consuming waterfowl contaminated at the INL Site, based on the highest concentrations of radionuclides measured in edible tissue of samples collected near the ATR Complex ponds, was estimated to be 0.0009 mrem (0.009  $\mu$ Sv). In 2022, none of the game samples collected (e.g., four elk and one pronghorn) had a detectable concentration of cesium-137 ( $^{137}\text{Cs}$ ) or other human-made radionuclides. When the dose estimated for the air pathway was summed with the dose from consuming contaminated waterfowl, assuming that the waterfowl is eaten by the same hypothetical individual, the representative person off the INL Site could potentially receive a total dose of 0.019 mrem (0.19  $\mu$ Sv) in 2022. This is 0.019% of the DOE health-based dose limit of 100 mrem/yr (1 mSv/yr) from all pathways for the INL Site.



**Table ES-2. Contribution to estimated annual dose from INL Site facilities by pathway (2022).**

PATHWAY	ANNUAL DOSE TO MAXIMALLY EXPOSED INDIVIDUAL		PERCENT OF DOE 100 mrem/YR LIMIT <sup>a</sup>	ESTIMATED POPULATION DOSE		POPULATION WITHIN 80 km	ESTIMATED BACKGROUND RADIATION POPULATION DOSE (PERSON-rem) <sup>b</sup>
	(mrem)	( $\mu$ Sv)		(PERSON-mrem)	(PERSON-Sv)		
Air	0.018	0.18	0.018	0.019	0.00019	349,242	134,109
Waterfowl	0.0009	0.009	0.0009	NA <sup>c</sup>	NA	NA	NA
Big game animals	0.000	0.00	NA	NA	NA	NA	NA
<b>Total pathways</b>	<b>0.019</b>	<b>0.19</b>	<b>0.019</b>	<b>0.019</b>	<b>0.00019</b>	<b>NA</b>	<b>NA</b>

- a. The DOE public dose limit from all sources of ionizing radiation and exposure pathways that could contribute significantly to the total dose is 100 mrem/yr (1 mSv/yr) total effective dose equivalent. It does not include dose from background radiation.
- b. The individual background dose was estimated to be 384 mrem or 0.384 rem in 2022, as shown previously in Table 7-8. The background population dose is calculated by multiplying the individual background dose by the population within 80 km (50 mi) of the INL Site.
- c. NA = Not applicable.

Tritium has been previously detected in two U.S. Geological Survey (USGS) monitoring wells located onsite along the southern boundary. A hypothetical individual ingesting the maximum concentration of tritium (3,970 pCi/L) via drinking water from these wells would receive a dose of approximately 0.2 mrem (0.002 mSv) in one year. This is an unrealistic pathway to humans because there are no drinking water wells located along the southern boundary of the INL Site. The maximum contaminant level established by EPA for tritium (20,000 pCi/L) corresponds to a dose of approximately 4 mrem (0.04 mSv [40  $\mu$ Sv/yr]).

A dose to a maximally exposed individual located in Idaho Falls, Idaho, near the DOE Radiological and Environmental Sciences Laboratory and the INL Research Center, within the Research and Education Campus, was calculated for compliance with the Clean Air Act. For 2022, the dose was conservatively estimated to be 0.004 mrem (0.04  $\mu$ Sv), which is less than 0.1% of the 10-mrem/yr federal standard.

Doses were also evaluated for nonhuman biota at the INL Site using a graded approach. Based on the conservative screening calculations, there is no evidence that INL Site-related radioactivity in soil or water is harming populations of plants or animals.

## ENVIRONMENTAL COMPLIANCE

One measure of the achievement of the environmental programs at the INL Site is compliance with applicable environmental regulations, which have been established to protect human health and the environment. INL Site and DOE-ID programs compliance with federal and state environmental protection requirements, such as statutes, acts, agreements, executive orders and DOE directives are presented in Table 2-1.



## ENVIRONMENTAL MONITORING OF AIR

Airborne releases of radionuclides from INL Site operations are reported annually in a document prepared in accordance with the 40 CFR 61, Subpart H, "National Emission Standards for Emissions of Radionuclides Other than radon from Department of Energy Facilities." An estimated total of 357 curies ( $1.32 \times 10^{13}$  Bq) of radioactivity, primarily in the form of short-lived noble gas isotopes, were released as airborne effluents in 2022. This was a significant decrease in emissions compared to the previous year and was primarily due to the shutdown of the ATR reactor. These airborne releases of radionuclides are reported to comply with regulatory requirements and are considered in the design and conduct of INL Site environmental surveillance activities.

The INL Site environmental surveillance programs, conducted by INL and ICP contractors emphasize the measurement of airborne radionuclides because air transport is considered the major potential pathway from INL Site releases to human receptors. During 2022, the INL contractor monitored ambient air at 34 locations (18 onsite, 8 boundary, and 8 offsite). The ICP contractor focused on ambient air monitoring of waste management facilities, namely INTEC and RWMC.

Air particulate samples were collected weekly by the INL contractor and biweekly by the ICP contractor. These samples were initially analyzed for gross alpha and gross beta activity. The particulate samples were then combined into monthly (ICP contractor) or quarterly (INL contractor) composite samples and were analyzed for gamma-emitting radionuclides such as  $^{137}\text{Cs}$ . Particulate filters were also composited quarterly by INL and ICP contractors. INL contractor analyzed for specific alpha- and beta-emitting radionuclides, specifically strontium-90 ( $^{90}\text{Sr}$ ), plutonium-238, plutonium-239/240, americium-241, uranium-233/234, and uranium-238. Charcoal cartridges were also collected weekly by the INL contractor and analyzed for radioiodine.

All radionuclide concentrations in ambient air samples were below DOE radiation protection standards for air. In addition, gross alpha and gross beta concentrations were analyzed statistically, and there were no differences between samples collected onsite, boundary, and offsite locations. Trends in the data appear to be seasonal in nature and do not demonstrate any INL Site influence. This indicates that INL Site airborne effluents were not measurable in environmental air samples.

The INL contractor collected atmospheric moisture samples at three stations onsite, three stations offsite, and two boundary stations in 2022. Precipitation was collected at one location onsite, two boundary locations, and one offsite location. The samples were all analyzed for tritium. The results were within measurements made historically and below DOE Derived Concentration Standards. Tritium measured in these samples is most likely the result of natural production in the atmosphere and not the result of INL Site effluent releases.

## ENVIRONMENTAL MONITORING OF GROUNDWATER, DRINKING, AND SURFACE WATER

The INL and ICP contractors monitor liquid effluents (wastewater), drinking water, groundwater, and storm water runoff at the INL Site, primarily for nonradioactive constituents, for compliance with applicable laws and regulations, DOE orders, and other requirements. Wastewater is typically discharged from INL Site facilities to infiltration ponds or to evaporation ponds. Wastewater effluent discharges occur at percolation ponds southwest of INTEC, a cold waste pond at the ATR Complex, and an industrial waste pond at MFC. DOE-ID complies with the state of Idaho groundwater quality, wastewater, and reuse rules for these effluents through reuse permits, which provide for monitoring of the wastewater and, in some instances, groundwater in the area. During 2022, liquid effluent and groundwater monitoring were conducted in support of reuse permit requirements. An annual site performance report for each permitted reuse facility was prepared and submitted to the Idaho Department of Environmental Quality. No permit limits were exceeded.

Additional liquid effluent monitoring was performed at the ATR Complex Cold Waste Pond, INTEC, and MFC Industrial Waste Pond to comply with environmental protection objectives of DOE orders. Most results were within historical measurements. All radioactive parameters were below health-based contaminant levels.

Drinking water parameters are regulated by the state of Idaho under the authority of the Safe Drinking Water Act. The INL and ICP contractors monitored 11 drinking water systems at the INL Site in 2022. (The NRF contractor monitors an



additional drinking water system, the results of which are reported separately by NRF.) Results were below limits for all relevant drinking water standards.

Surface water flows off the Subsurface Disposal Area (SDA) following periods of heavy precipitation or rapid snowmelt. During these times, water may be pumped out of the SDA retention basin into a drainage canal, potentially carrying radionuclides originating from radioactive waste or contaminated surface soil off the SDA. Surface water is collected when it is available. Americium-241, plutonium-239/240, and <sup>90</sup>Sr were detected in 2022 samples collected from the SDA Lift Station. The detected concentrations are well below standards established by DOE for radiation protection of the public and the environment.

## ENVIRONMENTAL MONITORING OF THE EASTERN SNAKE RIVER PLAIN AQUIFER

The eastern Snake River Plain Aquifer is perhaps the single-most important aquifer in Idaho. Composed of layered basalt lava flows and some sediment, it covers an area of approximately 27,972 km<sup>2</sup> (10,800 square miles). The highly productive aquifer has been declared a sole source aquifer by the EPA due to the nearly complete reliance on the aquifer for drinking water supplies in the area.

The USGS began monitoring the groundwater below the INL Site in 1949. Currently, the USGS performs groundwater monitoring, analyses, and studies of the eastern Snake River Plain Aquifer under and adjacent to the INL Site. These activities use an extensive network of strategically placed monitoring wells on and around the INL Site. In 2022, the USGS continued to monitor localized areas of chemical and radiochemical contamination beneath the INL Site produced by past waste disposal practices, in particular, the direct injection of wastewater into the aquifer at INTEC. Results for monitoring wells sampled within the plumes show nearly all wells had decreasing trends of tritium and <sup>90</sup>Sr concentrations over time.

Volatile organic compounds (VOCs) are present in water from the eastern Snake River Plain Aquifer because of historical waste disposal practices at the INL Site. Several purgeable VOCs were detected by USGS in 26 groundwater monitoring wells and one perched well sampled at the INL Site in 2022. Most concentrations of the 61 analyzed compounds were either below the laboratory reporting levels or their respective primary contaminant standards. Trend test results for tetrachloromethane concentrations in water from the RWMC production well show a decreasing trend in that well since 2005. The more recent decreasing trend indicates that remediation efforts designed to reduce VOC movement to the aquifer are having a positive effect. Concentrations of tetrachloromethane from USGS-87 and USGS-120, south of RWMC, have had an increasing trend since 1987; however, concentrations have decreased through time at USGS-88. Trichloroethylene was detected above the maximum contaminant level (MCL) in one well sampled by the USGS at TAN, which was expected as there is a known groundwater plume at this location as well as one perched well.

Groundwater surveillance monitoring continued for the Comprehensive Environmental Response, Compensation, and Liability Act WAGs onsite in 2022. At TAN (WAG 1), groundwater monitoring continues to monitor the progress of remediation of the plume of trichloroethylene and to monitor <sup>90</sup>Sr and <sup>137</sup>Cs. Remedial action consists of three components: in situ bioremediation, pump and treat, and monitored natural attenuation. Strontium-90 and <sup>137</sup>Cs were present in wells in the source area at levels higher than those prior to starting in situ bioremediation. The elevated concentrations of these radionuclides are due to in situ bioremediation activities. The radionuclide concentrations will continue to be evaluated to determine if they will meet remedial action objectives by 2095.

Groundwater samples were collected from six aquifer wells in the vicinity of ATR Complex (WAG 2) during 2022 and were analyzed for <sup>90</sup>Sr, cobalt-60 (<sup>60</sup>Co), tritium, and chromium. Chromium and tritium were the only analytes detected; however, neither of the concentrations were above their respective drinking water MCL established by the EPA.





Groundwater samples were collected from 13 aquifer monitoring wells at and near INTEC (WAG 3) during 2022 and analyzed for a suite of radionuclides and inorganic constituents. Strontium-90, technetium-99, and nitrate exceeded their respective drinking water MCLs in one or more aquifer monitoring wells at or near INTEC, with  $^{90}\text{Sr}$  exceeding its MCL by the greatest margin in a well south (downgradient) of the former INTEC injection well. All other well locations showed  $^{90}\text{Sr}$  levels similar to or slightly lower than those reported in previous samples.

Monitoring groundwater at CFA (WAG 4) consists of CFA landfill monitoring and monitoring of a nitrate plume south of the CFA. Wells at the landfill were monitored in 2022 for metals (filtered), VOCs, and anions (e.g., nitrate, chloride, fluoride, sulfate). No CFA landfill monitoring samples exceeded a MCL or secondary maximum contaminant level (SMCL). Nitrate continued to exceed the EPA MCL in one well in the plume south of the CFA in 2022; however, the data shows a downward trend since 2006.

Groundwater samples were collected from monitoring wells near and downgradient of the RWMC (WAG 7) in May 2022, which were analyzed for radionuclides, inorganic constituents, and VOCs. Carbon tetrachloride was detected slightly above the MCL (5 ug/L) in one regular sample and its field duplicate from Well M15S. Carbon tetrachloride concentrations in all other well locations were below the MCL and consistent with historical detections in May 2022.

Wells at MFC (as part of WAG 9, and the MFC Industrial Waste Pond Reuse Permit) were sampled for radionuclides, metals, and other water quality parameters in the spring and fall of 2022. Overall, the results were not above the primary constituent standard/secondary constituent standard and show no evidence of impacts from MFC activities.

Wells along the southern INL Site boundary (as part of WAG 10) are sampled every two years. Groundwater samples were not collected in 2022. WAG 10 monitoring wells will be sampled in 2023.

Groundwater is monitored at the Remote-Handled Low-Level Waste Facility for gross alpha, gross beta, carbon-14 ( $^{14}\text{C}$ ), iodine-129, technetium-99, and tritium. Samples were collected from three monitoring wells in the spring and fall of 2022. The results were not above the primary constituent standard/secondary constituent standard and show no discernable impacts to the aquifer from Remote-Handled Low-Level Waste Facility operations.

Drinking water and surface water samples were sampled downgradient of the INL Site and analyzed for gross alpha and beta activity and tritium. Tritium was not detected in any of these surface or drinking water samples. Gross alpha and beta results were within historical measurements and below the EPA's screening level. The data appear to show no discernable impacts from activities at the INL Site.

## MONITORING OF AGRICULTURAL PRODUCTS, WILDLIFE, SOIL, AND DIRECT RADIATION MEASUREMENTS

To help assess the impact of contaminants released to the environment by operations at the INL Site, agricultural products (e.g., milk, lettuce, alfalfa, grain, potatoes) and wildlife were sampled and analyzed for radionuclides in 2022. The agricultural products were collected onsite, offsite, and at INL boundary locations by the INL contractor.

Some human-made radionuclides were detected in agricultural products; however, measurements were consistent with those made historically.

No human-made radionuclides were detected in big game animal samples collected in 2022. Cobalt-60 and  $^{90}\text{Sr}$  were detected in tissues of waterfowl collected near the ATR Complex ponds, indicating that they accessed the contaminated ponds. Zinc-65 was detected in one waterfowl collected near TAN.

Cobalt-60,  $^{90}\text{Sr}$ , and  $^{137}\text{Cs}$  were detected in some composited bat samples, indicating that bats may have visited radioactive wastewater ponds such as those at the ATR Complex.

Direct radiation measurements made at onsite, offsite, and boundary locations were consistent with historical and natural background levels.



## NATURAL AND CULTURAL RESOURCES CONSERVATION AND MONITORING

Natural resources conservation, monitoring, and land stewardship activities onsite are organized in four categories: (1) frequently evaluating the regulatory rankings, distribution, and populations for special status species; (2) planning and implementing conservation efforts for high priority natural resources; (3) ongoing monitoring and research to provide baseline and trend data for specific taxa and broader ecological communities; and (4) conducting land stewardship activities to minimize impacts to natural resources and restore ecological condition, where appropriate.

The INL Site provides breeding and foraging habitat for a variety of animal species, including 24 species of birds and 12 species of mammals that are of elevated conservation concern by state or federal agencies. There are also currently 20 special status plant species that have been documented to occur onsite. Many of those species are rare and occur very infrequently within their optimal habitats. While several animals and plants listed as threatened or endangered under the Endangered Species Act are present in Idaho, none are known to occur onsite.

For some species of elevated concern or with extensive populations and key habitats onsite, DOE-ID has developed conservation plans to protect species and the valuable ecosystems they inhabit. Conservation plans that are specific to or include the INL Site are the Candidate Conservation Agreement for Greater sage-grouse (*Centrocercus urophasianus*), the INL Site Bat Protection Plan, the Sagebrush Steppe Ecosystem Reserve, the Migratory Bird Conservation Plan and Avian Protection Planning documents, and the DOE Conservation Action Plan. Many of these plans include conservation measures, best management practices, monitoring programs, and annual reports to facilitate, evaluate, and communicate results of conservation efforts for species with high conservation priority.

Additional ecological monitoring has been conducted for more than 70 years onsite, with some studies dating back to the 1950s. The focus of this work is to better understand the INL Site's ecosystem and biota and to determine the impact on populations of these species from activities conducted at the INL Site. Natural resource monitoring activities include breeding bird surveys, midwinter raptor survey, long-term vegetation transects, and vegetation mapping. Furthermore, the INL Site was designated as a National Environmental Research Park in 1975 and serves as an outdoor laboratory for environmental scientists to study Idaho's native plants and wildlife in an intact and relatively undisturbed ecosystem. Ongoing National Environmental Research Park activities range from characterizing sagebrush steppe ecohydrology to identifying high quality foodscape for sage-grouse.

Land stewardship involves managing ecosystems onsite through planning, assessment, restoration, and rehabilitation activities. Areas where DOE-ID is actively employing land stewardship activities include wildland fire protection planning, management, and recovery; restoration and revegetation; weed management; and ecological support for the National Environmental Policy Act.

The INL Cultural Resource Management Office coordinates cultural resource-related activities at the INL Site and implements the INL Cultural Resource Management Plan (DOE-ID 2016) with oversight by DOE-ID's Cultural Resource Coordinator. Cultural resource identification and evaluation studies in fiscal year 2022 included (1) archaeological field surveys, (2) cultural resource monitoring and site record updates related to INL Site project activities and research, and (3) comprehensive evaluations of pre-1980 built environment resources. Additionally, the Cultural Resource Management Office supports the DOE-ID with their government-to-government consultation and meaningful collaboration with members of the Shoshone-Bannock Tribes to include the Fort Hall Business Council, the Language and Cultural Committee, and the Heritage Tribal Office (known as the HeTO), as well as other public stakeholders.

## USGS RESEARCH

The USGS INL Project Office drills and maintains research wells that provide information about subsurface water, rock and sediment, and contaminant movement in the eastern Snake River Plain Aquifer at and near the INL Site. In 2022, the USGS published two research reports and one software release.



## QUALITY ASSURANCE

Quality assurance and quality control programs are maintained by contractors conducting environmental monitoring and by laboratories performing environmental analyses to help provide confidence in the data and ensure data completeness. Programs involved in environmental monitoring developed quality assurance programs and documentation, which follow requirements and criteria established by DOE. Environmental monitoring programs implemented quality assurance program elements through quality assurance project plans developed for each contractor.

Adherence to procedures and quality assurance project plans was maintained during 2022. Data reported in this document were obtained from several commercial, university, government, and government contractor laboratories. To ensure quality results, these laboratories participated in several laboratory quality check programs. Quality issues that arose with laboratories used by INL and ICP contractors and USGS during 2022 were addressed with the laboratories and have been or are being resolved.



*Western skink*

# Helpful Information:



## What is Radiation?

Much of the Annual Site Environmental Report deals with radioactivity levels measured in environmental media such as air, water, soil, and plants. The following information is intended for individuals with little or no familiarity with radiological data or radiation dose. It presents terminology and concepts used in the Annual Site Environmental Report to aid the reader.


Matter is composed of atoms. Some atoms are energetically unstable and change to become more stable. During this transformation, unstable or radioactive atoms give off energy called radiation in the form of particles or electromagnetic waves. Generally, we refer to the various radioactive atoms as radionuclides. The radiation released by radionuclides has enough energy to eject electrons from other atoms it encounters. The resulting charged atoms or molecules are called ions, and the energetic radiation that produced the ions is called ionizing radiation. Ionizing radiation is referred to simply as radiation throughout this report. The most common types of radiation are alpha particles, beta particles, X-rays, and gamma-rays. X-rays and gamma-rays, just like visible light and radio waves, are packets of electromagnetic radiation. Collectively, packets of electromagnetic radiation are called photons. One may, for instance, speak of X-ray photons or gamma-ray photons.

**Alpha Particles.** An alpha particle is a helium nucleus without orbital electrons. It is composed of two protons and two neutrons and has a positive charge of two. Because alpha particles are relatively heavy and have a double charge, they cause intense tracks of ionization but have little penetrating ability, as observed in Figure HI-1. Alpha particles can be stopped by thin layers of materials, such as a sheet of paper or a piece of aluminum foil. Examples of alpha-emitting radionuclides include radioactive atoms of radon, uranium, plutonium, and americium.

**Beta Particles.** Beta particles are electrons that are ejected from unstable atoms during the transformation or decay process. Beta particles penetrate more than alpha particles but are less penetrating than X-rays or gamma-rays of equivalent energies. A piece of wood or a thin block of plastic can stop beta particles, as can be seen in Figure HI-1. The ability of beta particles to penetrate matter increases with energy. Examples of beta-emitting radionuclides include tritium ( $^3\text{H}$ ) and radioactive strontium.

**X-Rays and Gamma-Rays.** X-rays and gamma-rays are photons with very short wave-lengths compared to other electromagnetic waves such as visible light, heat rays, and radio waves. Gamma-rays and X-rays have identical properties, behavior, and effects but differ in their origin. Gamma-rays originate from an atomic nucleus, and X-rays originate from interactions with the electrons orbiting around atoms. All photons travel at the speed of light. Their energies, however, vary over a large range. The penetration of X-ray or gamma-ray photons depend on the energy of the photons as well as the thickness, density, and composition of the shielding material. Concrete is a common material used to shield people from gamma-rays and X-rays, as shown in Figure HI-1.

Examples of gamma-emitting radionuclides include radioactive atoms of iodine and cesium. X-rays may be produced by medical X-ray machines in a doctor's office.



**Figure HI-1. Comparison of penetrating ability of alpha, beta, and gamma radiation.**

## How are Radionuclides Designated?

Radionuclides are frequently expressed with a one or two letter abbreviation for the element and a superscript to the left of the symbol that identifies the atomic weight of the isotope. The atomic weight is the number of protons and neutrons in the nucleus of the atom. Most radionuclide symbols used in this report are shown in Table HI-1. This table also shows the half-life of each radionuclide. Half-life refers to the time in which one-half of the atoms of a radioactive sample transforms or decays in the quest to achieve a more energetically stable nucleus. Most radionuclides do not decay directly to a stable element, but rather they undergo a series of decays until a stable element is reached. This series of decays is called a decay chain.

## How are Radioactivity and Radionuclides Detected?

Environmental samples of air, water, soil, and plants are collected in the field and then prepared and analyzed for radioactivity in a laboratory. A prepared sample is placed in a radiation counting system with a detector that converts the ionization produced by the radiation into electrical signals or pulses. The number of electrical pulses recorded over a unit of time is called a count rate. The count rate is proportional to the amount of radioactivity in the sample.

Air and water samples are often analyzed to determine the total amount of alpha-emitting and beta-emitting radioactivity present. This is referred to as a gross measurement because the radiation from all alpha-emitting and beta-emitting radionuclides in the sample is quantified. Such sample analyses measure both human-generated and naturally occurring radioactive material. Gross alpha and beta analyses are generally considered screening measurements since specific radionuclides are not identified. The amount of gross alpha-emitting and beta-emitting radioactivity in air samples is frequently measured to screen for the potential presence of man-made radionuclides. If the results are higher than normal, sources other than background radionuclides may be suspected, and other laboratory techniques may be used to identify the specific radionuclides in the sample. Gross alpha and beta activity also can be examined over time and between locations to detect trends.

The low penetration ability of alpha-emitting particles makes detection by any instrument difficult. Identifying specific alpha-emitting radionuclides typically involves chemical separations in the laboratory to purify the sample prior to analysis with an alpha detection instrument. Radiochemical analysis is very time-consuming and expensive.

Beta particles are easily detected by several types of instruments, including the common Geiger-Mueller counter. However, detection of specific beta-emitting radionuclides, such as  $^3\text{H}$  and  $^{90}\text{Sr}$ , requires chemical separation first.



**Table HI-1. Radionuclides and their half-lives.**

SYMBOL	RADIONUCLIDE	HALF-LIFE <sup>a,b</sup>	SYMBOL	RADIONUCLIDE	HALF-LIFE <sup>a,b</sup>
<sup>241</sup> Am	Americium-241	432.2 yr	<sup>54</sup> Mn	Manganese-54	312.12 d
<sup>243</sup> Am	Americium-243	7,370 yr	<sup>59</sup> Ni	Nickel-59	1.01 × 10 <sup>5</sup> yr
<sup>125</sup> Sb	Antimony-125	2.75856 yr	<sup>63</sup> Ni	Nickel-63	100.1 yr
<sup>41</sup> Ar	Argon-41	109.61 min	<sup>238</sup> Pu	Plutonium-238	87.7 yr
<sup>137m</sup> Ba	Barium-137m	2.552 min	<sup>239</sup> Pu	Plutonium-239	2.411 × 10 <sup>4</sup> yr
<sup>140</sup> Ba	Barium-140	12.752 d	<sup>240</sup> Pu	Plutonium-240	6,564 yr
<sup>7</sup> Be	Beryllium-7	53.22 d	<sup>241</sup> Pu	Plutonium-241	14.35 yr
<sup>14</sup> C	Carbon-14	5,700 yr	<sup>242</sup> Pu	Plutonium-242	3.75 × 10 <sup>5</sup> yr
<sup>141</sup> Ce	Cerium-141	32.508 d	<sup>40</sup> K	Potassium-40	1.251 × 10 <sup>9</sup> yr
<sup>144</sup> Ce	Cerium-144	284.91 d	<sup>226</sup> Ra	Radium-226	1,600 yr
<sup>134</sup> Cs	Cesium-134	2.0648 yr	<sup>228</sup> Ra	Radium-228	5.75 yr
<sup>137</sup> Cs	Cesium-137	30.1671 yr	<sup>220</sup> Rn	Radon-220	55.6 s
<sup>51</sup> Cr	Chromium-51	27.7025 d	<sup>222</sup> Rn	Radon-222	3.8235 d
<sup>60</sup> Co	Cobalt-60	5.2713 yr	<sup>103</sup> Ru	Ruthenium-103	39.26 d
<sup>152</sup> Eu	Europium-152	13.537 yr	<sup>106</sup> Ru	Ruthenium-106	373.59 d
<sup>154</sup> Eu	Europium-154	8.593 yr	<sup>90</sup> Sr	Strontium-90	28.79 yr
<sup>3</sup> H	Tritium	12.32 yr	<sup>99</sup> Tc	Technetium-99	2.111 × 10 <sup>5</sup> yr
<sup>129</sup> I	Iodine-129	1.57 × 10 <sup>7</sup> yr	<sup>232</sup> Th	Thorium-232	1.405 × 10 <sup>10</sup> yr
<sup>131</sup> I	Iodine-131	8.0207 d	<sup>233</sup> U	Uranium-233	1.592 × 10 <sup>5</sup> yr
<sup>55</sup> Fe	Iron-55	2.737 yr	<sup>234</sup> U	Uranium-234	2.455 × 10 <sup>5</sup> yr
<sup>59</sup> Fe	Iron-59	44.495 d	<sup>235</sup> U	Uranium-235	7.04 × 10 <sup>8</sup> yr
<sup>85</sup> Kr	Krypton-85	10.756 yr	<sup>238</sup> U	Uranium-238	4.468 × 10 <sup>9</sup> yr
<sup>87</sup> Kr	Krypton-87	76.3 min	<sup>90</sup> Y	Yttrium-90	64.1 hr
<sup>88</sup> Kr	Krypton-88	2.84 hr	<sup>65</sup> Zn	Zinc-65	244.06 d
<sup>212</sup> Pb	Lead-212	10.64 hr	<sup>95</sup> Zr	Zirconium-95	64.032 d

a. From ICRP Publication 107 (ICRP 2008).

b. d = days; hr = hours; min = minutes; s = seconds; yr = years.

The high-energy photons from gamma-emitting radionuclides are relatively easy to detect. Because the photons from each gamma-emitting radionuclide have a characteristic energy, gamma emitters can be simply identified in the laboratory with only minimal sample preparation prior to analysis. Gamma-emitting radionuclides, such as <sup>137</sup>Cs, can even be measured in soil by field detectors called in situ detectors.

Gamma radiation originating from naturally occurring radionuclides in soil and rocks on the earth's surface is a primary contributor to the background external radiation exposure measured in the air. Cosmic radiation from outer space is another contributor to the external radiation background. External radiation is easily measured with devices known as environmental dosimeters.



## How are Results Reported?

**Scientific Notation.** Concentrations of radionuclides detected in the environment are typically quite small. Scientific notation is used to express numbers that are very small or very large. A very small number may be expressed with a negative exponent, for example,  $1.3 \times 10^{-6}$  (or 1.3E-06). To convert this number to its decimal form, the decimal point is moved left by the number of places equal to the exponent (in this case, six). The number  $1.3 \times 10^{-6}$  may also be expressed as 0.0000013. When considering large numbers with a positive exponent, such as  $1.0 \times 10^6$ , the decimal point is moved to the right by the number of places equal to the exponent. In this case,  $1.0 \times 10^6$  represents one million and may also be written as 1,000,000.

**Unit Prefixes.** Units for very small and very large numbers are often expressed with a prefix. One common example is the prefix kilo (abbreviated k), which means 1,000 of a given unit. One kilometer, therefore, equals 1,000 meters. Table HI-2 defines the values of commonly used prefixes.

**Table HI-2. Multiples of units.**

MULTIPLE	DECIMAL EQUIVALENT	PREFIX	SYMBOL
$10^6$	1,000,000	mega-	M
$10^3$	1,000	kilo-	k
$10^2$	100	hecto-	h
10	10	deka-	da
$10^{-1}$	0.1	deci-	d
$10^{-2}$	0.01	centi-	c
$10^{-3}$	0.001	milli-	m
$10^{-6}$	0.000001	micro-	$\mu$
$10^{-9}$	0.000000001	nano-	n
$10^{-12}$	0.000000000001	pico-	p
$10^{-15}$	0.000000000000001	femto-	f
$10^{-18}$	0.000000000000000001	atto-	a

**Units of Radioactivity.** The basic unit of radioactivity used in this report is the curie (abbreviated Ci), which is based on the disintegration rate occurring in 1 gram of the radionuclide radium-226 ( $^{226}\text{Ra}$ ) that is 37 billion ( $3.7 \times 10^{10}$ ) disintegrations per second (becquerels). For any other radionuclide, 1 Ci is the amount of the radionuclide that produces this same decay rate.

**Units of Exposure and Dose (Table HI-3).** Exposure, or the amount of ionization produced by gamma or X-ray radiation in the air, is measured in terms of the roentgen ®. Dose is a general term to express how much radiation energy is deposited into something. The energy deposited can be expressed in terms of absorbed, equivalent, and effective dose. The term rad, which is short for radiation absorbed dose, is a measure of the energy absorbed in an organ or tissue. The equivalent dose, which considers the effect of different types of radiation on tissues and is therefore the potential for biological effects, is expressed as the R equivalent man or rem. Radiation exposures to the human body, whether from external or internal sources, can involve all or a portion of the body. To enable radiation protection specialists to express partial-body exposures (and the accompanying doses) to portions of the body in terms of an equal dose to the whole body, the concept of effective dose was developed.





**Table HI-3. Names and symbols for units of radioactivity and radiological dose used in this report.**

SYMBOL	NAME
Bq	Becquerel
Ci	Curie (37,000,000,000 Bq)
mCi	Millicurie ( $1 \times 10^{-3}$ Ci)
$\mu$ Ci	Microcurie ( $1 \times 10^{-6}$ Ci)
mrad	Millirad ( $1 \times 10^{-3}$ rad)
mrem	Millirem ( $1 \times 10^{-3}$ rem)
R	Roentgen
mR	Milliroentgen ( $1 \times 10^{-3}$ R)
$\mu$ R	Microroentgen ( $1 \times 10^{-6}$ R)
Sv	Sievert (100 rem)
mSv	Millisievert (100 mrem)
$\mu$ Sv	Microsievert (0.1 mrem)

The Syst me International (SI) is the official system of measurement used internationally to express units of radioactivity and radiation dose. The basic SI unit of radioactivity is the Becquerel (Bq), which is equivalent to one nuclear disintegration per second. The number of curies must be multiplied by  $3.7 \times 10^{10}$  to obtain the equivalent number of becquerels. The concept of dose may also be expressed using the SI units, Gray (Gy) for absorbed dose ( $1 \text{ Gy} = 100 \text{ rad}$ ) and sievert (Sv) for effective dose ( $1 \text{ Sv} = 100 \text{ rem}$ ).

**Concentrations of Radioactivity in Environmental Sample Media.** Table HI-4 shows the units used to identify the concentration of radioactivity in various sample media.

There is always uncertainty associated with the measurement of radioactivity in environmental samples. This is mainly because radioactive decay events are inherently random. Thus, when a radioactive sample is counted again and again for the same length of time, the results will differ slightly, but most of the results will be close to the true value of the activity of the radioactive material in the sample. Statistical methods are used to estimate the true value of a single measurement and the associated uncertainty of the measurement. The uncertainty of a measurement is reported by following the result with an uncertainty value that is preceded by the plus or minus symbol,  $\pm$  (e.g.,  $10 \pm 2 \text{ pCi/L}$ ). The uncertainty is often referred to as sigma (or  $\sigma$ ). For concentrations of greater than or equal to three times the uncertainty, there is 99% probability that the radionuclide was detected in a sample. For example, if a radionuclide is reported for a sample at a concentration of  $10 \pm 2 \text{ pCi/L}$ , then the radionuclide is considered to be detected in that sample because 10 is greater than  $3 \times 2$ , or 6. On the other hand, if the reported concentration of a radionuclide (e.g.,  $10 \pm 6 \text{ pCi/L}$ ) is smaller than three times its associated uncertainty, then the sample probably does not contain that radionuclide (i.e., 10 is less than  $3 \times 6$ , or 18). Such low concentrations are considered to be undetected by the method or instrumentation used.

**Table HI-4. Units of radioactivity.**

MEDIA	UNIT
Air	Microcuries per milliliter ( $\mu$ Ci/mL)
Liquid, such as water and milk	Picocuries per liter (pCi/L)
Soil and agricultural products	Picocuries per gram (pCi/kg) dry weight
Annual human radiation exposure, measured by environmental dosimeters	Milliroentgens (mR) or millirem (mrem), after being multiplied by an appropriate dose equivalent conversion factor




**Mean, Median, Maximum, and Minimum Values.** Descriptive statistics are often used to express the patterns and distribution of a group of results. The most common descriptive statistics used in this report are the mean, median, minimum, and maximum values. Mean and median values measure the central tendency of the data. The mean is calculated by adding up all the values in a set of data and then dividing that sum by the number of values in the dataset. The median is the middle value in a group of measurements. When the data are arranged from largest (maximum) to smallest (minimum), the result in the exact center of an odd number of results is the median. If there is an even number of results, the median is the average of the two central values. The maximum and minimum results represent the range of the measurements.

Statistical analysis of many of the air data reported in this annual report indicate that the median is a more appropriate representation of the central tendency of those results. For this reason, some of the figures present the median value of a data group. For example, Figure HI-2 is a box plot showing the minimum, maximum, and median of a set of air measurements.


## How are Data Represented Graphically?

Charts and graphs often are used to compare data and to visualize patterns, such as trends over time. Four kinds of graphics are used in this report to represent data: pie charts, column graphs, line plots, and contour lines.

A **pie chart** is used in this report to illustrate fractions of a whole. For example, Figure HI-3 shows the approximate contribution to dose that a typical person might receive while living in southeast Idaho. The percentages are derived from the table in the lower left-hand corner of the figure. The medical, consumer, and occupational/industrial portions are from the National Council on Radiation Protection and Measurements Report No. 160 (NCRP 2009). The contribution from background (e.g., natural radiation, mostly radon) is estimated in Table 7-7 of this report.



**Figure HI-2. A graphical representation of minimum, median, and maximum results with a box plot.** The 25th and 75th percentiles are the values such that 75% of the measurements in the dataset are greater than the 25th percentile, and 75% of the measurements are less than the 75th percentile.



**Figure HI-3. Data presented using a pie chart.**

A **column or bar chart** can show data changes over a period of time or illustrate comparisons among items. Figure HI-4 illustrates the maximum dose (mrem) calculated for the maximally exposed individual from 2013 through 2022. The maximally exposed individual is a hypothetical member of the public who is exposed to radionuclides from airborne releases through various environmental pathways and the media through which the radionuclides are transported (i.e., air, water, and food). The chart shows the general trend of the dose over time.






Figure HI-4. Data plotted using a column chart.

A **plot chart** can be useful to visualize differences in results over time. Figure HI-5 shows the <sup>90</sup>Sr measurements in three wells collected by USGS for 21 years (2002–2022). The results are plotted by year.



**Figure HI-5. Data plotted using a linear plot.**

**Contour lines** are sometimes drawn on a map to discern patterns over a geographical area. For example, Figure HI-6 shows the distribution of  $^{90}\text{Sr}$  in groundwater around INTEC. Each contour line, or isopleth, represents a specific concentration of the radionuclide in groundwater. It was estimated from measurements of samples collected from wells around INTEC. Each contour line separates areas that have concentrations above the contour line value from those that have concentrations below that value. The figure shows the highest concentration gradient near INTEC and the lowest farther away. It reflects the movement of the radionuclide in groundwater from INTEC where it was injected into the aquifer in the past.



**Figure HI-6. Data plotted using contour lines.** Each contour line drawn on this map connects points of equal <sup>90</sup>Sr concentration in water samples collected at the same depth from wells onsite.



## How Are Results Interpreted?

To better understand data, results are compared in one or more ways, including the following:

- Comparison of results collected at different locations. For example, measurements made at INL Site locations are compared with those made at locations near the boundary of the INL Site and offsite to find differences that may indicate an impact (Figure HI-2).
- Trends over time or space. Data collected during the year can be compared with data collected at the same location or locations during previous years to see if concentrations are increasing, decreasing, or remaining the same with time. See, for example, Figure HI-4, which shows a general decrease in dose from 2013 to 2018, followed by a slight increase in 2019. Figure HI-6 illustrates a clear spatial pattern of radionuclide concentrations in groundwater decreasing with distance from the source.
- Comparison with background measurements. Humans are now, and always have been, continuously exposed to ionizing radiation from natural background sources. Background sources include natural radiation and radioactivity as well as radionuclides from human activities. These sources are discussed in the following section.

## What Is Background Radiation?

Radioactivity from natural and fallout sources is detectable as background in all environmental media. Natural sources of radiation include (1) radiation of extraterrestrial origin (called cosmic rays), (2) radionuclides produced in the atmosphere by cosmic ray interaction with matter (called cosmogenic radionuclides), and (3) radionuclides present at the time of the formation of the earth (called primordial radionuclides). Radiation that has resulted from the activities of modern man is primarily fallout from past atmospheric testing of nuclear weapons. One of the challenges to environmental monitoring on and around the INL Site is to distinguish between what may have been released from the INL Site and what is already present in background from natural and fallout sources. These sources are discussed in more detail below.

Natural radiation and radioactivity in the environment, which is natural background, represent a major source of human radiation exposure (NCRP 1987, 2009). For this reason, natural radiation frequently is used as a standard of comparison for exposure to various human-generated sources of ionizing radiation. An individual living in southeast Idaho was estimated, in 2022, to receive an average dose of about 384 mrem/yr (3.8 mSv/yr) from natural background sources of radiation on earth, as observed in Figure HI-7. These sources include cosmic radiation and naturally occurring radionuclides.

Cosmic radiation is radiation that constantly bathes the earth in extraterrestrial sources. The atmosphere around the earth absorbs some of the cosmic radiation, so doses are lowest at sea level and increase sharply with altitude. Cosmic radiation is estimated using data in NCRP (2009) to produce a dose of about 57 mrem/yr (0.57 mSv/yr) to a typical individual living in southeast Idaho (Figure HI-7). Cosmic radiation also produces cosmogenic radionuclides, which are found naturally in all environmental media and are discussed in more detail below.

Naturally occurring radionuclides are of two general kinds: cosmogenic and primordial. Cosmogenic radionuclides are produced by the interaction of cosmic radiation within the atmosphere or in the earth. Cosmic rays have high enough energies to blast apart atoms in the earth's atmosphere. The result is the continuous production of radionuclides, such as  $^3\text{H}$ , beryllium-7, sodium-22, and  $^{14}\text{C}$ . Cosmogenic radionuclides, particularly  $^3\text{H}$  and  $^{14}\text{C}$ , have been measured in humans, animals, plants, soil, polar ice, surface rocks, sediments, the ocean floor, and the atmosphere. Concentrations are generally higher at mid-latitudes than at low- or high-latitudes. Cosmogenic radionuclides contribute only about 1 mrem/yr to the total average dose, mostly from  $^{14}\text{C}$ , that might be received by an adult living in the U.S. (NCRP 2009). Tritium and beryllium-7 are routinely detected in environmental samples collected by environmental monitoring programs on and around the INL Site, as observed in Figure HI-5, but these contribute little to the dose that might be received from natural background sources.




Figure HI-7. Calculated doses (mrem per year) from natural background sources for an average individual living in southeast Idaho (2022).

Table HI-5. Naturally occurring radionuclides that have been detected in environmental media collected on and around the INL Site.

RADIONUCLIDE	HALF-LIFE	HOW PRODUCED?	DETECTED OR MEASURED IN:
Beryllium-7 ( <sup>7</sup> Be)	53.22 da	Cosmic rays	Rain, air
Potassium-40 ( <sup>40</sup> K)	1.2516 × 10 <sup>9</sup> yr	Primordial	Water, air, soil, plants, animals
Radium-226 ( <sup>226</sup> Ra)	1,600 yr	<sup>238</sup> U progeny	Water
Thorium-232 ( <sup>232</sup> Th)	1.405 × 10 <sup>10</sup> yr	Primordial	Soil
Tritium ( <sup>3</sup> H)	12.32 yr	Cosmic rays	Water, rain, air moisture
Uranium-234 ( <sup>234</sup> U)	2.455 × 10 <sup>5</sup> yr	<sup>238</sup> U progeny	Water, air, soil
Uranium-238 ( <sup>238</sup> U)	4.468 × 10 <sup>9</sup> yr	Primordial	Water, air, soil





Primordial radionuclides are those that were present when the earth was formed. The primordial radionuclides detected today are billions of years old. The radiation dose to a person from primordial radionuclides comes from internally deposited radioactivity, inhaled radioactivity, and external radioactivity in soils and building materials. Three of the primordial radionuclides—potassium-40, uranium-238 ( $^{238}\text{U}$ ), and thorium-232 ( $^{232}\text{Th}$ )—are responsible for most of the dose received by people from natural background radioactivity. They have been detected in environmental samples collected on and around the INL Site (Table HI-5). The external dose to an adult living in southeast Idaho from terrestrial natural background radiation exposure (73 mrem/yr or 0.73 mSv/yr) has been estimated using concentrations of potassium-40,  $^{238}\text{U}$ , and  $^{232}\text{Th}$  measured in soil samples collected from areas surrounding the INL Site from 1976 through 1993. This number varies slightly from year to year based on the amount of snow cover. Uranium-238 and  $^{232}\text{Th}$  are also estimated to contribute 13 mrem/yr (0.13 mSv/yr) to an average adult through ingestion (NCRP 2009).

Potassium-40 is abundant and measured in living and nonliving matter. It is found in human tissue and is a significant source of internal dose to the human body (approximately 15 mrem/yr [0.15 mSv/yr] according to NCRP [2009]). Rubidium-87, another primordial radionuclide, contributes a small amount (< 1 mrem/yr) to the internal dose received by people but is not typically measured in INL Site samples.

Uranium-238 and  $^{232}\text{Th}$  initiate a decay chain of radionuclides. A radioactive decay chain starts with one type of radioactive atom called the parent that decays and changes into another type of radioactive atom called a progeny radionuclide. This system repeats, involving several different radionuclides. The parent radionuclide of the uranium decay chain is  $^{238}\text{U}$ . The most familiar element in the uranium series is radon, specifically radon-222. This is a gas that can accumulate in buildings. Radon and its progeny are responsible for most of the inhalation dose (e.g., an average of 200 mrem/yr [2.0 mSv/yr] nationwide) produced by naturally occurring radionuclides, as shown in Figure HI-7.

The parent radionuclide of the thorium series is  $^{232}\text{Th}$ . Another isotope of radon, called thoron, occurs in the thorium decay chain of radioactive atoms. Uranium-238,  $^{232}\text{Th}$ , and their progeny are often detected in environmental samples (Table HI-5).

**Global Fallout.** The U.S., the Union of Soviet Socialist Republics, and China tested nuclear weapons in the Earth's atmosphere in the 1950s and 1960s. This testing resulted in the release of radionuclides into the upper atmosphere, and such a release is referred to as fallout from weapons testing. Concerns over worldwide fallout rates eventually led to the Partial Test Ban Treaty in 1963, which limited signatories to underground testing. Not all countries stopped atmospheric testing with the treaty. France continued atmospheric testing until 1974, and China continued until 1980. Additional fallout, but to a substantially smaller extent, was produced by the Chernobyl and Fukushima nuclear accidents in 1986 and 2011, respectively.

Most of the radionuclides associated with nuclear weapons testing and the Chernobyl and Fukushima accidents have decayed and are no longer detected in environmental samples. Radionuclides that are currently detected in the environment and typically associated with global fallout include  $^{90}\text{Sr}$  and  $^{137}\text{Cs}$ . Strontium-90, a beta-emitter with a 29-year half-life, is important because it is chemically similar to calcium and tends to accumulate in bone tissues. Cesium-137, which has a 30-year half-life, is chemically similar to potassium and accumulates rather uniformly in muscle tissue throughout the body.

The deposition of these radionuclides on the earth's surface varies by latitude, with most occurring in the northern hemisphere at approximately 40 degrees. Variation within latitudinal belts is a function primarily of precipitation, topography, and wind patterns. The dose produced by global fallout from nuclear weapons testing has decreased steadily since 1970. The annual dose rate from fallout was estimated in 1987 to be less than 1 mrem (0.01 mSv) (NCRP 1987). It has been nearly 34 years since that estimate, so the current dose is assumed to be even lower.



## What are the Risks of Exposure to Low Levels of Radiation?

Radiation protection standards for the public have been established by state and federal agencies based mainly on recommendations of the International Commission on Radiological Protection and the National Council on Radiation Protection and Measurements. The International Commission on Radiological Protection is an association of scientists from many countries, including the U.S. The National Council on Radiation Protection and Measurements is a nonprofit corporation chartered by Congress. Through radiation protection standards, exposure of members of the general public to radiation is controlled so that risks are small enough to be considered insignificant compared to the risks undertaken during other activities deemed normal and acceptable in modern life.

A large amount of data exists concerning the effects of acute delivery (all at once) of high doses of radiation, especially in the range of 50–400 rem (0.5 to 4.0 Sv). Most of this information was gathered from the Japanese atomic bombing survivors and patients who were treated with substantial doses of X-rays. Conversely, information is limited, and therefore, it is difficult to estimate risks associated with low-level exposure. Risk can be defined in general as the probability (chance) of injury, illness, or death resulting from some activity. Low-dose effects are those that might be caused by doses of less than 20 rem (0.2 Sv), whether delivered acutely or spread out over a period as long as a year (Taylor 1996). Most of the radiation exposures that humans receive are very close to background levels. Moreover, many sources emit radiation that is well below natural background levels. This makes it extremely difficult to isolate its effects. For this reason, government agencies make the conservative (cautious) assumption that any increase in radiation exposure is accompanied by an increased risk of health effects. Cancer is considered by most scientists to be the primary health effect from long-term exposure to low levels of radiation while each radionuclide represents a somewhat different health risk. A 2011 report by the EPA estimated a  $5.8 \times 10^{-2} \text{ Gy}^{-1}$  cancer mortality risk coefficient for uniform whole-body exposure throughout life at a constant dose rate. Given a 1 gray (100 rad) ionizing radiation lifetime exposure, this corresponds to 580 deaths, above normal cancer mortality rates, within an exposure group of 10,000 people. For low-linear energy transfer radiation (i.e., beta and gamma radiation) the dose equivalent in Sv (100 rem) is numerically equal to the absorbed dose in Gy (100 rad). Therefore, if each person in a group of 10,000 people is exposed to 1 rem (0.01 Sv) of ionizing radiation in small doses over a lifetime, we would expect around six people to die of cancer than would otherwise. For perspective, most people living on the eastern Snake River Plain receive over 381 mrem (3.8 mSv) every year from natural background sources of radiation.

DOE limits the dose to a member of the public from all sources and pathways to 100 mrem (1 mSv) and the dose from the air pathway to 10 mrem (0.1 mSv) (DOE O 458.1). The doses estimated to maximally exposed individuals from INL Site releases are typically well below 1 mrem per year.

## References

- 40 CFR 61, Subpart H, 2023, “National Emission Standards for Emissions of Radionuclides Other Than Radon from Department of Energy Facilities,” Code of Federal Regulations, Office of the Federal Register, National Archives and Records Administration, <https://ecfr.io/Title-40/sp40.10.61.h>.
- DOE O 231.1B, 2011, “Environment, Safety, and Health Reporting,” U.S. Department of Energy.
- DOE O 458.1, 2020, “Radiation Protection of the Public and the Environment,” U.S. Department of Energy.
- ICRP, 2008, Nuclear Decay Data for Dosimetric Calculations, ICRP Publication 107, International Commission on Radiological Protection.
- NCRP, 1987, “Exposure of the Population in the United States and Canada from Natural Background Radiation,” NCRP Report No. 94, National Council on Radiation Protection and Measurements.
- NCRP, 2009, “Ionizing Radiation Exposure of the Population of the United States,” NCRP Report No. 160, National Council on Radiation Protection and Measurements.
- Taylor, L. S., 1996, “What You Need to Know About Radiation,” edited by Joyce Davis, <https://sites.google.com/isu.edu/health-physics-radinf/l-s-taylor>.

# Acronyms:



AFV	alternative fuel vehicle	CWP	Cold Waste Pond
ALLWDF	active low-level waste disposal facility	D&D	decontamination and decommissioning
ARP	Accelerated Retrieval Project	DCS	Derived Concentration Standard
ATR	Advanced Test Reactor	DEQ	Department of Environmental Quality (state of Idaho)
BBS	breeding bird survey	DEQ-IOP	Department of Environmental Quality – INL Oversight Program
BCG	Biota Concentration Guide	DOE	U.S. Department of Energy
BEA	Battelle Energy Alliance, LLC	DOE-ICP	DOE Idaho Cleanup Project
BLM	Bureau of Land Management	DOE-ID	U.S. Department of Energy, Idaho Operations Office
BMP	best management practices	DOSEMM	dose multi-media
BRR	Biological Resource Review	DQO	data quality objective
C&D	construction and demolition	EAs	Environmental Assessments
CA	corrective action	EBR-I	Experimental Breeder Reactor-I
CAA	Clean Air Act	ECP	Environmental Compliance Permits
CAP	criteria air pollutant	EFS	Experimental Field Station
CAP88-PC	Clean Air Act Assessment Package-1988 computer model, PC	EJ	environmental justice
CARP	Climate Adaptation and Resilience Plan	EJP	Environmental Justice Program
CCA	Candidate Conservation Agreement	EMS	Environmental Management System
CEJST	Climate and Economic Justice Screening Tool	EO	Executive Order
CEMML	Center for the Environmental Management of Military Lands	EPA	U.S. Environmental Protection Agency
CERCLA	Comprehensive Environmental Response, Compensation, and Liability Act	EPCRA	Emergency Planning and Community Right-to-Know Act
CFA	Central Facilities Area	EPEAT	Electronic Product Environmental Assessment Tool
CFR	Code of Federal Regulations	EPI	emergency plan implementing procedures
CITRC	Critical Infrastructure Test Range Complex	ESA	Endangered Species Act
CRMO	Cultural Resource Management Office	ESPC	Energy Savings Performance Contract
CTF	Contained Test Facility	EV	electric vehicle
CWA	Clean Water Act	FEC	facility emission cap
CWMA	Cooperative Weed Management Area		



FFA/CO	Federal Facility Agreement and Consent Order	LLW	low-level waste
FIFRA	Federal Insecticide, Fungicide, and Rodenticide Act	LOFT	Loss-of-Fluid Test
FY	fiscal year	LTS	Long-Term Stewardship
GPRS	Global Positioning Radiometric Scanner	LTV	long-term vegetation
HeTO	Heritage Tribal Office	MAPEP	Mixed Analyte Performance Evaluation Program
HFC	hydrofluorocarbons	MCL	maximum contaminant level
HLW	high-level waste	MEI	maximally exposed individual
HYSPLIT	Hybrid Single-particle Lagrangian Integrated Trajectory	MFC	Materials and Fuels Complex
IC	institutional control	NA	not applicable
ICDF	Idaho CERCLA Disposal Facility	NCRP	National Council on Radiation Protection and Measurements
ICP	Idaho Cleanup Project	ND	not detected
ICPP	Idaho Chemical Processing Plant	NEPA	National Environmental Policy Act
IDAPA	Idaho Administrative Procedures Act	NERP	National Environmental Research Park
IDFG	Idaho Department of Fish and Game	NESHAP	National Emission Standards for Hazardous Air Pollutants
IDNH	Idaho Museum of Natural History	NHPA	National Historic Preservation Act
IEC	Idaho Environmental Coalition, LLC	NM	not measured
INEEL	Idaho National Engineering and Environmental Laboratory	NOAA	National Oceanic and Atmospheric Administration
INL	Idaho National Laboratory	NON/CO	Notice of Noncompliance/Consent Order
INTEC	Idaho Nuclear Technology and Engineering Center (formerly Idaho Chemical Processing Plant)	NRF	Naval Reactors Facility
IRC	INL Research Center	NS	no sample
ISA	Idaho Settlement Agreement	O&M	Operations & Maintenance
ISB	in situ bioremediation	OSLD	optically stimulated luminescence dosimeter
ISO	International Organization for Standardization	PA	performance assessment
ISU-EAL	Idaho State University-Environmental Assessment Laboratory	PCB	polychlorinated biphenyls
ITEK	Indigenous and Traditional Ecological Knowledge	PCC	Precontact Context
IWCS	Industrial Wastewater Collection System	PCS	primary constituent standard
IWD	Industrial Waste Ditch	PE	performance evaluation
IWTU	Integrated Waste Treatment Unit	PFAS	perfluoroalkyl substances
		PL	primary line
		PT	performance testing
		PTC	permit to construct



PWS	public water system	VARP	Vulnerability Assessment and Resilience Plan
QA	Quality Assurance	VOC	volatile organic compound
QC	Quality Control	WAG	waste area group
RCRA	Resource Conservation and Recovery Act	WFMC	Wildland Fire Management Committee
REC	Research and Education Campus	WMF	Waste Management Facility
RESL	Radiological and Environmental Sciences Laboratory	XRF	x-ray fluorescence spectroscopy
RHLLW	Remote-Handled Low-level Waste Disposal Facility	YOY	year-over-year
RI/FS	Remedial Investigation/Feasibility Study		
ROD	Record of Decision		
RWMC	Radioactive Waste Management Complex		
SBL	Southwestern Branch Line		
SCS	Secondary Constituent Standard		
SDA	Subsurface Disposal Area		
SGCA	Sage-grouse Conservation Area		
SCGN	Species of Greatest Conservation Need		
SMC	Specific Manufacturing Capability		
SMCL	secondary maximum contaminant level		
SNF	spent nuclear fuel		
SSER	Sagebrush Steppe Ecosystem Reserve		
STP	Sewage Treatment Plant		
TAN	Test Area North		
TCE	trichloroethylene		
TFF	Tank Farm Facility		
TLD	thermoluminescent dosimeter		
TMI	Three Mile Island		
TREAT	Transient Reactor Experiment and Test Facility		
TRU	transuranic		
TSCA	Toxic Substances Control Act		
USFWS	U.S. Fish and Wildlife Service		
USGS	U.S. Geological Survey		
UTL	Upper Tolerance Limit		
UTV	utility task vehicle		



*Mountain bluebird*

# Units:



Bq	becquerel	MG	million gallons
C	Celsius	mGy	milligray ( $10^{-3}$ ) gray
cfm	cubic feet per minute	MI	million liters
CFU	colony forming unit	mi	mile
Ci	curie	min	minute
cm	centimeter	mL	milliliter ( $10^{-3}$ ) liter
cps	counts per second	mR	milliroentgen ( $10^{-3}$ ) roentgen
d	day	mrad	milliard ( $10^{-3}$ ) rad
F	Fahrenheit	mSv	millisievert ( $10^{-3}$ ) sievert
ft	feet	oz	ounce
g	gram	pCi	picocurie ( $10^{-12}$ curies)
gal	gallon	R	roentgen
Gy	gray	rad	radiation absorbed dose
ha	hectare	rem	roentgen equivalent man
keV	kilo-electron-volts	Sv	sievert
kg	kilograms ( $10^3$ ) gram	yd	yard
km	kilometer ( $10^3$ ) meter	yr	year
L	liter		
lb	pound		
m	meter		
$\mu$ Ci	microcurie ( $10^{-6}$ ) curies		
$\mu$ g	microgram ( $10^{-6}$ ) grams		
$\mu$ R	microroentgen ( $10^{-6}$ ) roentgen		
$\mu$ S	microsiemen ( $10^{-6}$ ) siemen		
$\mu$ Sv	microsievert ( $10^{-6}$ ) sievert		
Ma	million years		
mCi	millicurie ( $10^{-3}$ ) curies		
MeV	mega electron volt		
mg	milligram ( $10^{-3}$ ) grams		



*Sage-grouse habitat monitoring crew*



# Table of Contents:



ACKNOWLEDGEMENTS .....	v
TO OUR READERS .....	vii
EXECUTIVE SUMMARY .....	ix
HELPFUL INFORMATION .....	xxi
ACRONYMS.....	xxxv
UNITS.....	xxxix
1. INTRODUCTION .....	1-1
1.1 Site Location .....	1-1
1.2 Environmental Setting .....	1-3
1.3 History of the INL Site .....	1-7
1.4 Human Populations Near the INL Site .....	1-8
1.5 INL Site Primary Program Missions and Facilities .....	1-8
1.5.1 Idaho National Laboratory .....	1-8
1.5.2 Idaho Cleanup Project .....	1-9
1.5.3 Primary INL Site Facilities .....	1-9
1.5.4 Independent Oversight and Public Involvement and Outreach .....	1-12
1.5.5 Citizens Advisory Board .....	1-13
1.5.6 Site-wide Monitoring Committees .....	1-13
1.5.7 Environmental Oversight and Monitoring Agreement .....	1-13
1.5.8 Environmental Education Outreach .....	1-14
1.6 References .....	1-15
2. ENVIRONMENTAL COMPLIANCE SUMMARY .....	2-1
2.1 Enforcement and Compliance History Online Database .....	2-1
2.2 Compliance with Requirements .....	2-1
2.3 Environmental and Energy Justice .....	2-24
2.3.1 Initiatives .....	2-25
2.4 INL Site Agreements .....	2-26
2.5 Low-Level and Mixed Radioactive Waste .....	2-28
2.5.1 Spent Nuclear Fuel .....	2-28
2.6 Release and Inventory Reporting at the INL Site .....	2-29
2.6.1 Spills and Releases .....	2-29
2.6.2 Unplanned Releases .....	2-29
2.7 Environmental Permits .....	2-29
2.8 References .....	2-30



3.	ENVIRONMENTAL MANAGEMENT SYSTEMS .....	3-1
3.1	Environmental Management System Structure.....	3-2
3.2	Environmental Policy .....	3-2
3.3	Plan .....	3-2
3.3.1	Environmental Aspects.....	3-2
3.4	Do (Implementation and Operations) .....	3-4
3.4.1	Structure and Responsibility .....	3-4
3.4.2	Competence, Training, and Awareness .....	3-4
3.4.3	Communication .....	3-4
3.4.4	Operational Control .....	3-4
3.4.5	Document and Record Control.....	3-4
3.5	Check .....	3-5
3.6	Act .....	3-5
3.7	INL Site Resiliency .....	3-5
3.7.1	Performance Status.....	3-6
3.7.2	Plans and Projected Performance .....	3-7
3.8	Sustainability Goals.....	3-9
3.9	Environmental Operating Objectives and Targets .....	3-9
3.10	Accomplishments, Awards, and Recognition .....	3-9
3.11	References .....	3-17
4.	ENVIRONMENTAL MONITORING PROGRAMS – AIR.....	4-2
4.1	Organization of Air Monitoring Programs.....	4-2
4.2	Airborne Effluent Monitoring .....	4-6
4.2.1	Hydrofluorocarbon Phasedown .....	4-11
4.3	Ambient Air Monitoring.....	4-12
4.3.1	Ambient Air Monitoring System Design.....	4-12
4.3.2	Air Particulate, Radioiodine, and Tritium Sampling Methods.....	4-14
4.3.3	Ambient Air Monitoring Results.....	4-15
4.3.4	Atmospheric Moisture Monitoring Results.....	4-20
4.3.5	Precipitation Monitoring Results.....	4-20
4.4	Waste Management Environmental Surveillance Air Monitoring.....	4-22
4.4.1	Gross Activity .....	4-22
4.4.2	Specific Radionuclides .....	4-23
4.5	References.....	4-29
5.	ENVIRONMENTAL MONITORING PROGRAMS – LIQUID EFFLUENTS MONITORING .....	5-1
5.1	Liquid Effluent and Related Groundwater Compliance Monitoring.....	5-2
5.1.1	Advanced Test Reactor Complex Cold Waste Ponds .....	5-3
5.1.2	Idaho Nuclear Technology and Engineering Center New Percolation Ponds and Sewage Treatment Plant .....	5-5



5.1.3 Materials and Fuels Complex Industrial Waste Pond ..... 5-7

5.2 Liquid Effluent Surveillance Monitoring..... 5-9

5.2.1 Advanced Test Reactor Complex ..... 5-9

5.2.2 Idaho Nuclear Technology and Engineering Center..... 5-10

5.2.3 Materials and Fuels Complex ..... 5-10

5.3 Waste Management Surveillance Surface Water Sampling ..... 5-10

5.4 References..... 5-12

6. ENVIRONMENTAL MONITORING PROGRAMS – EASTERN SNAKE RIVER PLAIN AQUIFER..... 6-2

6.1 Summary of Monitoring Programs ..... 6-2

6.2 Hydrogeologic Data Management ..... 6-9

6.3 USGS Radiological Groundwater Monitoring at the INL Site ..... 6-9

6.4 USGS Non-radiological Groundwater Monitoring at the INL Site ..... 6-15

6.5 Comprehensive Environmental Response, Compensation, and Liability Act Groundwater Monitoring During 2022 ..... 6-17

6.5.1 Summary of Waste Area Group 1 Groundwater Monitoring Results..... 6-17

6.5.2 Summary of Waste Area Group 2 Groundwater Monitoring Results..... 6-20

6.5.3 Summary of Waste Area Group 3 Groundwater Monitoring Results..... 6-22

6.5.4 Summary of Waste Area Group 4 Groundwater Monitoring Results..... 6-25

6.5.5 Summary of Waste Area Group 7 Groundwater Monitoring Results..... 6-28

6.5.6 Summary of Waste Area Group 9 Groundwater Monitoring Results..... 6-32

6.5.7 Summary of Waste Area Group 10 Groundwater Monitoring Results..... 6-37

6.6 Remote-Handled Low-Level Waste Disposal Facility ..... 6-37

6.7 Onsite Drinking Water Sampling..... 6-39

6.7.1 Idaho National Laboratory Site Drinking Water Monitoring Results ..... 6-39

6.8 Offsite Drinking Water Sampling..... 6-43

6.9 Surface Water Sampling ..... 6-44

6.10 USGS 2022 Publication Abstracts ..... 6-46

6.10.1 Evaluation of Sample Preservation Methods for Analysis of Selected Volatile Organic Compounds in Groundwater at the Idaho National Laboratory, Idaho (Treinen and Bartholomay, 2022)..... 6-47

6.10.2 Historical Development of the U.S. Geological Survey Hydrological Monitoring and Investigative Programs at the Idaho National Laboratory, Idaho, 2002–2020 (Bartholomay, 2022) ..... 6-47

6.10.3 Inpubs—Bibliographic information for the U.S. Geological Survey Idaho National Laboratory Project Office (Fisher, 2022)..... 6-48

6.11 References..... 6-48

7. ENVIRONMENTAL MONITORING PROGRAMS – AGRICULTURAL PRODUCTS, WILDLIFE, SOIL, AND DIRECT RADIATION ..... 7-2

7.1 Agricultural Products and Biota Sampling ..... 7-2

7.2 Sampling Design for Agricultural Products ..... 7-3

7.2.1 Methods ..... 7-4



7.2.2 Milk Results ..... 7-4

7.2.3 Lettuce Results ..... 7-6

7.2.4 Grain Results ..... 7-7

7.2.5 Potato Results ..... 7-7

7.2.6 Alfalfa Results ..... 7-7

7.2.7 Big Game Animals Results ..... 7-7

7.2.8 Waterfowl Results ..... 7-7

7.2.9 Bats Results ..... 7-8

7.3 Soil Sampling ..... 7-9

7.3.1 Soil Sampling Design ..... 7-9

7.3.2 Methods ..... 7-11

7.3.3 Soil Sampling Results ..... 7-11

7.4 Direct Radiation ..... 7-11

7.4.1 Sampling Design ..... 7-11

7.4.2 Methods ..... 7-13

7.4.3 Results ..... 7-13

7.5 Waste Management Surveillance Sampling ..... 7-18

7.5.1 Vegetation Sampling at the Radioactive Waste Management Complex ..... 7-18

7.5.2 Soil Sampling at the Radioactive Waste Management Complex ..... 7-18

7.5.3 Surface Radiation Survey at the Radioactive Waste Management Complex and the Idaho CERCLA Disposal Facility ..... 7-19

7.6 References ..... 7-21

8. DOSE TO THE PUBLIC AND BIOTA ..... 8-1

8.1 Possible Exposure Pathways to the Public ..... 8-2

8.2 Dose to the Public from INL Site Air Emissions ..... 8-2

8.2.1 Maximally Exposed Individual Dose ..... 8-5

8.2.2 Eighty Kilometer (50 Mile) Population Dose ..... 8-8

8.3 Dose to the Public from Ingestion of Wild Game from the INL Site ..... 8-14

8.3.1 Waterfowl ..... 8-14

8.3.2 Big Game Animals ..... 8-14

8.4 Dose to the Public from Drinking Groundwater from the INL Site ..... 8-15

8.5 Dose to the Public from Direct Radiation Exposure along INL Site Borders ..... 8-15

8.6 Dose to the Public from All Pathways ..... 8-15

8.7 Dose to the Public from Operations on the INL Research and Education Campus ..... 8-16

8.8 Dose to Biota ..... 8-16

8.8.1 Introduction ..... 8-16

8.8.2 Terrestrial Evaluation ..... 8-17

8.8.3 Aquatic Evaluation ..... 8-18

8.9 Unplanned Releases ..... 8-23

8.10 References ..... 8-23



9. NATURAL AND CULTURAL RESOURCES CONSERVATION AND PLANNING ..... 9-2

9.1 Special Status Species ..... 9-2

9.1.1 Wildlife..... 9-2

9.1.2 Plants ..... 9-3

9.2 Conservation Planning..... 9-8

9.2.1 Candidate Conservation Agreement for Greater Sage-grouse Onsite ..... 9-8

9.2.2 Bat Protection Plan ..... 9-10

9.2.3 Sagebrush Steppe Ecosystem Reserve ..... 9-11

9.2.4 Migratory Bird Conservation and Avian Protection Planning ..... 9-12

9.2.5 Conservation Action Plan..... 9-13

9.3 Natural Resource Monitoring and Research..... 9-15

9.3.1 Breeding Bird Surveys ..... 9-15

9.3.2 Midwinter Raptor Survey..... 9-17

9.3.3 Long-term Vegetation Transects ..... 9-17

9.3.4 Vegetation Map..... 9-19

9.3.5 National Environmental Research Park..... 9-20

9.4 Land Stewardship ..... 9-22

9.4.1 Wildland Fire Protection Planning, Management, and Recovery ..... 9-22

9.4.2 Restoration and Revegetation ..... 9-23

9.4.3 Weed Management..... 9-25

9.4.4 Ecological Support for National Environmental Policy Act ..... 9-26

9.5 INL Site Cultural Resource Management ..... 9-26

9.5.1 INL Section 106 Project Reviews ..... 9-27

9.5.2 INL Section 110 Research ..... 9-27

9.5.3 Cultural Resource Monitoring ..... 9-29

9.5.4 Stakeholder, Tribal, Public, and Professional Outreach ..... 9-29

9.5.5 INL Archives and Special Collections ..... 9-30

9.6 References..... 9-30

10. QUALITY ASSURANCE OF ENVIRONMENTAL SURVEILLANCE PROGRAMS..... 10-1

10.1 Quality Assurance Policy and Requirements ..... 10-1

10.2 Program Elements and Supporting QA Process..... 10-2

10.2.1 Planning..... 10-2

10.2.2 Sample Collection and Handling ..... 10-4

10.2.3 Sample Analysis ..... 10-4

10.2.4 Data Review and Evaluation ..... 10-5

10.3 QC and PT Sample Results ..... 10-7

10.3.1 2022 MAPEP PT Results ..... 10-8

10.3.2 2022 Field QC Elements ..... 10-9

10.4 Conclusions..... 10-12



10.5 References ..... 10-13

Appendix A. Chapter 5 Addendum

Appendix B. Dosimeter Measurements and Locations

Appendix C. Glossary

## LIST OF FIGURES

Figure ES-1. Regional location of the INL Site .....vii

Figure ES-2. INL Site facilities ..... ix

Figure HI-1. Comparison of penetrating ability of alpha, beta, and gamma radiation..... xx

Figure HI-2. A graphical representation of minimum, median, and maximum results with a box plot .....xxiv

Figure HI-3. Data presented using a pie chart .....xxv

Figure HI-4. Data plotted using a column chart.....xxvi

Figure HI-5. Data plotted using a linear plot..... xxvii

Figure HI-6. Data plotted using contour lines ..... xxviii

Figure HI-7. Calculated doses (mrem per year) from natural background sources for an average individual living in southeast Idaho (2022).....xxx

Figure 1-1. Location of the INL Site ..... 1-2

Figure 1-2. Designated elk and pronghorn hunting boundary on the INL Site ..... 1-3

Figure 1-3. Big Lost River ..... 1-5

Figure 1-4. INL Site relation to the eastern Snake River Plain Aquifer ..... 1-6

Figure 1-5. Location of the INL Site, showing key facilities ..... 1-10

Figure 1-6. Teachers attending joint INL/Museum of Idaho Project Water Education Today workshop..... 1-14

Figure 1-7. Children participating in INL Earth Day activity at the Idaho Falls Zoo ..... 1-15

Figure 1-8. Members of public attending bat night led by INL staff at Idaho Falls Zoo ..... 1-15

Figure 2-1. Students from the Shoshone-Bannock Tribes discussing salmon migration with INL staff .....2-26

Figure 2-2. Radioactive Waste Management Complex Subsurface Disposal Area (2022).....2-28

Figure 4-1. INL Site conceptual model ..... 4-3

Figure 4-2. INL Site environmental surveillance radiological air sampling locations (regional [top] and onsite [bottom]) ..... 4-5

Figure 4-3. Percent contributions in Ci, by facility, to total INL Site airborne radiological releases (2022) .....4-7

Figure 4-4. Locations of INL contractor high-volume event monitors at NOAA weather stations .....4-14

Figure 4-5. Box plots of tritium concentrations measured in atmospheric moisture and in precipitation from 2012–2022. .... 4-22

Figure 4-6. Locations of ICP contractor low-volume air samplers at waste management areas (SDA [top] and ICDF [bottom])..... 4-24

Figure 4-7. Gross alpha (top) and gross beta (bottom) results from waste management site air samples ( $\mu\text{Ci}/\text{mL}$ ) compared to their respective DCSs ..... 4-26

Figure 4-8. Specific human-made radionuclide detections ( $\mu\text{Ci}/\text{mL}$ ) from waste management air samples compared to various fractions of their respective DCSs ..... 4-28

Figure 5-1. Permit monitoring locations for the ATR Complex Cold Waste Pond .....5-4



Figure 5-2. Reuse permit groundwater monitoring locations for INTEC New Percolation Ponds .....5-6

Figure 5-3. INTEC wastewater monitoring for reuse permit ..... 5-7

Figure 5-4. Wastewater and groundwater sampling locations MFC..... 5-8

Figure 5-5. Surface water sampling location at the RWMC SDA .....5-11

Figure 6-1. The eastern Snake River Plain Aquifer and direction of groundwater flow ..... 6-3

Figure 6-2. USGS groundwater monitoring locations on and off the INL Site ..... 6-4

Figure 6-3. Map of the INL Site showing locations of facilities and corresponding WAGs..... 6-6

Figure 6-4. Distribution of tritium (pCi/L) in the eastern Snake River Plain Aquifer onsite in 2018 (from Bartholomay et al. 2020) ..... 6-10

Figure 6-5. Long-term trend of tritium in wells USGS-065 and USGS-114 (2002–2022) ..... 6-11

Figure 6-6. Distribution of <sup>90</sup>Sr (pCi/L) in the eastern Snake River Plain Aquifer onsite in 2018 (from Bartholomay et al. 2020) ..... 6-12

Figure 6-7. Long-term trend of <sup>90</sup>Sr in wells USGS-047, USGS-057, and USGS-113 (2002–2022) ..... 6-13

Figure 6-8. Distribution of <sup>129</sup>I in the eastern Snake River Plain Aquifer onsite in 2017–2018 (from Maimer and Bartholomay 2019) ..... 6-14

Figure 6-9. TCE plume at TAN in 1997 ..... 6-18

Figure 6-10. Distribution of TCE in the Snake River Plain Aquifer from April–June 2022..... 6-19

Figure 6-11. Locations of WAG 2 aquifer monitoring wells ..... 6-21

Figure 6-12. Locations of WAG 3 monitoring wells ..... 6-23

Figure 6-13. Locations of WAG 4/CFA monitoring wells ..... 6-26

Figure 6-14. The WAG 7 aquifer well monitoring network at the RWMC (DOE-ID 2021c)..... 6-29

Figure 6-15. Carbon tetrachloride (CCl<sub>4</sub>) concentration trends in RWMC aquifer Wells M7S, M16S, M3S, and M6S ..... 6-30

Figure 6-16. Carbon tetrachloride (CCl<sub>4</sub>) concentration trends in RWMC aquifer Wells A11A31 and M15S..... 6-30

Figure 6-17. Concentration history of TCE in aquifer Wells M7S, M15S, M16S, A11A31, and M3S..... 6-31

Figure 6-18. Groundwater-level contours in the aquifer near the RWMC, based on 2022 measurements ..... 6-32

Figure 6-19. Locations of WAG 9 wells sampled in 2022..... 6-33

Figure 6-20. Well locations sampled for RHLLW Facility ..... 6-38

Figure 6-21. Detailed map of INL program surface water monitoring locations ..... 6-45

Figure 7-1. Locations of agricultural product samples collected (2022) ..... 7-3

Figure 7-2. Portable lettuce planter ..... 7-6

Figure 7-3. Soil sampling locations in 2022 ..... 7-10

Figure 7-4. Offsite and boundary direct radiation monitoring locations (2022)..... 7-14

Figure 7-5. Historical vegetation sampling areas at the RWMC..... 7-19

Figure 7-6. SDA surface radiation survey area (2022) ..... 7-20

Figure 7-7. ICDF surface radiation survey area (2022) ..... 7-21

Figure 8-1. INL Site major facility airborne source locations ..... 8-3

Figure 8-2. MEI dose from INL Site airborne releases estimated for 2013–2022 ..... 8-6

Figure 8-3. Radionuclides contributing to dose to MEI from INL Site airborne effluents as calculated using the CAP88-PC Model (2022) ..... 8-7

Figure 8-4. Percent contributions, by facility, to MEI dose from the INL Site airborne effluents as calculated using the CAP88-PC Model (2022)..... 8-8



Figure 8-5. Region within 80 km (50 miles) of INL Site facilities ..... 8-9

Figure 8-6. Effective dose (mrem) isopleth map with boundary receptor locations displayed (2022)..... 8-13

Figure 8-7. Radiation doses associated with some common sources ..... 8-16

Figure 9-1. Area defined by the CCA for greater sage-grouse onsite as a SGCA and location of baseline leks used for determining the population trigger..... 9-9

Figure 9-2. Remote and facility BBS routes and north and south midwinter raptor survey routes onsite..... 9-16

Figure 9-3. Locations for the LTV plots established onsite in 1950 and sampled regularly over the past 70 years shown with the INL Site vegetation community classification map published in 2019..... 9-18

Figure 9-4. Researchers studying the flora and fauna of the Idaho NERP ..... 9-20

Figure 9-5. Planters using hoedads to install big sagebrush seedlings onsite..... 9-24

Figure 10-1. Flow of environmental surveillance program elements and associated QA processes and activities..... 10-3

Figure 10-2. QC sampling elements ..... 10-4

Figure 10-3. Laboratory measurement elements ..... 10-5

Figure 10-4. Environmental surveillance field sampling data QA review process..... 10-6

Figure 10-5. 2022 MAPEP PT analyte performance ..... 10-8

Figure 10-6. INL contractor 2022 QC analyte results ..... 10-10

Figure 10-7. ICP contractor 2022 QC analyte results..... 10-11

Figure 10-8. USGS 2022 QC analyte results..... 10-12

Figure B-1. Environmental radiation measurements at Auxiliary Reactor Area (ARA) and Critical Infrastructure Test Range Complex (CITRC) (2022)..... B-1

Figure B-2. Environmental radiation measurements at Advanced Test Reactor (ATR) Complex and Remote-Handled Low-Level Waste Disposal Facility (RHLLW) (2022)..... B-3

Figure B-3. Environmental radiation measurements at Central Facilities Area (CFA) and Lincoln Boulevard (2022) ..... B-4

Figure B-4. Environmental radiation measurements at Idaho Nuclear Technology and Engineering Center (INTEC) (2022)..... B-6

Figure B-5. Environmental radiation measurements at Idaho National Laboratory Research Center Complex (IRC) (2022)..... B-7

Figure B-6. Environmental radiation measurements at Materials and Fuels Complex (MFC) and Transient Reactor Test (TREAT) Facility (2022) ..... B-9

Figure B-7. Environmental radiation measurements at Naval Reactors Facility (NRF) (2022) ..... B-10

Figure B-8. Environmental radiation measurements at IF-675 Portable Isotopic Neutron Spectroscopy (PINS) Laboratory (2022) ..... B-11

Figure B-9. Environmental radiation measurements at Radioactive Waste Management Complex (RWMC) (2022) ..... B-12

Figure B-10. Environmental radiation measurements at Specific Manufacturing Capability (SMC) (2022) ..... B-13

Figure B-11. Environmental radiation measurements at sitewide locations (2022)..... B-14

Figure B-12. Environmental radiation measurements at regional locations (2022) ..... B-16

Figure B-13. Environmental radiation measurements at Willow Creek Building (WCB) and Center for Advanced Energy Studies (CAES) (2022) ..... B-17

Figure B-14. Environmental radiation measurements at Experimental Breeder Reactor I (EBR-I) (2022) ..... B-18

Figure B-15. Environmental radiation measurements at Energy Innovation Laboratory (EIL) (2022)..... B-19





Figure B-16. Environmental radiation measurements at Lindsay Building IF-652A (2022) ..... B-20

## LIST OF TABLES

Table ES-1.	Major INL Site areas and missions .....	x
Table ES-2.	Contribution to estimated annual dose from INL Site facilities by pathway (2022).....	xii
Table HI-1.	Radionuclides and their half-lives .....	xxi
Table HI-2.	Multiples of units.....	xxii
Table HI-3.	Names and symbols for units of radioactivity and radiological dose used in this report.....	xxiii
Table HI-4.	Units of radioactivity .....	xxiii
Table HI-5.	Naturally occurring radionuclides that have been detected in environmental media collected on and around the INL Site .....	xxx
Table 2-1.	Federal, state, and local laws and regulations established for protection of human health and the environment.....	2-2
Table 2-2.	2022 status of active Waste Area Groups.....	2-21
Table 2-3.	Radioactive wastes managed at the INL Site .....	2-23
Table 2-4.	Listing of the status of each phase of the LLW management process for sites authorized to manage a LLW facility .....	2-23
Table 2-5.	Environmental permits for the INL Site (2022).....	2-29
Table 3-1.	Summary table of DOE sustainability goals (DOE-ID 2023).....	3-11
Table 4-1.	Radiological air monitoring activities by organization .....	4-4
Table 4-2.	Radionuclide composition of INL Site airborne effluents (2022) .....	4-10
Table 4-3.	INL Site and regional ambient air monitoring summary (2022).....	4-13
Table 4-4.	Median annual gross alpha concentrations in ambient air samples collected by the INL contractor in 2022.....	4-16
Table 4-5.	Median annual gross beta concentrations in ambient air samples collected the INL contractor in 2022.....	4-18
Table 4-6.	Human-made radionuclides detected in ambient air samples collected by the INL contractor in 2022 .....	4-20
Table 4-7.	Tritium concentrations in atmospheric moisture samples collected by the INL contractor onsite and offsite in 2022 .....	4-21
Table 4-8.	Tritium concentrations in precipitation samples collected by the INL contractor in 2022 .....	4-21
Table 4-9.	Median annual gross alpha concentration in air samples collected at waste management sites in 2022.....	4-25
Table 4-10.	Median annual gross beta concentration in air samples collected at waste management sites in 2022.....	4-25
Table 4-11.	Human-made radionuclides detected in air samples collected at waste management sites in 2022.....	4-27
Table 5-1.	Liquid effluent monitoring at the INL Site .....	5-2
Table 5-2.	2022 status of reuse permits.....	5-3
Table 5-3.	Radionuclides detected in surface water runoff at the RWMC SDA (2022) .....	5-12
Table 6-1.	USGS monitoring program summary (2022).....	6-5
Table 6-2.	ICP contractor drinking water program summary (2022) .....	6-7
Table 6-3.	INL contractor drinking water program summary (2022) .....	6-8
Table 6-4.	INL surface water and offsite drinking water summary (2022).....	6-8



Table 6-5. Purgeable organic compounds in annual USGS groundwater well samples (2022) ..... 6-16

Table 6-6. Purgeable organic compounds in monthly production well samples at the RWMC (2022) ..... 6-17

Table 6-7. WAG 2 aquifer groundwater quality summary (October 2022) ..... 6-22

Table 6-8. Summary of constituents detected in WAG 3 aquifer monitoring wells (FY 2022) ..... 6-24

Table 6-9. Comparison of CFA landfill groundwater sampling results to regulatory levels (August 2022) ..... 6-27

Table 6-10. Summary of WAG 7 aquifer analyses for May 2022 sampling ..... 6-28

Table 6-11. Comparisons of detected analytes to groundwater standards at WAG 9 monitoring wells (2022)..... 6-34

Table 6-12. Radioactivity detected in surveillance groundwater samples collected at the RHLLW Facility (2022)..... 6-37

Table 6-13. Summary of INL Site drinking water results (2022)..... 6-40

Table 6-14. Gross alpha, gross beta, and tritium concentrations in offsite drinking water samples collected by the INL contractor in 2022 ..... 6-43

Table 6-15. Gross alpha, gross beta, and tritium concentrations in surface water samples collected along the Big Lost River by the INL contractor in 2022..... 6-46

Table 7-1. Environmental monitoring of agricultural products, biota, soil, and direct radiation on and around the INL Site..... 7-2

Table 7-2. Strontium and tritium concentrations in milk samples collected offsite in 2022 ..... 7-5

Table 7-3. Radionuclide concentrations detected in waterfowl collected in 2022 ..... 7-8

Table 7-4. Radionuclide concentrations measured in bats collected in 2022 ..... 7-9

Table 7-5. 2022 Soil results compared to background ..... 7-12

Table 7-6. Annual environmental radiation doses using OSLDs at all offsite locations (2018–2022) ..... 7-15

Table 7-7. Dosimetry locations above the six-month background UTL (2022) ..... 7-16

Table 7-8. Calculated effective dose from natural background sources (2022)..... 7-17

Table 8-1. Summary of radionuclide composition of INL Site airborne effluents (2022) ..... 8-4

Table 8-2. Particulate radionuclide source term (Ci yr<sup>-1</sup>) for radionuclide-facility combinations that contributed greater than 0.01% of the total dose for INL Site facilities at the MEI location (2023)..... 8-10

Table 8-3. Noble gases, iodine, tritium and carbon-14 source term (Ci yr<sup>-1</sup>) for radionuclide-facility combinations that contributed greater than 0.01% of the total dose for INL Site facilities at the MEI location (2023)..... 8-11

Table 8-4. Radionuclide source term (Ci yr<sup>-1</sup>) for radionuclides that contributed greater than 0.1% of the total dose for INL in-town facilities (2023) ..... 8-12

Table 8-5. Contribution to estimated annual dose from INL Site facilities by pathway (2022)..... 8-14

Table 8-6. Concentrations of radionuclides in INL Site soils, by area ..... 8-19

Table 8-7. RESRAD Biota assessment (screening level) of terrestrial ecosystems on the INL Site (2022)..... 8-21

Table 8-8. RESRAD Biota assessment (Level 3 analysis) of terrestrial ecosystems on the INL Site using measured bat tissue data (2022) ..... 8-22

Table 8-9. RESRAD Biota assessment (screening level) of aquatic ecosystems on the INL Site (2022) ..... 8-22

Table 8-10. RESRAD Biota assessment (Level 3 analysis) of aquatic ecosystems on the INL Site using measured waterfowl tissue data (2022) ..... 8-23

Table 9-1. Special status animal taxa documented to occur onsite ..... 9-4

Table 9-2. Special status plant taxa documented to occur onsite ..... 9-7

Table 10-1. 2022 analytical laboratories used to analyze surveillance media ..... 10-7



Table A-1. Advanced Test Reactor Complex cold waste pond effluent permit-required monitoring results (2022) ..... A-1

Table A-2. Hydraulic loading rates for the Advanced Test Reactor Complex cold waste pond (2022)..... A-1

Table A-3a. Advanced Test Reactor Complex cold waste pond industrial wastewater reuse permit monitoring well results (2022) ..... A-2

Table A-3b. Advanced Test Reactor Complex cold waste pond industrial wastewater reuse permit monitoring well results (2022) ..... A-4

Table A-4. Idaho Nuclear Technology and Engineering Center sewage treatment plant influent monitoring results at CPP-769 (2022)..... A-4

Table A-5. Idaho Nuclear Technology and Engineering Center sewage treatment plant effluent monitoring results at CPP-773 (2022)..... A-5

Table A-6. Idaho Nuclear Technology and Engineering Center new percolation ponds effluent monitoring results at CPP-797 (2022)..... A-5

Table A-7. Hydraulic loading rates for the Idaho Nuclear Technology and Engineering Center new percolation ponds (2022) ..... A-5

Table A-8. Idaho Nuclear Technology and Engineering Center new percolation ponds aquifer monitoring well groundwater results (2022). ..... A-6

Table A-9. Idaho Nuclear Technology and Engineering Center new percolation ponds perched water monitoring well groundwater results (2022) ..... A-7

Table A-10. Materials and Fuels Complex industrial waste pond effluent monitoring results for the reuse permit (2022) ..... A-8

Table A-11. Materials and Fuels Complex effluent hydraulic loading to the industrial waste pond (2022) ..... A-8

Table A-12. Materials and Fuels Complex industrial waste pond summary of groundwater quality data collected for the reuse permit (2022)..... A-9

Table A-13. Advanced Test Reactor Complex cold waste ponds effluent surveillance monitoring results (2022) ..... A-10

Table A-14. Radioactivity detected in surveillance groundwater samples collected at the Advanced Test Reactor Complex (2022)..... A-10

Table A-15. Liquid effluent radiological monitoring results for the Idaho Nuclear Technology and Engineering Center New Percolation Ponds CPP-797 (2022)..... A-11

Table A-16. Groundwater radiological monitoring results for the Idaho Nuclear Technology and Engineering Center (2022) ..... A-12

Table A-17. Radiological Monitoring Results for Materials and Fuels Complex industrial waste pond (2022) ..... A-12

Table B-1. Results of environmental radiation measurements at Auxiliary Reactor Area (ARA) and Critical Infrastructure Test Range Complex (CITRC) (2022) ..... B-1

Table B-2. Results of environmental radiation measurements at Advanced Test Reactor (ATR) Complex and Remote-Handled Low-Level Waste Disposal Facility (RHLLW) (2022)..... B-2

Table B-3. Results of environmental radiation measurements at Central Facilities Area (CFA) and Lincoln Boulevard (2022) ..... B-4

Table B-4. Results of environmental radiation measurements at Idaho Nuclear Technology and Engineering Center (INTEC) (2022)..... B-5

Table B-5. Results of environmental radiation measurements at Idaho National Laboratory Research Center Complex (IRC) (2022)..... B-7

Table B-6. Results of environmental radiation measurements at Materials and Fuels Complex (MFC) and Transient Reactor Test (TREAT) Facility (2022) ..... B-8

Table B-7. Results of environmental radiation measurements at Naval Reactors Facility (NRF) (2022) ..... B-10

Table B-8. Results of environmental radiation measurements at IF-675 Portable Isotopic Neutron Spectroscopy (PINS) Laboratory (2022)..... B-11



Table B-9. Results of environmental radiation measurements at Radioactive Waste Management Complex (RWMC) (2022)..... B-12

Table B-10. Results of environmental radiation measurements at Specific Manufacturing Capability (SMC) (2022) .... B-13

Table B-11. Results of environmental radiation measurements at sitewide locations (2022)..... B-14

Table B-12. Environmental radiation measurements at regional locations (2022)..... B-15

Table B-13. Results of environmental radiation measurements at Willow Creek Building (WCB) and Center for Advanced Energy Studies (CAES) (2022)..... B-17

Table B-14. Results of environmental radiation measurements at Experimental Breeder Reactor I (EBR-I) (2022) ..... B-18

Table B-15. Results of environmental radiation measurements at Energy Innovation Laboratory (EIL) (2022)..... B-19

Table B-16. Results of environmental radiation measurements at Lindsay Building IF-652A (2022) ..... B-20