

S.M. Stoller Corporation
Environmental Surveillance, Education, and Research Program
ISSN NUMBER 1089-5469

Idaho National Laboratory Site Offsite Environmental Surveillance Program Report: Fourth Quarter 2008

May 2009



*Contributors:
Marilyn Case, Russ Mitchell*

Program conducted for the U.S. Department of Energy, Idaho Operations Office
Under Contract DE-AC07-06ID14680

By the S.M. Stoller Corporation,
Environmental Surveillance, Education and Research Program
Douglas K. Halford, Program Manager
120 Technology Dr., Idaho Falls, Idaho 83402
www.stoller-eser.com

EXECUTIVE SUMMARY

Most of the radionuclides detected in samples collected during the fourth quarter of 2008 could not be directly linked with INL Site activities. One exception was waterfowl collected directly from waste ponds located at one INL facility. Levels of other detected radionuclides were no different than values measured at other locations across the United States or were consistent with levels measured historically at the INL Site. All detected radionuclide concentrations were well below guidelines set by the U.S. Department of Energy (DOE) and regulatory standards established by the U.S. Environmental Protection Agency (EPA) for protection of the public.

This report for the fourth quarter of 2008 contains results from the Environmental Surveillance, Education and Research (ESER) Program's monitoring of the Department of Energy's Idaho National Laboratory (INL) Site's offsite environment, October 1 through December 31, 2008. All sample types (media) and the sampling schedule followed during 2008 are listed in Appendix A. Specifically, this report contains the results for the following:

- Air sampling, including particulate air filters, charcoal cartridges and atmospheric moisture
- Precipitation sampling
- Milk sampling
- Potato sampling
- Game animal sampling
- Waterfowl Sampling
- Soil Sampling

Table E-1 Summary of results for the Fourth Quarter of 2008.

Media	Sample Type	Analysis	Results
Air	Filters	Gross alpha, gross beta	A statistical difference was found between quarterly gross beta concentrations but the pattern of data did not appear to be unusual. There were no other statistical differences in gross alpha or gross beta quarterly or monthly data in the fourth quarter. A couple of statistical differences were noted in weekly results but these appeared to result from natural variability in the data. No result exceeded the DCG for gross alpha or gross beta activity in air.
		Gamma-emitting radionuclides, select actinides (^{241}Am , ^{238}Pu , and $^{239,240}\text{Pu}$), ^{90}Sr	Americium-241 was detected on three composites at concentrations similar to those found previously. Americium-241, Plutonium-238 and Plutonium-239/240 were reported on composites from duplicate sampling locations. After review of the data and discussions with the laboratory, the data for these locations were considered invalid.
	Charcoal Cartridge	Iodine-131	No detections of ^{131}I were made during the fourth quarter.
Atmospheric Moisture	Liquid	Tritium	A total of 14 samples were collected. Seven of these samples had a tritium result greater than the 3s uncertainty. No sample result exceeded the DCG for tritium in air.
Precipitation	Liquid	Tritium	Eleven samples were collected. Three of the results were greater than the 3s uncertainty. The concentration was consistent with those reported across the region and with previous results.
Milk	Liquid	Iodine-131, gamma-emitting radionuclides, ^{90}Sr , tritium	Thirty-four samples, including two duplicates, were collected. No Iodine-131 or other manmade gamma-emitting radionuclides were detected in any sample. Strontium-90 was detected in three of four samples and tritium was detected in two of four samples at levels similar to previous measurements.
Potatoes	Solid	Gamma-emitting radionuclides, ^{90}Sr	Eleven samples were collected, including one duplicate. No man-made gamma-emitting radionuclides were detected. Strontium-90 was detected in six samples at levels consistent with historical measurements and can be attributed to residual ^{90}Sr from past nuclear weapons testing.
Soil	Solid	Gamma emitting radionuclides, ^{90}Sr , ^{241}Am , and plutonium	Cesium-137 was detected in all samples collected. Strontium-90 was detected in most of the samples. The origin of these nuclides is probably deposition of fallout from atmospheric nuclear weapons testing. Americium and Plutonium were not detected in any samples.
Large Game Animals	Tissue	Iodine-131, gamma emitting radionuclides	Two animals were sampled. One mule deer had Cesium-137 in the thyroid at a concentration just above the minimal detectable concentration.
Waterfowl	Tissue	Gamma-emitting	Five radionuclides were found in tissues from

		radionuclides, ⁹⁰ Sr, select actinides (²⁴¹ Am, ²³⁸ Pu, and ^{239,240} Pu)	ducks collected at the Advanced Test Reactor Complex, including edible tissue. Concentrations were generally similar to those found in the previous two years. The estimated dose from eating the entire edible mass of the duck with the highest concentrations was estimated at 0.05 mrem.
Environmental Radiation	TLD	Ambient ionizing radiation	Values were consistent with expected exposures given the altitude and location of the TLDs. There were no statistical differences between Boundary and Distant location results.

LIST OF ABBREVIATIONS

AEC	Atomic Energy Commission
CFA	Central Facilities Area
CMS	community monitoring station
DCG	Derived Concentration Guide
DOE	Department of Energy
DOE – ID	Department of Energy Idaho Operations Office
EAL	Environmental Assessment Laboratory
EFS	Experimental Field Station
EPA	Environmental Protection Agency
ERAMS	Environmental Radiation Ambient Monitoring System
ESER	Environmental Surveillance, Education, and Research
ICP	Idaho Cleanup Project
INL	Idaho National Laboratory
INEL	Idaho National Engineering Laboratory
INEEL	Idaho National Engineering and Environmental Laboratory
ISU	Idaho State University
MDC	minimum detectable concentration
NRTS	National Reactor Testing Station

LIST OF UNITS

Bq	becquerel
Ci	curie
g	gram
L	liter
μ Ci	microcurie
mL	milliliter
mR	milliRoentgen
pCi	picocurie

1. ESER PROGRAM DESCRIPTION

Operations at the Idaho National Laboratory (INL) Site are conducted under requirements imposed by the U.S. Department of Energy (DOE) under authority of the Atomic Energy Act, and the U.S. Environmental Protection Agency (EPA) under a number of acts (e.g. the Clean Air Act and Safe Drinking Water Act). The requirements imposed by DOE are specified in DOE Orders. These requirements include those to monitor the effects of DOE activities both inside and outside the boundaries of DOE facilities (DOE 2003). During calendar year 2008, environmental monitoring within the INL Site boundaries was primarily the responsibility of the INL and Idaho Cleanup Project (ICP) contractors, while monitoring outside the INL Site boundaries was conducted under the Environmental Surveillance, Education and Research (ESER) Program. The ESER Program is led by the S.M. Stoller Corporation in cooperation with its team members, including the University of Idaho, Idaho State University (ISU), the Wildlife Conservation Society and Teledyne Brown Engineering. This report contains monitoring results from the ESER Program for samples collected during the fourth quarter of 2008 (October 1-December 31, 2008).

The surveillance portion of the ESER Program is designed to satisfy the following program objectives:

- Verify compliance with applicable environmental laws, regulations, and DOE Orders
- Characterize and define trends in the physical, chemical, and biological condition of environmental media on and around the INL Site
- Assess the potential radiation dose to members of the public from INL Site effluents
- Present program results clearly and concisely through the use of reports, presentations, newsletter articles and press releases.

The goal of the surveillance program is to monitor different media at a number of potential exposure points within the various exposure pathways, including air, water, agricultural products, wildlife and soil, that could possibly contribute to the radiation dose received by the public.

Environmental samples collected include:

- air at 16 locations on and around the INL Site
- moisture in air at four locations around the INL Site
- precipitation from three locations on and around the INL Site
- agricultural products, including milk at eight dairies around the INL Site, potatoes from at least five local producers, wheat from approximately 10 local producers, and lettuce from approximately nine home-owned and portable gardens on and around the INL
- soil from 12 locations around the INL Site biennially
- environmental dosimeters from 15 locations semi-annually
- various numbers of wildlife including big game (pronghorn, mule deer, and elk) and waterfowl sampled on and near the INL Site.

Table A-1 in Appendix A lists samples, sampling locations and collection frequency for the ESER Program.

The ESER Program used two laboratories to perform analyses on routine environmental samples collected during the quarter reported here. The ISU Environmental Assessment Laboratory (EAL) performed routine gross alpha, gross beta, tritium and gamma spectrometry analyses. Analyses requiring radiochemistry including strontium-90 (⁹⁰Sr), plutonium-238

(²³⁸Pu), plutonium-239/240 (^{239/240}Pu) and americium-241 (²⁴¹Am) were performed by Teledyne Brown Engineering, Inc. of Knoxville, Tennessee.

In the event of non-routine occurrences, such as suspected releases of radioactive material, the ESER Program may increase the frequency of sampling and/or the number of sampling locations based on the nature of the release and wind distribution patterns. Any data found to be outside historical norms in the ESER Program is thoroughly investigated to determine if an INL Site origin is likely. Investigation may include re-sampling and/or re-analysis of prior samples.

In the event of any suspected worldwide nuclear incidents, like the 1986 Chernobyl accident, the EPA may request additional sampling be performed through RadNet [previously known as the Environmental Radiation Ambient Monitoring System (ERAMS) network] (EPA 2008). The EPA established the ERAMS network in 1973 with an emphasis on identifying trends in the accumulation of long-lived radionuclides in the environment. ERAMS was renamed RadNet in 2005 to reflect a new mission. RadNet is comprised of a nationwide network of sampling stations that provide air, precipitation, drinking water and milk samples. The ESER Program currently operates a high-volume air sampler and collects precipitation and drinking water in Idaho Falls for this national program and routinely sends samples to EPA's Eastern Environmental Radiation Facility for analyses. The RadNet data collected at Idaho Falls are not reported by the ESER Program but are available through the EPA RadNet website (<http://www.epa.gov/narel/radnet/>).

Once samples have been collected and analyzed, the ESER Program has the responsibility for quality control of the data and for preparing quarterly reports on results from the environmental surveillance program. The quarterly reports are then consolidated into the INL Site Environmental Report for each calendar year. These annual reports also include data collected by other INL Site contractors.

The results reported in the quarterly and annual reports are assessed in terms of data quality and statistical significance with respect to laboratory analytical uncertainties, sample locations, reported INL Site releases, meteorological data, and worldwide events that might conceivably have an effect on the INL Site environment. First, field collection and laboratory information are reviewed to determine identifiable errors that would invalidate or limit use of the data. Examples of such limitations include insufficient sample volume, torn filters, evidence of laboratory cross-contamination or quality control issues. Data that pass initial screening are further evaluated using statistical methods. Statistical tools are necessary for data evaluation particularly since environmental measurements typically involve the determination of minute concentrations, which are difficult to detect and even more difficult to distinguish from other measurements.

Results are presented in this report with an analytical uncertainty term, *s*, where "s" is the estimated sample standard deviation (σ), assuming a Gaussian or normal distribution. All results are reported in this document, even those that do not necessarily represent detections. The term "detected", as used for the discussion of results in this report, does not imply any degree of risk to the public or environment, but rather indicates that the radionuclide was measured at a concentration sufficient for the analytical instrument to record a value that is statistically different from background. The ESER has adopted guidelines developed by the United States Geological Survey (Bartholomay, et al. 2003), based on an extension of a method proposed by Currie (1984), to interpret analytical results and make decisions concerning detection. Most of the following discussion is taken from Bartholomay et al (2003).

Laboratory measurements involve the analysis of a target sample and the analysis of a prepared laboratory blank (i.e., a sample which is identical to the sample collected in the environment, except that the radionuclide of interest is absent). Instrument signals for the target and blank vary randomly about the true signals and may overlap making it difficult to distinguish between radionuclide activities in blank and in environmental samples (Figure 1). That is, the variability around the sample result may substantially overlap the variability around a net activity of zero for samples with no radioactivity. In order to conclude that a radionuclide has been detected, it is essential to consider two fundamental aspects of the problem of detection: (1) the instrument signal for the sample must be greater than that observed for the blank before the decision can be made that the radionuclide has been detected; and (2) an estimate must be made of the minimum radionuclide concentration that will yield a sufficiently large observed signal before the correct decision can be made for detection or non-detection.




Figure 1. Example of overlap of blank and sample measurement distributions.

In the laboratory, instrument signals must exceed a critical level of $1.6s$ before the qualitative decision can be made as to whether the radionuclide was detected in a sample. At $1.6s$ there is about a 95-percent probability that the correct conclusion—not detected—will be made. Given a large number of samples, approximately 5 percent of the samples with measured concentrations greater than or equal to $1.6s$, which were concluded as being detected, might not contain the radionuclide. These are referred to as false positives. For purposes of simplicity and consistency with past reporting, the ESER has rounded the $1.6s$ critical level estimate to $2s$.

Once the critical level has been defined, the minimum detectable concentration may be determined. Concentrations that equal $3s$ represent a measurement at the detection level or minimum detectable concentration. For true concentrations of $3s$ or greater, there is a 95-percent probability that the radionuclide was detected in the target sample. In a large number of samples, the conclusion—not detected—will be made in 5 percent of the samples with true concentrations at the minimum detectable concentration of $3s$. These measurements are known as false negatives. The ESER reports measured radionuclide concentrations greater than or equal to their respective $3s$ uncertainties as being “detected with confidence.”

Concentrations between $2s$ and $3s$ are reported as “questionably detected”. That is, the radionuclide may be present in the sample; however, the detection may not be reliable.

Measurements made between 2s and 3s are examined further to determine if they are a part of a pattern (temporal or spatial) that might warrant further investigation or recounting. For example, if a particular radionuclide is typically detected at > 3s at a specific location, a sample result between 2s and 3s might be considered detected.

If a result is less than or equal to 2s there is little confidence that the radionuclide is present in the sample. Analytical results in this report are presented as the result value \pm one standard deviation (1s) for reporting consistency with the annual report. To obtain the 2s or 3s values simply multiply the uncertainty term by 2 or 3.

For more information concerning the ESER Program, contact the S.M. Stoller Corporation at (208) 525-9358, or visit the Program's web page (<http://www.stoller-eser.com>).

2. THE INL SITE

The INL Site is a nuclear energy and homeland security research and environmental management facility. It is owned and administered by the U.S. Department of Energy, Idaho Operations Office (DOE-ID) and occupies about 890 mi² (2300 km²) of the upper Snake River Plain in Southeastern Idaho. The history of the INL Site began during World War II when the U.S. Naval Ordnance Station was located in Pocatello, Idaho. This station, one of two such installations in the U.S., retooled large guns from U.S. Navy warships. The retooled guns were tested on the nearby, uninhabited plain, known as the Naval Proving Ground. In the years following the war, as the nation worked to develop nuclear power, the Atomic Energy Commission (AEC), predecessor to the DOE, became interested in the Naval Proving Ground and made plans for a facility to build, test, and perfect nuclear power reactors.

The Naval Proving Ground became the National Reactor Testing Station (NRTS) in 1949, under the AEC. By the end of 1951, a reactor at the NRTS became the first to produce useful amounts of electricity. Over time the site has operated 52 various types of reactors, associated research centers, and waste handling areas. The NRTS was renamed the Idaho National Engineering Laboratory (INEL) in 1974, and the Idaho National Engineering and Environmental Laboratory (INEEL) in January 1997. With renewed interest in nuclear power the DOE announced in 2003 that Argonne National Laboratory and the INEEL would be the lead laboratories for development of the next generation of power reactors. On February 1, 2005 the INEEL and Argonne National Laboratory-West became the INL. The INL is committed to providing international nuclear leadership for the 21st Century, developing and demonstrating compelling national security technologies, and delivering excellence in science and technology as one of the Department of Energy's multiprogram national laboratories.

The cleanup operation, the ICP, is now a separately managed effort. The ICP is charged with safely and cost-effectively completing the majority of cleanup work from past laboratory missions by 2012.



3. AIR SAMPLING

The primary pathway by which radionuclides can move off the INL Site is through the air and for this reason the air pathway is the primary focus of monitoring on and around the INL Site. Samples for particulates and iodine-131 (^{131}I) gas in air were collected weekly for the duration of the quarter at 16 locations using low-volume air samplers. Moisture in the atmosphere was sampled at four locations around the INL Site and analyzed for tritium. Air sampling activities and results for the fourth quarter of 2008 are discussed below. A summary of approximate minimum detectable concentrations (MDCs) for radiological analyses and DOE Derived Concentration Guide (DCG) (DOE 1993) values is provided in Appendix B.

LOW-VOLUME AIR SAMPLING

Radioactivity associated with airborne particulates was monitored continuously by 18 low-volume air samplers (two of which are used as replicate samplers) at 16 locations during the fourth quarter of 2008 (Figure 2). Three of these samplers are located on the INL Site, nine are situated off the INL Site near the boundary and six have been placed at locations distant to the INL Site. Samplers are divided into INL Site, Boundary and Distant groups to determine if there is a gradient of radionuclide concentrations, increasing towards the INL Site. Each replicate sampler is relocated every other year to a new location. At the start of 2008, one replicate sampler was moved to Blue Dome (a Boundary location) and one was moved to Atomic City (also a Boundary location). An average of 15,381 ft³ (436 m³) of air was sampled at each location, each week, at an average flow rate of 1.53 ft³/min (0.04 m³/min). Particulates in air were collected on membrane particulate filters (1.2- μm pore size). Gases passing through the filter were collected with an activated charcoal cartridge.




Figure 2. Low-volume air sampler locations.

Filters and charcoal cartridges were changed weekly at each station during the quarter. Each particulate filter was analyzed for gross alpha and gross beta radioactivity using thin-window gas flow proportional counting systems after waiting about four days for naturally-occurring daughter products of radon and thorium to decay.

The weekly particulate filters collected during the quarter for each location were composited and analyzed for gamma-emitting radionuclides. Selected composites were also analyzed by location for ^{90}Sr , ^{238}Pu , $^{239/240}\text{Pu}$ and ^{241}Am as determined by a rotating quarterly schedule.

Charcoal cartridges were analyzed for gamma-emitting radionuclides, specifically for iodine-131 (^{131}I). Iodine-131 is of particular interest because it is produced in relatively large quantities by nuclear fission, is readily accumulated in human and animal thyroids, and has a half-life of eight days. This means that any elevated level of ^{131}I in the environment could be from a recent release of fission products.

Gross alpha results are reported in Table C-1. Median gross alpha concentrations in air for INL Site, Boundary, and Distant locations for the fourth quarter of 2008 are shown in Figure 3. Gross alpha data are tested for normality prior to statistical analyses, and generally show no consistent discernable distribution. Box and whisker plots are commonly used when there is no assumed distribution. Each data group in Figure 3 is presented as a box and whisker plot, with a median (small red square), a box enclosing values between the 25th and 75th percentiles, and whiskers representing the non-outlier range. Outliers and extreme values are identified separately from the box and whiskers. Outliers and extreme values are atypical, infrequent data points that are far from the middle of the data distribution. For this report, outliers are defined as values that are greater than 1.5 times the height of the box, above or below the box. Extreme values are greater than 2 times the height of the box, above or below the box. Outliers and extreme values may reflect inherent variability, may be due to errors associated with transcription or measurement, or may be related to other anomalies. A careful review of the data collected during the fourth quarter indicates that the outlier values were not due to mistakes in collection, analysis, or reporting procedures, but rather reflect natural variability in the measurements. The outlier values lie within the range of measurements made within the past several years. Thus, rather than dismissing the outliers, they were included in the subsequent statistical analyses.

Because there is no discernable distribution of the data, the nonparametric Kruskal-Wallis test of multiple independent groups was used to test for statistical differences between INL Site, Boundary and Distant locations. The use of nonparametric tests, such as Kruskal-Wallis, gives less weight to outlier and extreme values thus allowing a more appropriate comparison of data groups. A statistically significant difference exists between data groups if the (p) value is less than 0.05. Values greater than 0.05 translate into a 95 percent confidence that the medians are statistically the same. The p-value for each comparison is shown in Table D-1. For the quarter, there was no statistical difference noted. Figure 3 graphically shows that the gross alpha measurements made at INL Site, Boundary and Distant locations are similar for the fourth quarter. If the INL Site were a significant source of offsite contamination, concentrations of contaminants could be statistically greater at Boundary locations than at Distant locations.

Comparisons of gross alpha concentrations were made for each month of the quarter (Figures 4 – 6). Again the Kruskal-Wallis test of multiple independent groups was used to determine if statistical differences exist between INL Site, Boundary and Distant data groups.




Figure 3. Gross alpha concentrations in air at ESER INL Site, Boundary and Distant locations for the fourth quarter of 2008.




Figure 4. October gross alpha concentrations in air at ESER INL Site, Boundary and Distant locations. Number of samples (N) = 5 at each location, except FAA Tower (N=4) and Jackson (N = 3).




Figure 5. November gross alpha concentrations in air at ESER INL Site, Boundary and Distant locations. Number of samples (N) = 4 at each location, except Jackson (N = 3).




Figure 6. December gross alpha concentrations in air at ESER INL Site, Boundary and Distant locations. Number of samples (N) = 5 at each location, except Blue Dome and Van Buren Gate (N = 4).

No statistical differences in gross alpha concentrations between groups were noted during any month of the quarter (Table D-1).

As an additional check, comparisons between gross alpha concentrations measured at Boundary and Distant locations were made on a weekly basis. The Mann-Whitney U test was used to compare the Boundary and Distant data because it is the most powerful nonparametric alternative to the t-test for independent samples. INL Site sample results were not included in this analysis because the onsite data, collected at only three locations, are not representative of the entire INL Site and would not aid in determining offsite impacts. In the fourth quarter, there was one week (November 5) where a statistical difference existed between the two sample groups (Table D-2). The Distant group was statistically greater than the Boundary group in this case.

Gross beta results are presented in Table C-1. Gross beta concentrations in air for INL Site, Boundary and Distant locations for the fourth quarter of 2008 are shown in Figure 7. The data were tested and found to be neither normally nor log-normally distributed. Box and whiskers plots were used for presentation of the data. Outliers and extreme values were retained in subsequent statistical analyses because they are within the range of measurements made in the past five years, and because these values could not be attributed to mistakes in collection, analysis, or reporting procedures. A statistical difference was noted in the quarterly data using the Kruskal-Wallis test (Table D-1). Figure 7, however, shows the values to be similar for all three groups and a review of gross beta data for the quarter did not indicate any unusual patterns in the distribution of results.

Monthly median gross beta concentrations in air for each sampling group are shown in Figures 8 – 10. Statistical data are presented in Table D-1. There were no statistical differences in gross beta between groups for any month during the quarter.

Comparison of weekly Boundary and Distant gross beta data sets, using the Mann Whitney U test, showed statistical differences between Boundary and Distant measurements during three weeks in the fourth quarter (Table D-2). In all three cases, the Boundary group was higher than the Distant group. Analysis of the data for these weeks showed no particular pattern in the results. On December 10, for example, the Distant group data was influenced by a low concentration for the Jackson sampler, which ran for a longer period of time due to a scheduling conflict with the station manager the previous week and thus had a higher volume. Similarly, in the November 12 data set, Rexburg had a lower concentration than the rest of the data set.

No ^{131}I was detected in any of the charcoal cartridge batches collected during the fourth quarter of 2008. Weekly ^{131}I results for each location are listed in Table C-2 of Appendix C. Gamma spectrographic analysis is also done with the ^{131}I analysis. Cesium-137 was not detected in any of the 26 measured batches of cartridges this quarter. The analytical laboratory considers occasional detections of this nuclide a result of the materials used in the charcoal filters.

Weekly filters for the fourth quarter of 2008 were composited by location. All samples were analyzed for gamma-emitting radionuclides, including ^{137}Cs (see Table C-3, Appendix C). No manmade gamma-emitting radionuclides were detected.

Composites were also analyzed for ^{90}Sr , ^{238}Pu , $^{239/240}\text{Pu}$ and ^{241}Am (see Table C-3, Appendix C). Americium-241 was detected on composites from Atomic City, Howe, and the Experimental Field Station at levels consistent with previous detections. In addition, ^{241}Am and ^{238}Pu were detected on the duplicate sampler located at Atomic City and $^{239/240}\text{Pu}$ was detected on the composite from the duplicate sampler at Blue Dome. The concentrations found on the duplicate sampler composites were far higher than the normal range of detections. After a

thorough review of the data and discussions with the analytical laboratory, these detections are considered invalid for the following reasons:

- Results for the duplicate samplers were completely inconsistent with those of the co-located regular samplers.
- Recoveries for the radionuclides detected were reported to be poor by the laboratory.
- The deposition on the sample mount appeared to be non-uniform by the laboratory.
- The laboratory recounted the samples at the request of the ESER Program and results of the recounts were inconsistent with the original counts.

ATMOSPHERIC MOISTURE SAMPLING

Fourteen atmospheric moisture samples were obtained during the fourth quarter of 2008 from Atomic City, Blackfoot CMS, Idaho Falls and Rexburg CMS. Atmospheric moisture is collected by pulling air through a column of absorbent material (molecular sieve material) to absorb water vapor. The water is then extracted from the absorbent material by heat distillation. The resulting water samples are then analyzed for tritium using liquid scintillation.

Seven of the 14 samples exceeded the 3s uncertainty level for tritium, with similar results reported from all four locations. All samples were significantly below the DOE DCG for tritium in air of $1 \times 10^{-7} \mu\text{Ci}/\text{mL}_{\text{air}}$ with a maximum reported value of $(13.8 \pm 1.9) \times 10^{-13} \mu\text{Ci}/\text{mL}_{\text{air}}$. All results are shown in Table C-4, Appendix C.




Figure 7. Gross beta concentrations in air at ESER INL Site, Boundary and Distant locations for the fourth quarter of 2008.




Figure 8. October gross beta concentrations in air at ESER INL Site, Boundary and Distant locations. Number of samples (N) = 5 at each location, except FAA Tower (N = 4) and Jackson (N = 3).




Figure 9. November gross beta concentrations in air at ESER INL Site, Boundary and Distant locations. Number of samples (N) = 4 at each location, except Jackson (N = 3).




Figure 10. December gross beta concentrations in air at ESER INL Site, Boundary and Distant locations. Number of samples (N) = 5 at each location, except Blue Dome and Van Buren Gate (N = 4).

4. PRECIPITATION SAMPLING

Precipitation samples are gathered when sufficient precipitation occurs to allow for the collection of the minimum sample volume of approximately 20 mL. Samples are taken of monthly composites from Idaho Falls and CFA, and weekly from the EFS. Precipitation samples are analyzed for tritium. Storm events in the fourth quarter of 2008 produced sufficient precipitation to yield 11 samples.

Tritium was measured above the 3s value in three of the eleven samples collected during the fourth quarter of 2008. Low levels of tritium exist in the environment at all times as a result of cosmic ray reactions with water molecules in the upper atmosphere. The EPA's RadNet program collects precipitation samples from across the United States. From 1980 to 2005, tritium measured in samples from Region 10 (which includes Idaho) ranged from -200 to 7500 pCi/L (EPA 2007). Data for all fourth quarter 2008 precipitation samples collected by the ESER Program were at the low end of this range and are listed in Table C-5 (Appendix C).



5. AGRICULTURAL PRODUCT, WILDLIFE AND SOIL SAMPLING

Another potential pathway for contaminants to reach humans is through the food chain. The ESER Program samples multiple agricultural products and game animals from around the INL Site and Southeast Idaho. Specifically, milk, wheat, potatoes, garden lettuce, big game, and waterfowl are sampled. Milk is sampled throughout the year and large game animals are sampled whenever large game animals are killed onsite from vehicle collisions. Lettuce and wheat are sampled during the third quarter, while potatoes are collected during the fourth quarter. Waterfowl are collected in either the third or fourth quarter. See Table A-1, Appendix A, for more details on agricultural product and wildlife sampling. This section discusses results from milk, lettuce, wheat and large game animals sampled during the fourth quarter of 2008.

MILK SAMPLING

Milk samples were collected weekly in Idaho Falls. Monthly samples were collected at seven other locations around the INL Site (Figure 11) during the fourth quarter of 2008. All samples were analyzed for gamma emitting radionuclides. During the second quarter, samples from half of the locations are analyzed for ^{90}Sr and half are analyzed for tritium. In the fourth quarter the analyses are reversed, so that each location receives one analysis for ^{90}Sr and tritium each year.

No Iodine-131 or other gamma-emitting radionuclides were detected in any sample. Data for ^{131}I and ^{137}Cs in milk samples are listed in Appendix C, Table C-6.

Strontium-90 was detected in three of four samples analyzed at levels within historical measurements, ranging from 0.24 to 0.67 pCi/L (Table C-7 in Appendix C.) Tritium was detected in one of four samples analyzed, also well within the range of historical measurements (Table C-7).




Figure 11. ESER milk sampling locations.

POTATO SAMPLING

Eight potato samples were collected from area growers and from two out-of-state locations (Colorado and Oregon). All samples were analyzed for gamma emitting radionuclides and ^{90}Sr . Cesium-137 was not measured in any samples. Strontium-90 was detected in two of the samples—Idaho Falls and Minidoka. All values were within historic concentrations measured in potatoes collected from farms surrounding the INL and out-of-state areas.

Data for ^{137}Cs and ^{90}Sr in all potato samples taken during the fourth quarter are listed in Table C -8 (Appendix C).

SOIL SAMPLING

Thirteen soil samples (including one duplicate at Mud Lake #1) were collected at boundary and offsite locations in the third quarter and full results were available in the fourth quarter. All samples were analyzed for gamma-emitting radionuclides, ^{241}Am , ^{238}Pu , $^{239/240}\text{Pu}$, and ^{90}Sr (Tables C-9 and C-10). Cesium-137 was detected in all samples at concentrations consistent with historical measurements and is most likely present from past atmospheric nuclear weapons testing fallout. Similarly ^{90}Sr , another fallout radionuclide, was detected in nine of the 13 soil samples at levels within historical measurements.

None of the transuranic radionuclides were detected in any of the samples.

LARGE GAME ANIMAL SAMPLING

Two large game animals (one mule deer and one elk) were sampled on the INL Site during the fourth quarter of 2008. No manmade radionuclides were found in any of the muscle and liver samples. No ^{131}I was detected in either of the thyroid samples, but ^{137}Cs was reported just above the detection limit in the mule deer thyroid.

The data for ^{137}Cs and ^{131}I are listed in Appendix C, Table C-11.

WATERFOWL SAMPLING

Nine ducks were collected during 2008. Four were collected from wastewater ponds located at the Advanced Test Reactor Complex (ATR-C), three came from wastewater ponds near the Materials and Fuels Complex (MFC), and two control samples were collected from Mud Lake. Each sample was divided into the following three sub-samples: 1) edible tissue (muscle, gizzard, heart and liver), 2) viscera, and 3) all remaining tissue (bones, feathers, feet, bill, head, and residual muscle). All were analyzed for gamma-emitting radionuclides, ^{90}Sr , plutonium-238 (^{238}Pu), plutonium-239,240 ($^{239/240}\text{Pu}$), and americium-241 (^{241}Am). Concentrations of radionuclides measured in the edible tissues of 2008 waterfowl are shown in Table C-12 (Appendix C).

Several manmade radionuclides were detected in the samples taken from the ATR-C ponds, including ^{241}Am , ^{137}Cs , cobalt-60 (^{60}Co), ^{90}Sr , and zinc-65 (^{65}Zn). All were also found in at least one edible tissue sample. Birds from the MFC ponds contained ^{137}Cs and ^{90}Sr but neither was found in edible tissues. One detection each for ^{241}Am , ^{238}Pu and $^{239/240}\text{Pu}$ were reported in the control samples.

Since manmade radionuclides were found more frequently and at higher concentrations in ducks taken from the INL Site than in those from other locations, it is assumed that the INL Site is the source of these radionuclides. Concentrations of the detected radionuclides from ATR-C were similar to those from 2006 and 2007, or significantly lower in the case of ^{137}Cs than those found in 2005. The ducks were not taken directly from the two-celled hypalon-lined radioactive wastewater evaporation pond but rather from an adjacent sewage lagoon. However, it is likely

that the birds also used the evaporation pond. Measured concentrations were also lower than those in ducks taken during a 1994-1998 study (Warren et al. 2001).

Waterfowl hunting is not allowed on the INL Site, but a maximum potential exposure scenario to humans would be someone collecting a contaminated duck directly from the ponds and immediately consuming all muscle, liver, heart, and gizzard tissue. The maximum potential dose from eating 225 g (8 oz) of meat from the most contaminated waterfowl collected in 2008 was estimated to be 0.05 mrem. This dose is lower than dose estimates for some previous periods. The dose estimated for 2007 was 0.015 mrem. The maximum dose estimated for the period from 1993 through 1998 was 0.89 mrem and from 2000 through 2004 was 0.08 mrem. In the late 1970s, when the percolation ponds were still in use, the maximum dose estimated from eating a contaminated duck was estimated to be 54 mrem.

6. ENVIRONMENTAL RADIATION

An array of thermoluminescent dosimeters (TLDs) is distributed throughout the Eastern Snake River Plain to monitor for environmental radiation (Figure 12). TLDs are changed out in May and again in November after six months in the field. The results of the TLDs exposed from November 2007 to May 2008 are discussed below.




Figure 12. TLD locations.

Similar to the low-volume air results the environmental dosimeter locations are also divided into Boundary and Distant groupings. Boundary average exposure rates ranged from a low of 0.30 mR/day at Blue Dome and Birch Creek to a high of 0.36 mR/day at Mud Lake. The overall Boundary average was 0.33 mR/day. The Distant group had a high of 0.40 mR/day at Rexburg and a low of 0.28 mR/day at the Dubois location. The overall average Distant value was also 0.33 mR/day. There was no statistical difference between Boundary and Distant locations. Furthermore, all values are consistent with past readings. All results are listed in Appendix C, Table C-13.

7. QUALITY ASSURANCE

The ESER Quality Assurance Program consists of five ongoing tasks which measure:

1. method uncertainty
2. data completeness
3. data accuracy, using spike, performance evaluation and laboratory control samples
4. data precision, using split samples, duplicate samples and recounts
5. presence of contamination in samples, using blanks.

Sample results are compared to criteria described in the Quality Assurance Project Plan for the INL Site Offsite Environmental Surveillance Program (Stoller 2007). The following table summarizes the results of the quality assurance program for the fourth quarter of 2008.

QA Sample Type	Number of Sample Results	Number of Results Meeting Criteria	Percentage Meeting Criteria
Spikes/Laboratory Control Samples	243	240	98.8
Field Duplicates	85	73	85.9
Laboratory Splits	32	32	100.0
Recounts	177	176	99.4
Blanks	85	83	97.6
Method Uncertainty	1888	1839	97.4

8. REFERENCES

- Bartholomay, R.C., Knobel, L.L., and Rousseau, J.P., 2003, *Field Methods and Quality Plan for Quality-of-Water Activities*, U.S. Geological Survey, Idaho National Engineering and Environmental Laboratory, Idaho, DOE/ID-22182, January 2003.
- Code of Federal Regulations (CFR), 2007, 40 CFR 50.6, "National Primary and Secondary Ambient Air Quality Standards for Particulate Matter," Code of Federal Regulations, Office of the Federal Register, July 1, 2007.
- Currie, L.A., 1984, *Lower Limit of Detection: Definition and Elaboration of a Proposed Position for Radiological Effluent and Environmental Measurements*, NUREG/CR-4007, U.S. Nuclear Regulatory Commission, Washington, D.C., September 1984.
- DOE, 2003, "Environmental Management System," U.S. Department of Energy Order 450.1, January 2003.
- DOE, 1993, "Radiation Protection of the Public and the Environment," U.S. Department of Energy Order 5400.5, January 1993.
- EPA, 2008, RadNet—Tracking Environmental Radiation Nationwide, Web-page: <http://www.epa.gov/narel/radnet/>
- Stoller, 2007, *Quality Assurance Project Plan for the INL Site Offsite Environmental Surveillance Program*, Environmental Surveillance, Education and Research Program, February 2007.

APPENDIX A
SUMMARY OF SAMPLING SCHEDULE

Table A-1. Summary of the ESER Program's Sampling Schedule

Sample Type Analysis	Collection Frequency	LOCATIONS		
		Distant	Boundary	INL Site
AIR SAMPLING				
<i>LOW-VOLUME AIR</i>				
Gross Alpha, Gross Beta, ¹³¹ I	weekly	Blackfoot, Craters of the Moon, Dubois, Idaho Falls, Jackson WY, Rexburg	Arco, Atomic City, FAA Tower, Howe, Montevieu, Mud Lake, Blue Dome	Main Gate, EFS, Van Buren
Gamma Spec	quarterly	Blackfoot, Craters of the Moon, Dubois, Idaho Falls, Jackson WY, Rexburg	Arco, Atomic City, FAA Tower, Howe, Montevieu, Mud Lake, Blue Dome	Main Gate, EFS, Van Buren
⁹⁰ Sr, Transuranics	quarterly	Rotating schedule	Rotating schedule	Rotating schedule
<i>ATMOSPHERIC MOISTURE</i>				
Tritium	2 to 13 weeks	Blackfoot, Idaho Falls, Rexburg	Atomic City	None
<i>PRECIPITATION</i>				
Tritium	monthly	Idaho Falls	None	CFA
Tritium	weekly	None	None	EFS
ENVIRONMENTAL RADIATION SAMPLING				
<i>TLDs</i>				
Gamma Radiation	semiannual	Aberdeen, Blackfoot (2), Craters of the Moon, Dubois, Idaho Falls, Jackson WY, Minidoka, Rexburg, Roberts	Arco, Atomic City, Birch Creek, Blue Dome, Howe, Montevieu, Mud Lake	None
SOIL SAMPLING				
<i>SOIL</i>				
Gamma Spec, ⁹⁰ Sr, Transuranics	biennially	Carey, Crystal Ice Caves (Aberdeen), Blackfoot, St. Anthony	Butte City, Montevieu, Atomic City, FAA Tower, Howe, Mud Lake (2), Birch Creek	None

Table A-1. Summary of the ESER Program's Sampling Schedule (continued)

Sample Type Analysis	Collection Frequency	LOCATIONS		
		Distant	Boundary	INL Site
FOODSTUFF SAMPLING				
<i>MILK</i>				
Gamma Spec (¹³¹ I)	weekly	Idaho Falls	None	None
Gamma Spec (¹³¹ I)	monthly	Blackfoot, Dietrich, Fort Hall, Idaho Falls, Minidoka, Moreland	Howe, Terreton	None
Tritium, ⁹⁰ Sr	Semi-annually	Blackfoot, Dietrich, Fort Hall, Idaho Falls, Minidoka, Moreland	Howe, Terreton	None
<i>POTATOES</i>				
Gamma Spec, ⁹⁰ Sr	annually	Aberdeen, Blackfoot, Fort Hall, Idaho Falls, Rupert, Taber, occasional samples across the U.S.	Arco, Monteview, Mud Lake, Terreton	None
<i>WHEAT</i>				
Gamma Spec, ⁹⁰ Sr	annually	American Falls, Blackfoot, Dietrich, Idaho Falls, Minidoka, Carey	Arco, Monteview, Mud Lake, Taber, Terreton	None
<i>LETTUCE</i>				
Gamma Spec, ⁹⁰ Sr	annually	Blackfoot, Carey, Idaho Falls, Pocatello	Arco, Atomic City, FAA Tower, Howe, Monteview	EFS
<i>BIG GAME</i>				
Gamma Spec	varies	Occasional samples across the U.S.	Public Highways	INL Site roads
<i>WATERFOWL</i>				
Gamma Spec, ⁹⁰ Sr, Transuranics	annually	Varies among: Heise, Firth, Fort Hall, Mud Lake, Market Lake, and American Falls	None	Wastewater disposal ponds

APPENDIX B
SUMMARY OF MDCs AND DCGs

Table B-1. Summary of Approximate Minimum Detectable Concentrations for Radiological Analyses Performed During Fourth Quarter 2008

Sample Type	Analysis	Approximate Minimum Detectable Concentration ^a (MDC)	Derived Concentration Guide ^b (DCG)
Air (particulate filter) ^e	Gross alpha ^c	4.62×10^{-16} $\mu\text{Ci/mL}$	2×10^{-14} $\mu\text{Ci/mL}$
	Gross beta ^d	1.57×10^{-15} $\mu\text{Ci/mL}$	3×10^{-12} $\mu\text{Ci/mL}$
	Specific gamma (¹³⁷ Cs)	1.99×10^{-16} $\mu\text{Ci/mL}$	4×10^{-10} $\mu\text{Ci/mL}$
	²³⁸ Pu	2.57×10^{-17} $\mu\text{Ci/mL}$	3×10^{-14} $\mu\text{Ci/mL}$
	^{239/240} Pu	1.10×10^{-17} $\mu\text{Ci/mL}$	2×10^{-14} $\mu\text{Ci/mL}$
	²⁴¹ Am	4.37×10^{-18} $\mu\text{Ci/mL}$	2×10^{-14} $\mu\text{Ci/mL}$
	⁹⁰ Sr	4.62×10^{-17} $\mu\text{Ci/mL}$	9×10^{-12} $\mu\text{Ci/mL}$
Air (charcoal cartridge) ^e	¹³¹ I	3.53×10^{-16} $\mu\text{Ci/mL}$	4×10^{-10} $\mu\text{Ci/mL}$
Air (atmospheric moisture)	³ H	109.3 pCi/L _{water}	1×10^{-7} $\mu\text{Ci/mL}_{\text{air}}$
Air (precipitation)	³ H	111.5 pCi/L	2×10^{-3} $\mu\text{Ci/mL}$
Milk	¹³¹ I	0.72 pCi/L	--
	¹³⁷ Cs	1.53 pCi/L	--
	⁹⁰ Sr	0.17 pCi/L	--
	³ H	113.6 pCi/L	--
Potatoes	⁹⁰ Sr	0.0044 pCi/g	--
Soil	¹³⁷ Cs	2.62×10^{-9} $\mu\text{Ci/g}$	--
	⁹⁰ Sr	0.04 pCi/g	--
	²³⁸ Pu	0.04 pCi/g	--
	^{239/240} Pu	0.03 pCi/g	--
	²⁴¹ Am	0.065 pCi/g	--
<p>a The MDC is an estimate of the concentration of radioactivity in a given sample type that can be identified with a 95 percent level of confidence and precision of plus or minus 100 percent under a specified set of typical laboratory measurement conditions.</p> <p>b DCGs, set by the DOE, represent reference values for radiation exposure. They are based on a radiation dose of 100 mrem/yr for exposure through a particular exposure mode such as direct exposure, inhalation, or ingestion of water.</p> <p>c The DCG for gross alpha is equivalent to the DCGs for ^{239,240}Pu and ²⁴¹Am.</p> <p>d The DCG for gross beta is equivalent to the DCGs for ²²⁸Ra</p> <p>e The approximate MDC is based on an average filtered air volume (pressure corrected) of 445 m³/week.</p>			

APPENDIX C
SAMPLE ANALYSIS RESULTS

TABLE C-1. Weekly Gross Alpha and Gross Beta Concentrations in Air

Sampling Group and Location	Sampling Date	GROSS ALPHA			GROSS BETA			
		Result ± 1s Uncertainty (x 10 ⁻¹⁵ µCi/mL)		Result > 3s	Result ± 1s Uncertainty (x 10 ⁻¹⁵ µCi/mL)		Result > 3s	
ARCO	10/1/2008	1.99 ± 0.31	7.36 ± 1.13	Y	42.90 ± 1.16	158.73 ± 4.29	Y	
	10/8/2008	1.65 ± 0.26	6.11 ± 0.96	Y	24.20 ± 0.88	89.54 ± 3.27	Y	
	10/15/2008	1.51 ± 0.20	5.59 ± 0.73	Y	27.20 ± 0.71	100.64 ± 2.62	Y	
	10/22/2008	1.95 ± 0.28	7.22 ± 1.03	Y	36.40 ± 1.01	134.68 ± 3.74	Y	
	10/29/2008	1.37 ± 0.28	5.07 ± 1.02	Y	37.20 ± 1.10	137.64 ± 4.07	Y	
	11/5/2008	1.43 ± 0.23	5.29 ± 0.85	Y	29.00 ± 0.92	107.30 ± 3.41	Y	
	11/12/2008	1.39 ± 0.25	5.14 ± 0.91	Y	24.40 ± 0.90	90.28 ± 3.32	Y	
	11/19/2008	2.36 ± 0.29	8.73 ± 1.06	Y	37.30 ± 1.02	138.01 ± 3.77	Y	
	11/26/2008	1.70 ± 0.22	6.29 ± 0.81	Y	33.70 ± 0.80	124.69 ± 2.95	Y	
	12/3/2008	1.54 ± 0.22	5.70 ± 0.83	Y	30.30 ± 0.77	112.11 ± 2.83	Y	
	12/10/2008	1.46 ± 0.21	5.40 ± 0.76	Y	35.10 ± 0.80	129.87 ± 2.96	Y	
	12/17/2008	1.43 ± 0.19	5.29 ± 0.71	Y	29.80 ± 0.73	110.26 ± 2.71	Y	
	12/24/2008	1.04 ± 0.18	3.85 ± 0.66	Y	23.10 ± 0.68	85.47 ± 2.50	Y	
	12/31/2008	0.58 ± 0.14	2.16 ± 0.52	Y	18.65 ± 0.59	69.00 ± 2.17	Y	
	ATOMIC CITY	10/1/2008	2.59 ± 0.31	9.58 ± 1.15	Y	49.00 ± 1.15	181.30 ± 4.26	Y
		10/8/2008	1.36 ± 0.27	5.03 ± 1.01	Y	28.00 ± 1.02	103.60 ± 3.77	Y
10/15/2008		1.15 ± 0.18	4.26 ± 0.66	Y	26.60 ± 0.70	98.42 ± 2.59	Y	
10/22/2008		1.69 ± 0.29	6.25 ± 1.08	Y	41.40 ± 1.15	153.18 ± 4.26	Y	
10/29/2008		2.03 ± 0.28	7.51 ± 1.05	Y	39.90 ± 1.04	147.63 ± 3.85	Y	
11/5/2008		1.36 ± 0.23	5.03 ± 0.85	Y	29.10 ± 0.93	107.67 ± 3.46	Y	
11/12/2008		1.01 ± 0.21	3.74 ± 0.78	Y	22.90 ± 0.82	84.73 ± 3.03	Y	
11/19/2008		1.72 ± 0.25	6.36 ± 0.94	Y	38.00 ± 1.02	140.60 ± 3.77	Y	
11/26/2008		1.44 ± 0.19	5.33 ± 0.72	Y	36.40 ± 0.77	134.68 ± 2.85	Y	
12/3/2008		1.69 ± 0.28	6.25 ± 1.03	Y	42.30 ± 1.03	156.51 ± 3.81	Y	
12/10/2008		1.30 ± 0.20	4.81 ± 0.73	Y	37.60 ± 0.82	139.12 ± 3.02	Y	
12/17/2008		1.63 ± 0.21	6.03 ± 0.77	Y	33.30 ± 0.78	123.21 ± 2.90	Y	
12/24/2008		0.77 ± 0.16	2.86 ± 0.59	Y	21.90 ± 0.65	81.03 ± 2.41	Y	
12/31/2008		0.79 ± 0.16	2.92 ± 0.60	Y	18.94 ± 0.63	70.07 ± 2.33	Y	
QA-1 (ATOMIC CITY)		10/1/2008	2.52 ± 0.35	9.32 ± 1.28	Y	54.60 ± 1.32	202.02 ± 4.88	Y
		10/8/2008	1.44 ± 0.29	5.33 ± 1.05	Y	23.70 ± 1.00	87.69 ± 3.70	Y
	10/15/2008	1.28 ± 0.19	4.74 ± 0.70	Y	28.10 ± 0.73	103.97 ± 2.70	Y	
	10/22/2008	2.50 ± 0.35	9.25 ± 1.30	Y	37.30 ± 1.18	138.01 ± 4.37	Y	
	10/29/2008	2.10 ± 0.30	7.77 ± 1.11	Y	38.40 ± 1.07	142.08 ± 3.96	Y	
	11/5/2008	2.22 ± 0.32	8.21 ± 1.19	Y	30.90 ± 1.14	114.33 ± 4.22	Y	
	11/12/2008	1.10 ± 0.22	4.07 ± 0.82	Y	22.40 ± 0.84	82.88 ± 3.10	Y	
	11/19/2008	1.90 ± 0.28	7.03 ± 1.04	Y	39.70 ± 1.10	146.89 ± 4.07	Y	
	11/26/2008	1.13 ± 0.18	4.18 ± 0.67	Y	29.00 ± 0.72	107.30 ± 2.68	Y	
	12/3/2008	1.48 ± 0.24	5.48 ± 0.88	Y	35.50 ± 0.87	131.35 ± 3.21	Y	
	12/10/2008	1.27 ± 0.21	4.70 ± 0.78	Y	37.70 ± 0.88	139.49 ± 3.27	Y	
	12/17/2008	1.24 ± 0.19	4.59 ± 0.70	Y	33.50 ± 0.79	123.95 ± 2.91	Y	
	12/24/2008	0.77 ± 0.16	2.85 ± 0.58	Y	20.40 ± 0.63	75.48 ± 2.32	Y	
	^a 12/31/2008	0.60 ± 0.82	2.21 ± 3.03	Y	29.40 ± 3.29	108.78 ± 12.16	Y	
	BLUE DOME	10/1/2008	1.96 ± 0.28	7.25 ± 1.03	Y	41.00 ± 1.05	151.70 ± 3.89	Y
		10/8/2008	1.58 ± 0.26	5.85 ± 0.97	Y	23.80 ± 0.90	88.06 ± 3.32	Y
10/15/2008		0.57 ± 0.14	2.12 ± 0.53	Y	21.60 ± 0.65	79.92 ± 2.39	Y	
10/22/2008		1.28 ± 0.26	4.74 ± 0.94	Y	33.70 ± 1.01	124.69 ± 3.74	Y	
10/29/2008		1.37 ± 0.27	5.07 ± 0.98	Y	29.50 ± 0.98	109.15 ± 3.63	Y	
11/5/2008		1.05 ± 0.21	3.89 ± 0.79	Y	30.90 ± 0.95	114.33 ± 3.53	Y	
11/12/2008		0.65 ± 0.20	2.39 ± 0.73	Y	20.40 ± 0.82	75.48 ± 3.02	Y	
11/19/2008		1.37 ± 0.24	5.07 ± 0.88	Y	33.70 ± 0.98	124.69 ± 3.63	Y	
11/26/2008		1.23 ± 0.19	4.55 ± 0.68	Y	31.30 ± 0.74	115.81 ± 2.72	Y	
12/3/2008		0.68 ± 0.22	2.50 ± 0.82	Y	25.50 ± 0.86	94.35 ± 3.17	Y	
12/10/2008	1.03 ± 0.16	3.81 ± 0.60	Y	26.80 ± 0.65	99.16 ± 2.42	Y		

TABLE C-1. Weekly Gross Alpha and Gross Beta Concentrations in Air

Sampling Group and Location	Sampling Date	GROSS ALPHA						GROSS BETA					
		Result ± 1s Uncertainty (x 10 ⁻¹⁵ µCi/mL)		Result ± 1s Uncertainty (x 10 ⁻¹¹ Bq/mL)		Result > 3s	Result ± 1s Uncertainty (x 10 ⁻¹⁵ µCi/mL)		Result ± 1s Uncertainty (x 10 ⁻¹¹ Bq/mL)		Result > 3s		
a	12/17/2008	1.12 ± 0.19	4.14 ± 0.69	Y	27.00 ± 0.75	99.90 ± 2.76	Y						
	12/24/2008	2.04 ± 0.20	7.55 ± 0.74	Y	16.20 ± 0.52	59.94 ± 1.94	Y						
	12/31/2008	0.57 ± 0.45	2.09 ± 1.68		17.03 ± 1.75	63.00 ± 6.47	Y						
	10/1/2008	1.65 ± 0.32	6.11 ± 1.18	Y	44.30 ± 1.28	163.91 ± 4.74	Y						
QA-2 (BLUE DOME)	10/8/2008	1.11 ± 0.25	4.11 ± 0.93	Y	24.80 ± 0.96	91.76 ± 3.56	Y						
	10/15/2008	0.66 ± 0.16	2.43 ± 0.58	Y	23.10 ± 0.70	85.47 ± 2.60	Y						
	10/22/2008	1.79 ± 0.26	6.62 ± 0.97	Y	33.20 ± 0.94	122.84 ± 3.49	Y						
	10/29/2008	1.29 ± 0.35	4.77 ± 1.29	Y	33.70 ± 1.33	124.69 ± 4.92	Y						
	11/5/2008	1.68 ± 0.27	6.22 ± 0.98	Y	26.90 ± 0.99	99.53 ± 3.64	Y						
	11/12/2008	1.70 ± 0.29	6.29 ± 1.08	Y	21.80 ± 0.98	80.66 ± 3.61	Y						
	11/19/2008	1.50 ± 0.22	5.55 ± 0.83	Y	32.70 ± 0.89	120.99 ± 3.29	Y						
	11/26/2008	1.48 ± 0.19	5.48 ± 0.69	Y	30.00 ± 0.69	111.00 ± 2.55	Y						
	12/3/2008	1.00 ± 0.17	3.70 ± 0.64	Y	22.90 ± 0.61	84.73 ± 2.26	Y						
	12/10/2008	1.30 ± 0.21	4.81 ± 0.76	Y	31.70 ± 0.80	117.29 ± 2.97	Y						
	12/17/2008	1.48 ± 0.19	5.48 ± 0.70	Y	26.50 ± 0.68	98.05 ± 2.53	Y						
	12/24/2008	0.93 ± 0.16	3.44 ± 0.59	Y	12.20 ± 0.51	45.14 ± 1.89	Y						
	12/31/2008	0.66 ± 0.13	2.46 ± 0.49	Y	13.86 ± 0.50	51.29 ± 1.83	Y						
	a	10/1/2008	5.04 ± 0.55	18.65 ± 2.04	Y	51.10 ± 1.64	189.07 ± 6.07	Y					
		10/8/2008	1.36 ± 0.27	5.03 ± 0.99	Y	24.90 ± 0.97	92.13 ± 3.60	Y					
		10/15/2008	0.80 ± 0.16	2.97 ± 0.60	Y	23.50 ± 0.68	86.95 ± 2.53	Y					
10/22/2008		1.59 ± 0.68	5.88 ± 2.50	Y	49.30 ± 2.60	182.41 ± 9.62	Y						
10/29/2008		1.61 ± 0.30	5.96 ± 1.11	Y	36.70 ± 1.13	135.79 ± 4.18	Y						
11/5/2008		1.61 ± 0.26	5.96 ± 0.96	Y	29.00 ± 1.00	107.30 ± 3.69	Y						
11/12/2008		0.74 ± 0.21	2.74 ± 0.79	Y	19.80 ± 0.85	73.26 ± 3.16	Y						
11/19/2008		1.48 ± 0.25	5.48 ± 0.91	Y	31.70 ± 0.98	117.29 ± 3.61	Y						
11/26/2008		1.33 ± 0.21	4.92 ± 0.77	Y	32.10 ± 0.81	118.77 ± 2.98	Y						
12/3/2008		1.20 ± 0.25	4.44 ± 0.91	Y	37.30 ± 0.95	138.01 ± 3.52	Y						
12/10/2008		1.18 ± 0.22	4.37 ± 0.81	Y	33.30 ± 0.90	123.21 ± 3.31	Y						
12/17/2008		1.30 ± 0.23	4.81 ± 0.84	Y	28.70 ± 0.88	106.19 ± 3.25	Y						
12/24/2008		0.78 ± 0.16	2.88 ± 0.61	Y	15.90 ± 0.60	58.83 ± 2.23	Y						
12/31/2008		0.35 ± 0.15	1.29 ± 0.55	Y	11.12 ± 0.60	41.13 ± 2.24	Y						
HOWE		10/1/2008	1.81 ± 0.25	6.70 ± 0.93	Y	41.60 ± 0.98	153.92 ± 3.61	Y					
		10/8/2008	1.77 ± 0.26	6.55 ± 0.97	Y	24.40 ± 0.88	90.28 ± 3.25	Y					
	10/15/2008	0.95 ± 0.17	3.52 ± 0.62	Y	28.90 ± 0.72	106.93 ± 2.67	Y						
	10/22/2008	2.36 ± 0.36	8.73 ± 1.33	Y	38.80 ± 1.25	143.56 ± 4.63	Y						
	10/29/2008	2.31 ± 0.36	8.55 ± 1.33	Y	43.10 ± 1.29	159.47 ± 4.77	Y						
	11/5/2008	1.36 ± 0.25	5.03 ± 0.93	Y	30.10 ± 1.03	111.37 ± 3.81	Y						
	11/12/2008	1.10 ± 0.23	4.07 ± 0.85	Y	26.10 ± 0.91	96.57 ± 3.35	Y						
	11/19/2008	1.69 ± 0.29	6.25 ± 1.07	Y	41.60 ± 1.20	153.92 ± 4.44	Y						
	11/26/2008	1.84 ± 0.22	6.81 ± 0.81	Y	36.80 ± 0.80	136.16 ± 2.96	Y						
	12/3/2008	1.40 ± 0.27	5.18 ± 1.00	Y	31.70 ± 0.95	117.29 ± 3.50	Y						
	12/10/2008	1.71 ± 0.19	6.33 ± 0.71	Y	36.80 ± 0.72	136.16 ± 2.68	Y						
	12/17/2008	1.47 ± 0.22	5.44 ± 0.80	Y	33.50 ± 0.85	123.95 ± 3.15	Y						
	12/24/2008	0.84 ± 0.15	3.10 ± 0.54	Y	23.70 ± 0.60	87.69 ± 2.22	Y						
	12/31/2008	0.64 ± 0.15	2.39 ± 0.56	Y	19.10 ± 0.62	70.68 ± 2.30	Y						
	MONTEVIEW	10/1/2008	2.32 ± 0.32	8.58 ± 1.18	Y	46.50 ± 1.19	172.05 ± 4.40	Y					
		10/8/2008	1.33 ± 0.25	4.92 ± 0.93	Y	26.70 ± 0.93	98.79 ± 3.45	Y					
10/15/2008		1.69 ± 0.31	6.25 ± 1.14	Y	24.60 ± 1.05	91.02 ± 3.89	Y						
10/22/2008		1.80 ± 0.28	6.66 ± 1.02	Y	34.40 ± 1.00	127.28 ± 3.69	Y						
10/29/2008		3.02 ± 0.36	11.17 ± 1.31	Y	38.50 ± 1.14	142.45 ± 4.22	Y						
11/5/2008		1.66 ± 0.23	6.14 ± 0.86	Y	29.20 ± 0.88	108.04 ± 3.26	Y						
11/12/2008		1.13 ± 0.22	4.18 ± 0.80	Y	24.20 ± 0.83	89.54 ± 3.07	Y						
11/19/2008		2.34 ± 0.27	8.66 ± 1.00	Y	34.90 ± 0.95	129.13 ± 3.50	Y						
11/26/2008		2.22 ± 0.23	8.21 ± 0.83	Y	33.40 ± 0.74	123.58 ± 2.74	Y						
12/3/2008		2.03 ± 0.24	7.51 ± 0.87	Y	43.60 ± 0.85	161.32 ± 3.13	Y						

TABLE C-1. Weekly Gross Alpha and Gross Beta Concentrations in Air

Sampling Group and Location	Sampling Date	GROSS ALPHA				GROSS BETA			
		Result ± 1s Uncertainty (x 10 ⁻¹⁵ µCi/mL)		Result ± 1s Uncertainty (x 10 ⁻¹¹ Bq/mL)		Result ± 1s Uncertainty (x 10 ⁻¹⁵ µCi/mL)		Result ± 1s Uncertainty (x 10 ⁻¹¹ Bq/mL)	
	12/10/2008	1.40 ± 0.17	5.18 ± 0.63	Y	33.20 ± 0.67	122.84 ± 2.46	Y		
	12/17/2008	1.65 ± 0.20	6.11 ± 0.74	Y	29.80 ± 0.71	110.26 ± 2.64	Y		
	12/24/2008	0.77 ± 0.13	2.85 ± 0.48	Y	13.00 ± 0.45	48.10 ± 1.67	Y		
	12/31/2008	0.48 ± 0.12	1.76 ± 0.46	Y	16.70 ± 0.53	61.80 ± 1.97	Y		
MUD LAKE	10/1/2008	2.60 ± 0.32	9.62 ± 1.18	Y	52.90 ± 1.20	195.73 ± 4.44	Y		
	10/8/2008	1.29 ± 0.25	4.77 ± 0.91	Y	27.40 ± 0.94	101.38 ± 3.46	Y		
	10/15/2008	0.82 ± 0.15	3.02 ± 0.56	Y	26.20 ± 0.66	96.94 ± 2.45	Y		
	10/22/2008	2.52 ± 0.31	9.32 ± 1.15	Y	42.60 ± 1.09	157.62 ± 4.03	Y		
	10/29/2008	2.32 ± 0.31	8.58 ± 1.15	Y	48.70 ± 1.17	180.19 ± 4.33	Y		
	11/5/2008	1.41 ± 0.23	5.22 ± 0.85	Y	28.90 ± 0.91	106.93 ± 3.38	Y		
	11/12/2008	0.96 ± 0.21	3.54 ± 0.78	Y	23.00 ± 0.84	85.10 ± 3.09	Y		
	11/19/2008	2.01 ± 0.29	7.44 ± 1.06	Y	45.00 ± 1.15	166.50 ± 4.26	Y		
	11/26/2008	2.20 ± 0.23	8.14 ± 0.83	Y	37.30 ± 0.77	138.01 ± 2.86	Y		
	12/3/2008	1.79 ± 0.29	6.62 ± 1.07	Y	51.80 ± 1.12	191.66 ± 4.14	Y		
	12/10/2008	2.03 ± 0.22	7.51 ± 0.83	Y	43.10 ± 0.84	159.47 ± 3.09	Y		
	12/17/2008	1.69 ± 0.21	6.25 ± 0.78	Y	35.60 ± 0.80	131.72 ± 2.95	Y		
	12/24/2008	0.63 ± 0.14	2.34 ± 0.51	Y	24.60 ± 0.62	91.02 ± 2.31	Y		
	12/31/2008	0.49 ± 0.13	1.83 ± 0.47	Y	18.34 ± 0.56	67.87 ± 2.08	Y		
DISTANT									
BLACKFOOT CMS	10/1/2008	2.68 ± 0.41	9.92 ± 1.53	Y	46.80 ± 1.47	173.16 ± 5.44	Y		
	10/8/2008	1.78 ± 0.34	6.59 ± 1.26	Y	27.00 ± 1.17	99.90 ± 4.33	Y		
	10/15/2008	1.41 ± 0.21	5.22 ± 0.79	Y	26.90 ± 0.79	99.53 ± 2.91	Y		
	10/22/2008	2.00 ± 0.38	7.40 ± 1.40	Y	40.50 ± 1.39	149.85 ± 5.14	Y		
	10/29/2008	2.29 ± 0.32	8.47 ± 1.20	Y	35.90 ± 1.10	132.83 ± 4.07	Y		
	11/5/2008	2.38 ± 0.30	8.81 ± 1.12	Y	31.40 ± 1.05	116.18 ± 3.89	Y		
	11/12/2008	1.01 ± 0.24	3.74 ± 0.88	Y	18.80 ± 0.87	69.56 ± 3.23	Y		
	11/19/2008	1.81 ± 0.29	6.70 ± 1.05	Y	35.80 ± 1.10	132.46 ± 4.07	Y		
	11/26/2008	2.26 ± 0.25	8.36 ± 0.93	Y	36.40 ± 0.84	134.68 ± 3.12	Y		
	12/3/2008	1.73 ± 0.25	6.40 ± 0.94	Y	44.00 ± 0.95	162.80 ± 3.52	Y		
	12/10/2008	0.90 ± 0.18	3.34 ± 0.68	Y	32.80 ± 0.81	121.36 ± 3.00	Y		
	12/17/2008	1.54 ± 0.19	5.70 ± 0.68	Y	30.90 ± 0.69	114.33 ± 2.55	Y		
	12/24/2008	0.80 ± 0.14	2.95 ± 0.53	Y	18.10 ± 0.54	66.97 ± 2.01	Y		
	12/31/2008	0.48 ± 0.11	1.79 ± 0.42	Y	13.25 ± 0.45	49.02 ± 1.68	Y		
CRATERS OF THE MOON	10/1/2008	2.44 ± 0.35	9.03 ± 1.28	Y	41.40 ± 1.22	153.18 ± 4.51	Y		
	10/8/2008	0.84 ± 0.24	3.12 ± 0.90	Y	21.90 ± 0.96	81.03 ± 3.54	Y		
	10/15/2008	1.15 ± 0.20	4.26 ± 0.73	Y	26.10 ± 0.77	96.57 ± 2.83	Y		
	10/22/2008	1.88 ± 0.31	6.96 ± 1.15	Y	32.80 ± 1.09	121.36 ± 4.03	Y		
	10/29/2008	1.24 ± 0.27	4.59 ± 1.01	Y	33.90 ± 1.08	125.43 ± 4.00	Y		
	11/5/2008	1.53 ± 0.23	5.66 ± 0.84	Y	23.80 ± 0.84	88.06 ± 3.10	Y		
	11/12/2008	1.02 ± 0.21	3.77 ± 0.79	Y	20.30 ± 0.80	75.11 ± 2.96	Y		
	11/19/2008	1.70 ± 0.24	6.29 ± 0.89	Y	30.90 ± 0.91	114.33 ± 3.35	Y		
	11/26/2008	0.66 ± 0.17	2.43 ± 0.61	Y	30.80 ± 0.79	113.96 ± 2.91	Y		
	12/3/2008	1.82 ± 0.24	6.73 ± 0.87	Y	36.90 ± 0.82	136.53 ± 3.05	Y		
	12/10/2008	1.10 ± 0.18	4.07 ± 0.67	Y	28.70 ± 0.73	106.19 ± 2.69	Y		
	12/17/2008	1.14 ± 0.18	4.22 ± 0.65	Y	28.10 ± 0.70	103.97 ± 2.60	Y		
	12/24/2008	0.80 ± 0.16	2.96 ± 0.58	Y	16.20 ± 0.57	59.94 ± 2.11	Y		
	12/31/2008	0.39 ± 0.12	1.44 ± 0.45	Y	14.92 ± 0.53	55.20 ± 1.96	Y		
DUBOIS	10/1/2008	2.64 ± 0.30	9.77 ± 1.11	Y	43.30 ± 1.05	160.21 ± 3.89	Y		
	10/8/2008	1.21 ± 0.25	4.48 ± 0.94	Y	25.00 ± 0.95	92.50 ± 3.51	Y		
	10/15/2008	1.23 ± 0.16	4.55 ± 0.61	Y	23.70 ± 0.60	87.69 ± 2.23	Y		
	10/22/2008	1.45 ± 0.25	5.37 ± 0.92	Y	37.00 ± 0.99	136.90 ± 3.65	Y		
	10/29/2008	1.91 ± 0.28	7.07 ± 1.03	Y	34.20 ± 0.99	126.54 ± 3.66	Y		
	11/5/2008	1.68 ± 0.25	6.22 ± 0.92	Y	27.60 ± 0.92	102.12 ± 3.42	Y		
	11/12/2008	0.95 ± 0.22	3.50 ± 0.81	Y	19.70 ± 0.83	72.89 ± 3.06	Y		
	11/19/2008	1.50 ± 0.26	5.55 ± 0.95	Y	30.20 ± 1.00	111.74 ± 3.70	Y		

TABLE C-1. Weekly Gross Alpha and Gross Beta Concentrations in Air

Sampling Group and Location	Sampling Date	GROSS ALPHA				GROSS BETA			
		Result ± 1s Uncertainty (x 10 ⁻¹⁵ µCi/mL)		Result ± 1s Uncertainty (x 10 ⁻¹¹ Bq/mL)		Result ± 1s Uncertainty (x 10 ⁻¹⁵ µCi/mL)		Result ± 1s Uncertainty (x 10 ⁻¹¹ Bq/mL)	
				Result > 3s					Result > 3s
	11/26/2008	1.38 ± 0.18	5.11 ± 0.66	Y	29.10 ± 0.67	107.67 ± 2.46	Y		
	12/3/2008	1.79 ± 0.28	6.62 ± 1.03	Y	42.80 ± 1.02	158.36 ± 3.77	Y		
	12/10/2008	1.10 ± 0.20	4.07 ± 0.74	Y	30.50 ± 0.81	112.85 ± 3.00	Y		
	12/17/2008	1.89 ± 0.23	6.99 ± 0.86	Y	31.30 ± 0.81	115.81 ± 3.00	Y		
	12/24/2008	1.23 ± 0.16	4.55 ± 0.61	Y	13.10 ± 0.49	48.47 ± 1.81	Y		
	12/31/2008	0.69 ± 0.16	2.56 ± 0.60	Y	14.37 ± 0.60	53.15 ± 2.21	Y		
IDAHO FALLS	10/1/2008	2.58 ± 0.37	9.55 ± 1.36	Y	47.90 ± 1.33	177.23 ± 4.92	Y		
	10/8/2008	2.48 ± 0.35	9.18 ± 1.28	Y	25.70 ± 1.06	95.09 ± 3.92	Y		
	10/15/2008	1.24 ± 0.20	4.59 ± 0.74	Y	28.30 ± 0.78	104.71 ± 2.87	Y		
	10/22/2008	2.74 ± 0.34	10.14 ± 1.25	Y	39.00 ± 1.13	144.30 ± 4.18	Y		
	10/29/2008	3.02 ± 0.37	11.17 ± 1.38	Y	39.70 ± 1.20	146.89 ± 4.44	Y		
	11/5/2008	2.59 ± 0.36	9.58 ± 1.32	Y	32.90 ± 1.23	121.73 ± 4.55	Y		
	11/12/2008	1.26 ± 0.24	4.66 ± 0.87	Y	17.30 ± 0.80	64.01 ± 2.95	Y		
	11/19/2008	1.50 ± 0.26	5.55 ± 0.97	Y	37.10 ± 1.09	137.27 ± 4.03	Y		
	11/26/2008	1.73 ± 0.23	6.40 ± 0.84	Y	34.00 ± 0.82	125.80 ± 3.05	Y		
	12/3/2008	1.54 ± 0.24	5.70 ± 0.90	Y	44.30 ± 0.94	163.91 ± 3.49	Y		
	12/10/2008	1.44 ± 0.21	5.33 ± 0.77	Y	34.30 ± 0.81	126.91 ± 3.00	Y		
	12/17/2008	1.34 ± 0.21	4.96 ± 0.77	Y	32.00 ± 0.83	118.40 ± 3.08	Y		
	12/24/2008	1.29 ± 0.19	4.77 ± 0.70	Y	19.50 ± 0.63	72.15 ± 2.31	Y		
	12/31/2008	0.39 ± 0.13	1.46 ± 0.48	Y	13.77 ± 0.54	50.94 ± 2.02	Y		
JACKSON	10/1/2008	2.20 ± 0.34	8.14 ± 1.26	Y	42.60 ± 1.25	157.62 ± 4.63	Y		
a	10/8/2008								
a	10/15/2008								
	10/22/2008	1.62 ± 0.28	5.99 ± 1.03	Y	30.40 ± 1.00	112.48 ± 3.70	Y		
	10/29/2008	2.33 ± 0.32	8.62 ± 1.18	Y	33.30 ± 1.06	123.21 ± 3.92	Y		
	11/5/2008	1.58 ± 0.27	5.85 ± 1.00	Y	30.30 ± 1.06	112.11 ± 3.92	Y		
	11/12/2008	0.76 ± 0.22	2.82 ± 0.81	Y	17.00 ± 0.84	62.90 ± 3.09	Y		
	11/19/2008	1.37 ± 0.25	5.07 ± 0.91	Y	26.50 ± 0.95	98.05 ± 3.51	Y		
	11/26/2008	1.67 ± 0.22	6.18 ± 0.83	Y	35.10 ± 0.83	129.87 ± 3.07	Y		
a	12/3/2008	3.57 ± 0.70	13.21 ± 2.60	Y	98.90 ± 13.23	365.93 ± 9.73	Y		
	12/10/2008	1.02 ± 0.14	3.77 ± 0.50	Y	20.40 ± 0.50	75.48 ± 1.83	Y		
	12/17/2008	1.06 ± 0.18	3.92 ± 0.65	Y	18.40 ± 0.63	68.08 ± 2.32	Y		
	12/24/2008	0.90 ± 0.18	3.32 ± 0.65	Y	17.00 ± 0.63	62.90 ± 2.33	Y		
	12/31/2008	0.44 ± 0.12	1.64 ± 0.45	Y	12.95 ± 0.49	47.91 ± 1.82	Y		
REXBURG CMS	10/1/2008	2.51 ± 0.31	9.29 ± 1.14	Y	47.80 ± 1.13	176.86 ± 4.18	Y		
	10/8/2008	1.46 ± 0.25	5.40 ± 0.93	Y	26.80 ± 0.91	99.16 ± 3.36	Y		
	10/15/2008	1.21 ± 0.16	4.48 ± 0.60	Y	22.00 ± 0.58	81.40 ± 2.16	Y		
	10/22/2008	1.70 ± 0.27	6.29 ± 0.98	Y	37.80 ± 1.02	139.86 ± 3.77	Y		
	10/29/2008	2.03 ± 0.30	7.51 ± 1.11	Y	40.60 ± 1.10	150.22 ± 4.07	Y		
	11/5/2008	2.39 ± 0.26	8.84 ± 0.97	Y	30.50 ± 0.89	112.85 ± 3.28	Y		
	11/12/2008	0.68 ± 0.18	2.51 ± 0.65	Y	13.00 ± 0.65	48.10 ± 2.41	Y		
	11/19/2008	1.87 ± 0.26	6.92 ± 0.96	Y	33.00 ± 0.96	122.10 ± 3.56	Y		
	11/26/2008	1.46 ± 0.20	5.40 ± 0.72	Y	32.20 ± 0.74	119.14 ± 2.74	Y		
	12/3/2008	1.12 ± 0.23	4.14 ± 0.84	Y	34.40 ± 0.87	127.28 ± 3.23	Y		
	12/10/2008	1.83 ± 0.23	6.77 ± 0.84	Y	32.00 ± 0.79	118.40 ± 2.94	Y		
	12/17/2008	1.57 ± 0.19	5.81 ± 0.70	Y	27.00 ± 0.67	99.90 ± 2.48	Y		
	12/24/2008	0.51 ± 0.13	1.89 ± 0.48	Y	16.70 ± 0.54	61.79 ± 2.01	Y		
	12/31/2008	0.59 ± 0.13	2.20 ± 0.49	Y	13.66 ± 0.50	50.54 ± 1.87	Y		
INL SITE									
EFS	10/1/2008	2.20 ± 0.35	8.14 ± 1.28	Y	52.20 ± 1.36	193.14 ± 5.03	Y		
	10/8/2008	1.40 ± 0.27	5.18 ± 1.00	Y	26.50 ± 0.99	98.05 ± 3.67	Y		
	10/15/2008	1.60 ± 0.21	5.92 ± 0.77	Y	28.00 ± 0.74	103.60 ± 2.72	Y		
	10/22/2008	1.91 ± 0.29	7.07 ± 1.09	Y	37.90 ± 1.08	140.23 ± 4.00	Y		
	10/29/2008	1.62 ± 0.29	5.99 ± 1.07	Y	43.20 ± 1.16	159.84 ± 4.29	Y		
	11/5/2008	1.93 ± 0.27	7.14 ± 1.01	Y	30.60 ± 1.00	113.22 ± 3.70	Y		

TABLE C-1. Weekly Gross Alpha and Gross Beta Concentrations in Air

Sampling Group and Location	Sampling Date	GROSS ALPHA						GROSS BETA					
		Result ± 1s Uncertainty (x 10 ⁻¹⁵ µCi/mL)		Result ± 1s Uncertainty (x 10 ⁻¹¹ Bq/mL)		Result > 3s	Result ± 1s Uncertainty (x 10 ⁻¹⁵ µCi/mL)		Result ± 1s Uncertainty (x 10 ⁻¹¹ Bq/mL)		Result > 3s		
	11/12/2008	1.56	± 0.26	5.77	± 0.95	Y	25.20	± 0.91	93.24	± 3.36	Y		
	11/19/2008	1.81	± 0.27	6.70	± 0.99	Y	42.80	± 1.09	158.36	± 4.03	Y		
	11/26/2008	1.27	± 0.20	4.70	± 0.74	Y	37.90	± 0.84	140.23	± 3.10	Y		
	12/3/2008	1.15	± 0.24	4.26	± 0.90	Y	39.70	± 0.98	146.89	± 3.61	Y		
	12/10/2008	1.78	± 0.26	6.59	± 0.97	Y	40.90	± 1.01	151.33	± 3.74	Y		
	12/17/2008	1.39	± 0.19	5.14	± 0.69	Y	31.70	± 0.73	117.29	± 2.71	Y		
	12/24/2008	1.06	± 0.17	3.92	± 0.64	Y	27.00	± 0.68	99.90	± 2.52	Y		
	12/31/2008	0.72	± 0.15	2.67	± 0.55	Y	19.60	± 0.60	72.51	± 2.21	Y		
MAIN GATE	10/1/2008	2.46	± 0.44	9.10	± 1.62	Y	50.60	± 1.62	187.22	± 5.99	Y		
	10/8/2008	1.38	± 0.27	5.11	± 0.98	Y	26.60	± 0.98	98.42	± 3.61	Y		
	10/15/2008	1.04	± 0.18	3.85	± 0.66	Y	27.80	± 0.74	102.86	± 2.73	Y		
	10/22/2008	2.26	± 0.31	8.36	± 1.14	Y	37.50	± 1.07	138.75	± 3.96	Y		
	10/29/2008	1.79	± 0.33	6.62	± 1.23	Y	44.10	± 1.29	163.17	± 4.77	Y		
	11/5/2008	2.41	± 0.43	8.92	± 1.59	Y	35.70	± 1.58	132.09	± 5.85	Y		
	11/12/2008	1.43	± 0.23	5.29	± 0.87	Y	21.10	± 0.81	78.07	± 2.98	Y		
	11/19/2008	1.56	± 0.24	5.77	± 0.90	Y	38.40	± 1.00	142.08	± 3.70	Y		
	11/26/2008	1.43	± 0.21	5.29	± 0.77	Y	34.90	± 0.82	129.13	± 3.02	Y		
	12/3/2008	1.62	± 0.24	5.99	± 0.90	Y	36.30	± 0.87	134.31	± 3.23	Y		
	12/10/2008	2.26	± 0.37	8.36	± 1.36	Y	45.70	± 1.36	169.09	± 5.03	Y		
	12/17/2008	1.94	± 0.34	7.18	± 1.25	Y	40.20	± 1.29	148.74	± 4.77	Y		
	12/24/2008	0.91	± 0.19	3.36	± 0.70	Y	18.60	± 0.70	68.82	± 2.57	Y		
	12/31/2008	0.57	± 0.14	2.10	± 0.51	Y	20.03	± 0.60	74.09	± 2.22	Y		
VAN BUREN GATE	10/1/2008	3.52	± 0.46	13.02	± 1.69	Y	46.10	± 1.48	170.57	± 5.48	Y		
	10/8/2008	1.51	± 0.28	5.59	± 1.02	Y	22.70	± 0.95	83.99	± 3.50	Y		
	10/15/2008	0.94	± 0.17	3.47	± 0.63	Y	29.10	± 0.73	107.67	± 2.72	Y		
	10/22/2008	2.20	± 0.32	8.14	± 1.18	Y	42.10	± 1.17	155.77	± 4.33	Y		
	10/29/2008	1.66	± 0.30	6.14	± 1.10	Y	41.60	± 1.16	153.92	± 4.29	Y		
	11/5/2008	2.12	± 0.28	7.84	± 1.02	Y	33.80	± 1.02	125.06	± 3.77	Y		
	11/12/2008	0.99	± 0.22	3.65	± 0.82	Y	23.60	± 0.87	87.32	± 3.23	Y		
	11/19/2008	1.61	± 0.26	5.96	± 0.94	Y	40.00	± 1.06	148.00	± 3.92	Y		
	11/26/2008	1.31	± 0.19	4.85	± 0.72	Y	36.30	± 0.80	134.31	± 2.94	Y		
	12/3/2008	1.00	± 0.23	3.70	± 0.87	Y	43.90	± 1.00	162.43	± 3.70	Y		
	12/10/2008	1.42	± 0.22	5.25	± 0.83	Y	35.50	± 0.89	131.35	± 3.29	Y		
	12/17/2008	1.12	± 0.20	4.14	± 0.74	Y	30.70	± 0.84	113.59	± 3.10	Y		
a	12/24/2008		±		±			±		±			
	12/31/2008	0.49	± 0.12	1.80	± 0.46	Y	18.87	± 0.55	69.80	± 2.03	Y		

a. Invalid Sample Result

TABLE C-2. Weekly Iodine-131 Activity in Air.

Sampling Group and Location	Sampling Date	Result ± 1s Uncertainty (x 10 ⁻¹⁵ µCi/mL)			Result ± 1s Uncertainty (x 10 ⁻¹¹ Bq/mL)			Result > 3s
BOUNDARY								
ARCO	10/01/2008	-0.87	±	1.61	-3.23	±	5.95	
	10/08/2008	0.35	±	1.40	1.29	±	5.18	
	10/15/2008	0.21	±	1.36	0.76	±	5.04	
	10/22/2008	-1.37	±	1.39	-5.06	±	5.14	
	10/29/2008	-0.03	±	1.63	-0.10	±	6.05	
	11/05/2008	0.39	±	1.33	1.45	±	4.92	
	11/12/2008	-1.23	±	1.35	-4.53	±	4.99	
	11/19/2008	0.24	±	1.47	0.87	±	5.44	
	11/26/2008	-2.29	±	1.51	-8.49	±	5.57	
	12/03/2008	-0.98	±	1.41	-3.64	±	5.21	
	12/10/2008	2.59	±	1.20	9.58	±	4.43	
	12/17/2008	-0.67	±	1.36	-2.47	±	5.03	
	12/24/2008	-0.41	±	1.42	-1.50	±	5.27	
	12/31/2008	-0.86	±	1.34	-3.19	±	4.97	
ATOMIC CITY	10/01/2008	-0.79	±	1.45	-2.91	±	5.36	
	10/08/2008	0.41	±	1.63	1.50	±	6.01	
	10/15/2008	0.21	±	1.36	0.76	±	5.03	
	10/22/2008	-1.55	±	1.57	-5.74	±	5.83	
	10/29/2008	-0.02	±	1.44	-0.09	±	5.34	
	11/05/2008	0.40	±	1.35	1.47	±	5.01	
	11/12/2008	-1.10	±	1.21	-4.08	±	4.49	
	11/19/2008	0.23	±	1.44	0.85	±	5.32	
	11/26/2008	-2.04	±	1.34	-7.55	±	4.95	
	12/03/2008	-1.29	±	1.84	-4.76	±	6.81	
	12/10/2008	2.53	±	1.17	9.37	±	4.34	
	12/17/2008	-0.69	±	1.41	-2.56	±	5.22	
	12/24/2008	-0.39	±	1.38	-1.46	±	5.12	
	12/31/2008	-0.96	±	1.49	-3.55	±	5.53	
QA-1	10/01/2008	-0.93	±	1.71	-3.43	±	6.32	
	10/08/2008	0.43	±	1.71	1.58	±	6.31	
	10/15/2008	0.21	±	1.40	0.79	±	5.19	
	10/22/2008	-1.73	±	1.75	-6.39	±	6.49	
	10/29/2008	-0.02	±	1.55	-0.09	±	5.72	
	11/05/2008	0.52	±	1.77	1.93	±	6.56	
	11/12/2008	-1.15	±	1.27	-4.27	±	4.69	
	11/19/2008	0.26	±	1.59	0.95	±	5.90	
	11/26/2008	-2.16	±	1.42	-8.00	±	5.25	
	12/03/2008	-1.09	±	1.56	-4.03	±	5.76	
	12/10/2008	2.92	±	1.35	10.79	±	4.99	
	12/17/2008	-0.69	±	1.41	-2.57	±	5.23	
	12/24/2008	-0.39	±	1.36	-1.43	±	5.03	
	a	12/31/2008	-8.25	±	12.86	-30.53	±	47.58
BLUE DOME	10/01/2008	-1.26	±	1.30	-4.65	±	4.82	
	10/08/2008	1.85	±	1.32	6.85	±	4.90	
	10/15/2008	1.43	±	1.28	5.30	±	4.72	
	10/22/2008	0.14	±	1.34	0.53	±	4.96	
	10/29/2008	-2.25	±	1.50	-8.32	±	5.55	
	11/05/2008	-2.06	±	1.23	-7.63	±	4.55	

TABLE C-2. Weekly Iodine-131 Activity in Air.

Sampling Group and Location	Sampling Date	Result ± 1s Uncertainty (x 10 ⁻¹⁵ µCi/mL)			Result ± 1s Uncertainty (x 10 ⁻¹¹ Bq/mL)			Result > 3s
BOUNDARY								
	11/12/2008	-1.84	±	1.17	-6.80	±	4.32	
	11/19/2008	0.22	±	1.32	0.81	±	4.89	
	11/26/2008	0.44	±	1.27	1.63	±	4.69	
	12/03/2008	-0.11	±	1.75	-0.40	±	6.47	
	12/10/2008	0.80	±	1.66	2.95	±	6.14	
	12/17/2008	0.39	±	1.41	1.45	±	5.20	
	12/24/2008	-0.16	±	1.07	-0.61	±	3.95	
a	12/31/2008	-2.13	±	6.34	-7.90	±	23.44	
QA-2	10/01/2008	-1.65	±	1.70	-6.09	±	6.30	
	10/08/2008	2.01	±	1.44	7.45	±	5.33	
	10/15/2008	1.57	±	1.40	5.83	±	5.19	
	10/22/2008	0.13	±	1.22	0.48	±	4.51	
	10/29/2008	-3.32	±	2.22	-12.28	±	8.20	
	11/05/2008	-2.32	±	1.38	-8.59	±	5.12	
	11/12/2008	-2.28	±	1.45	-8.43	±	5.35	
	11/19/2008	0.19	±	1.15	0.71	±	4.25	
	11/26/2008	0.40	±	1.16	1.50	±	4.29	
	12/03/2008	-0.07	±	1.07	-0.25	±	3.98	
	12/10/2008	1.01	±	2.10	3.73	±	7.77	
	12/17/2008	0.34	±	1.23	1.27	±	4.53	
	12/24/2008	-0.19	±	1.22	-0.69	±	4.50	
	12/31/2008	-0.39	±	1.16	-1.45	±	4.30	
FAA TOWER	10/01/2008	-2.23	±	2.31	-8.24	±	8.53	
	10/08/2008	2.05	±	1.46	7.57	±	5.42	
	10/15/2008	1.49	±	1.32	5.50	±	4.89	
a	10/22/2008	0.48	±	4.48	1.77	±	16.58	
	10/29/2008	-2.49	±	1.66	-9.22	±	6.16	
	11/05/2008	-2.28	±	1.36	-8.45	±	5.04	
	11/12/2008	-1.99	±	1.26	-7.37	±	4.68	
	11/19/2008	0.23	±	1.36	0.84	±	5.04	
	11/26/2008	0.51	±	1.45	1.87	±	5.38	
	12/03/2008	-0.10	±	1.62	-0.37	±	5.98	
	12/10/2008	1.17	±	2.44	4.34	±	9.05	
	12/17/2008	0.49	±	1.77	1.83	±	6.56	
	12/24/2008	-0.21	±	1.36	-0.78	±	5.04	
	12/31/2008	-0.60	±	1.77	-2.20	±	6.54	
HOWE	10/01/2008	-1.11	±	1.15	-4.10	±	4.24	
	10/08/2008	1.77	±	1.26	6.54	±	4.68	
	10/15/2008	1.41	±	1.26	5.23	±	4.65	
	10/22/2008	0.18	±	1.72	0.68	±	6.37	
	10/29/2008	-2.80	±	1.87	-10.35	±	6.91	
	11/05/2008	-2.36	±	1.41	-8.74	±	5.21	
	11/12/2008	-1.91	±	1.21	-7.08	±	4.49	
	11/19/2008	0.27	±	1.60	0.99	±	5.93	
	11/26/2008	0.45	±	1.30	1.67	±	4.80	
	12/03/2008	-0.11	±	1.79	-0.41	±	6.64	
	12/10/2008	0.75	±	1.56	2.77	±	5.78	
	12/17/2008	0.42	±	1.51	1.56	±	5.57	

TABLE C-2. Weekly Iodine-131 Activity in Air.

Sampling Group and Location	Sampling Date	Result ± 1s Uncertainty (x 10 ⁻¹⁵ µCi/mL)		Result ± 1s Uncertainty (x 10 ⁻¹¹ Bq/mL)		Result > 3s
BOUNDARY						
	12/24/2008	-0.16	± 1.05	-0.60	± 3.90	
	12/31/2008	-0.46	± 1.38	-1.72	± 5.09	
MONTEVIEW	10/01/2008	-1.42	± 1.47	-5.25	± 5.43	
	10/08/2008	1.85	± 1.33	6.86	± 4.90	
	10/15/2008	2.85	± 2.54	10.54	± 9.38	
	10/22/2008	0.14	± 1.30	0.51	± 4.82	
	10/29/2008	-2.43	± 1.62	-9.00	± 6.01	
	11/05/2008	-1.88	± 1.12	-6.95	± 4.14	
	11/12/2008	-1.74	± 1.10	-6.44	± 4.09	
	11/19/2008	0.20	± 1.22	0.75	± 4.52	
	11/26/2008	0.42	± 1.22	1.57	± 4.51	
	12/03/2008	-0.07	± 1.17	-0.27	± 4.31	
	12/10/2008	0.70	± 1.46	2.59	± 5.39	
	12/17/2008	0.34	± 1.21	1.25	± 4.49	
	12/24/2008	-0.15	± 0.97	-0.55	± 3.59	
	12/31/2008	-0.39	± 1.17	-1.45	± 4.32	
MUD LAKE	10/01/2008	-1.33	± 1.38	-4.93	± 5.11	
	10/08/2008	1.84	± 1.31	6.80	± 4.86	
	10/15/2008	1.31	± 1.17	4.85	± 4.32	
	10/22/2008	0.14	± 1.32	0.52	± 4.89	
	10/29/2008	-2.22	± 1.48	-8.20	± 5.47	
	11/05/2008	-2.00	± 1.19	-7.40	± 4.41	
	11/12/2008	-1.80	± 1.14	-6.67	± 4.23	
	11/19/2008	0.24	± 1.44	0.88	± 5.32	
	11/26/2008	0.42	± 1.20	1.55	± 4.45	
	12/03/2008	-0.11	± 1.69	-0.39	± 6.25	
	12/10/2008	0.86	± 1.78	3.16	± 6.59	
	12/17/2008	0.36	± 1.29	1.33	± 4.76	
	12/24/2008	-0.17	± 1.10	-0.63	± 4.07	
	12/31/2008	-0.40	± 1.20	-1.49	± 4.43	
DISTANT						
BLACKFOOT CMS	10/01/2008	-1.20	± 2.21	-4.44	± 8.19	
	10/08/2008	0.50	± 2.02	1.86	± 7.47	
	10/15/2008	0.25	± 1.64	0.92	± 6.08	
	10/22/2008	-2.11	± 2.15	-7.83	± 7.95	
	10/29/2008	-0.03	± 1.67	-0.10	± 6.20	
	11/05/2008	0.45	± 1.55	1.68	± 5.72	
	11/12/2008	-1.32	± 1.45	-4.88	± 5.37	
	11/19/2008	0.27	± 1.69	1.01	± 6.27	
	11/26/2008	-2.39	± 1.57	-8.83	± 5.79	
	12/03/2008	-1.09	± 1.55	-4.02	± 5.75	
	12/10/2008	2.79	± 1.29	10.33	± 4.78	
	12/17/2008	-0.59	± 1.19	-2.17	± 4.41	
	12/24/2008	-0.33	± 1.16	-1.23	± 4.31	
	12/31/2008	-0.71	± 1.10	-2.61	± 4.07	
CRATERS	10/01/2008	-0.97	± 1.79	-3.59	± 6.61	
	10/08/2008	0.41	± 1.65	1.53	± 6.12	
	10/15/2008	0.24	± 1.60	0.90	± 5.93	

TABLE C-2. Weekly Iodine-131 Activity in Air.

Sampling Group and Location	Sampling Date	Result ± 1s Uncertainty (x 10 ⁻¹⁵ µCi/mL)			Result ± 1s Uncertainty (x 10 ⁻¹¹ Bq/mL)			Result > 3s
BOUNDARY								
	10/22/2008	-1.63	±	1.66	-6.05	±	6.14	
	10/29/2008	-0.03	±	1.67	-0.10	±	6.19	
	11/05/2008	0.37	±	1.27	1.38	±	4.70	
	11/12/2008	-1.13	±	1.24	-4.18	±	4.59	
	11/19/2008	0.22	±	1.35	0.80	±	5.01	
	11/26/2008	-2.40	±	1.58	-8.89	±	5.83	
	12/03/2008	-0.97	±	1.38	-3.58	±	5.12	
	12/10/2008	2.53	±	1.17	9.38	±	4.34	
	12/17/2008	-0.65	±	1.32	-2.40	±	4.90	
	12/24/2008	-0.38	±	1.35	-1.42	±	4.98	
	12/31/2008	-0.84	±	1.31	-3.11	±	4.84	
DUBOIS								
	10/01/2008	-1.21	±	1.26	-4.49	±	4.65	
	10/08/2008	1.96	±	1.40	7.26	±	5.19	
	10/15/2008	1.19	±	1.06	4.41	±	3.93	
	10/22/2008	0.13	±	1.23	0.48	±	4.53	
	10/29/2008	-2.09	±	1.40	-7.74	±	5.17	
	11/05/2008	-2.08	±	1.24	-7.70	±	4.59	
	11/12/2008	-1.90	±	1.21	-7.04	±	4.47	
	11/19/2008	0.24	±	1.45	0.89	±	5.37	
	11/26/2008	0.39	±	1.12	1.44	±	4.14	
	12/03/2008	-0.10	±	1.65	-0.38	±	6.09	
	12/10/2008	1.06	±	2.20	3.91	±	8.15	
	12/17/2008	0.41	±	1.45	1.50	±	5.38	
	12/24/2008	-0.17	±	1.09	-0.62	±	4.02	
	12/31/2008	-0.51	±	1.53	-1.90	±	5.65	
IDAHO FALLS								
	10/01/2008	-1.68	±	1.74	-6.22	±	6.44	
	10/08/2008	2.28	±	1.63	8.44	±	6.04	
	10/15/2008	1.62	±	1.44	6.00	±	5.34	
	10/22/2008	0.16	±	1.46	0.58	±	5.41	
	10/29/2008	-2.62	±	1.75	-9.70	±	6.48	
	11/05/2008	-2.93	±	1.75	-10.85	±	6.47	
	11/12/2008	-1.91	±	1.21	-7.05	±	4.48	
	11/19/2008	0.24	±	1.47	0.90	±	5.43	
	11/26/2008	0.50	±	1.45	1.86	±	5.35	
	12/03/2008	-0.09	±	1.40	-0.32	±	5.19	
	12/10/2008	0.97	±	2.02	3.58	±	7.47	
	12/17/2008	0.42	±	1.50	1.55	±	5.55	
	12/24/2008	-0.20	±	1.28	-0.73	±	4.73	
	12/31/2008	-0.46	±	1.36	-1.69	±	5.03	
JACKSON								
	10/01/2008	-0.99	±	1.82	-3.65	±	6.72	
a	10/08/2008	3.03	±	12.12	11.20	±	44.86	
a	10/15/2008		±			±		
	10/22/2008	-1.50	±	1.52	-5.54	±	5.63	
	10/29/2008	-0.03	±	1.64	-0.10	±	6.06	
	11/05/2008	0.47	±	1.60	1.74	±	5.91	
	11/12/2008	-1.29	±	1.42	-4.79	±	5.27	
	11/19/2008	0.25	±	1.58	0.94	±	5.84	
	11/26/2008	-0.03	±	2.32	-0.11	±	8.57	

TABLE C-2. Weekly Iodine-131 Activity in Air.

Sampling Group and Location	Sampling Date	Result ± 1s Uncertainty (x 10 ⁻¹⁵ µCi/mL)			Result ± 1s Uncertainty (x 10 ⁻¹¹ Bq/mL)			Result > 3s
BOUNDARY								
a	12/03/2008	-4.20	±	7.18	-15.55	±	26.55	
	12/10/2008	1.67	±	0.77	6.19	±	2.86	
	12/17/2008	-0.71	±	1.45	-2.64	±	5.37	
	12/24/2008	-0.43	±	1.53	-1.61	±	5.65	
	12/31/2008	-0.81	±	1.27	-3.02	±	4.70	
REXBURG CMS	10/01/2008	-1.29	±	1.33	-4.76	±	4.93	
	10/08/2008	1.78	±	1.27	6.57	±	4.70	
	10/15/2008	1.19	±	1.06	4.40	±	3.91	
	10/22/2008	0.14	±	1.27	0.50	±	4.70	
	10/29/2008	-2.26	±	1.51	-8.35	±	5.58	
	11/05/2008	-1.85	±	1.10	-6.83	±	4.07	
	11/12/2008	-1.60	±	1.02	-5.94	±	3.77	
	11/19/2008	0.22	±	1.30	0.80	±	4.81	
	11/26/2008	0.44	±	1.25	1.61	±	4.63	
	12/03/2008	-0.09	±	1.48	-0.34	±	5.48	
	12/10/2008	0.98	±	2.04	3.63	±	7.56	
	12/17/2008	0.33	±	1.17	1.21	±	4.33	
	12/24/2008	-0.17	±	1.13	-0.64	±	4.17	
	12/31/2008	-0.41	±	1.21	-1.51	±	4.47	
INL SITE								
EFS	10/01/2008	-1.00	±	1.84	-3.70	±	6.82	
	10/08/2008	0.40	±	1.60	1.48	±	5.91	
	10/15/2008	0.22	±	1.43	0.80	±	5.28	
	10/22/2008	-1.49	±	1.52	-5.53	±	5.62	
	10/29/2008	-0.03	±	1.63	-0.10	±	6.03	
	11/05/2008	0.43	±	1.47	1.59	±	5.42	
	11/12/2008	-1.23	±	1.35	-4.54	±	5.00	
	11/19/2008	0.24	±	1.51	0.89	±	5.57	
	11/26/2008	-2.29	±	1.50	-8.47	±	5.56	
	12/03/2008	-1.23	±	1.76	-4.55	±	6.51	
	12/10/2008	3.47	±	1.61	12.85	±	5.94	
	12/17/2008	-0.64	±	1.30	-2.36	±	4.81	
	12/24/2008	-0.37	±	1.30	-1.37	±	4.81	
	12/31/2008	-0.86	±	1.35	-3.19	±	4.98	
MAIN GATE	10/01/2008	-1.35	±	2.48	-4.98	±	9.18	
	10/08/2008	0.39	±	1.55	1.43	±	5.74	
	10/15/2008	0.22	±	1.45	0.81	±	5.36	
	10/22/2008	-1.48	±	1.51	-5.49	±	5.57	
	10/29/2008	-0.03	±	1.92	-0.11	±	7.09	
	11/05/2008	0.79	±	2.67	2.91	±	9.89	
	11/12/2008	-1.12	±	1.23	-4.14	±	4.55	
	11/19/2008	0.23	±	1.41	0.84	±	5.20	
	11/26/2008	-2.33	±	1.53	-8.60	±	5.65	
	12/03/2008	-1.08	±	1.55	-4.00	±	5.73	
	12/10/2008	5.27	±	2.44	19.51	±	9.02	
	12/17/2008	-1.41	±	2.87	-5.20	±	10.60	
	12/24/2008	-0.48	±	1.70	-1.79	±	6.27	
	12/31/2008	-0.42	±	1.26	-1.57	±	4.66	

TABLE C-2. Weekly Iodine-131 Activity in Air.

Sampling Group and Location	Sampling Date	Result ± 1s Uncertainty (x 10⁻¹⁵ µCi/mL)			Result ± 1s Uncertainty (x 10⁻¹¹ Bq/mL)			Result > 3s
BOUNDARY								
VAN BUREN GATE	10/01/2008	-1.23	±	2.26	-4.54	±	8.36	
	10/08/2008	0.40	±	1.60	1.47	±	5.90	
	10/15/2008	0.21	±	1.39	0.78	±	5.13	
	10/22/2008	-1.58	±	1.60	-5.84	±	5.93	
	10/29/2008	-0.03	±	1.68	-0.10	±	6.22	
	11/05/2008	0.42	±	1.42	1.54	±	5.25	
	11/12/2008	-1.20	±	1.31	-4.42	±	4.86	
	11/19/2008	0.24	±	1.49	0.89	±	5.52	
	11/26/2008	-2.16	±	1.41	-7.97	±	5.23	
	12/03/2008	-1.20	±	1.72	-4.44	±	6.35	
	12/10/2008	3.07	±	1.42	11.36	±	5.25	
	12/17/2008	-0.82	±	1.68	-3.05	±	6.21	
	12/24/2008	-0.48	±	1.70	-1.79	±	6.27	
	12/31/2008	-0.77	±	1.20	-2.84	±	4.43	
a. Invalid Sample Result								

TABLE C-3. Quarterly Americium-241, Cesium-137, Plutonium-238, Plutonium-239/240, and Strontium-90 Concentrations in Composite Air Filters.

Sampling Group and Location	Sampling Date	Analyte	Result ± 1s Uncertainty (x 10 ⁻¹⁸ µCi/mL)			Result ± 1s Uncertainty (x 10 ⁻¹³ Bq/mL)			Result > 3s
BOUNDARY									
ARCO	12/31/2008	CESIUM-137	130.00	±	137.00	481.00	±	506.90	
ATOMIC CITY	12/31/2008	AMERICIUM-241	3.30	±	0.62	12.22	±	2.28	Y
		CESIUM-137	9.00	±	87.60	33.30	±	324.12	
		PLUTONIUM-238	0.59	±	0.30	2.19	±	1.10	
		PLUTONIUM-239/40	0.30	±	0.42	1.10	±	1.55	
ATOMIC CITY (QA-1)	12/31/2008	AMERICIUM-241	796.90	±	50.90	2948.53	±	188.33	Y
		CESIUM-137	45.90	±	143.00	169.83	±	529.10	
		PLUTONIUM-238	63.16	±	4.01	233.69	±	14.84	Y
		PLUTONIUM-239/40	-0.42	±	0.72	-1.54	±	2.66	
BLUE DOME	12/31/2008	AMERICIUM-241	2.69	±	3.70	9.94	±	13.67	
		CESIUM-137	59.90	±	132.00	221.63	±	488.40	
		PLUTONIUM-238	0.00	±	0.00	0.00	±	0.00	
		PLUTONIUM-239/40	0.45	±	0.34	1.67	±	1.24	
BLUE DOME (QA-2)	12/31/2008	AMERICIUM-241	0.94	±	0.66	3.46	±	2.45	
		CESIUM-137	-302.00	±	152.00	-1117.40	±	562.40	
		PLUTONIUM-238	0.91	±	1.57	3.35	±	5.81	
		PLUTONIUM-239/40	128.20	±	9.12	474.34	±	33.73	Y
FAA TOWER	12/31/2008	CESIUM-137	204.00	±	167.00	754.80	±	617.90	
		STRONTIUM-90	43.70	±	14.60	161.69	±	54.02	
HOWE	12/31/2008	AMERICIUM-241	7.60	±	1.48	28.11	±	5.48	Y
		CESIUM-137	36.40	±	134.00	134.68	±	495.80	
		PLUTONIUM-238	-0.58	±	0.58	-2.14	±	2.14	
		PLUTONIUM-239/40	1.16	±	0.82	4.28	±	3.03	
MONTEVIEW		CESIUM-137	-34.30	±	131.00	-126.91	±	484.70	
MUD LAKE		CESIUM-137	-396.00	±	266.00	-1465.20	±	984.20	
		STRONTIUM-90	7.11	±	12.50	26.31	±	46.25	

TABLE C-3. Quarterly Americium-241, Cesium-137, Plutonium-238, Plutonium-239/240, and Strontium-90 Concentrations in Composite Air Filters.

Sampling Group and Location	Sampling Date	Analyte	Result ± 1s Uncertainty (x 10 ⁻¹⁸ µCi/mL)			Result ± 1s Uncertainty (x 10 ⁻¹³ Bq/mL)			Result > 3s
DISTANT									
BLACKFOOT	12/31/2008	CESIUM-137	87.70	±	128.00	324.49	±	473.60	
CRATERS	12/31/2008	CESIUM-137	-14.10	±	280.00	-52.17	±	1036.00	
DUBOIS	12/31/2008	CESIUM-137	64.90	±	81.60	240.13	±	301.92	
IDAHO FALLS	12/31/2008	CESIUM-137	192.00	±	146.00	710.40	±	540.20	
		STRONTIUM-90	29.40	±	15.85	108.78	±	58.65	
JACKSON	12/31/2008	CESIUM-137	-314.00	±	149.00	-1161.80	±	551.30	
REXBURG CMS	12/31/2008	AMERICIUM-241	0.83	±	0.72	3.08	±	2.67	
		CESIUM-137	-348.00	±	257.00	-1287.60	±	950.90	
		PLUTONIUM-238	3.87	±	2.00	14.31	±	7.39	
		PLUTONIUM-239/40	2.76	±	1.24	10.22	±	4.59	
INL SITE									
EFS	12/31/2008	AMERICIUM-241	16.23	±	1.76	60.05	±	6.50	Y
		CESIUM-137	45.00	±	131.00	166.50	±	484.70	
		PLUTONIUM-238	0.00	±	0.00	0.00	±	0.00	
		PLUTONIUM-239/40	1.05	±	1.05	3.87	±	3.88	
MAIN GATE	12/31/2008	CESIUM-137	30.80	±	188.00	113.96	±	695.60	
		STRONTIUM-90	-3.43	±	16.25	-12.69	±	60.13	
VAN BUREN GATE	12/31/2008	CESIUM-137	86.90	±	95.20	321.53	±	352.24	
		STRONTIUM-90	21.80	±	12.40	80.66	±	45.88	

TABLE C-4. Tritium Concentrations in Atmospheric Moisture.

Sampling Group and Location	Start Date	Sampling Date	Result ± 1s Uncertainty			Result ± 1s Uncertainty			Collection Medium	Result > 3s
			(x 10 ⁻¹³ μCi/mL _{air})			(x 10 ⁻⁹ Bq/mL _{air})				
BOUNDARY										
ATOMIC CITY	09/03/2008	10/01/2008	4.14	±	1.27	15.33	±	4.71	Molecular Sieve	Y
ATOMIC CITY	10/01/2008	10/22/2008	5.69	±	1.90	21.07	±	7.04	Molecular Sieve	
ATOMIC CITY	10/22/2008	11/19/2008	5.11	±	1.88	18.89	±	6.97	Molecular Sieve	
ATOMIC CITY	11/19/2008	12/31/2008	1.24	±	1.37	4.57	±	5.07	Molecular Sieve	
DISTANT										
BLACKFOOT	09/19/2008	10/03/2008	7.15	±	2.06	26.47	±	7.62	Molecular Sieve	Y
BLACKFOOT	10/03/2008	10/22/2008	5.83	±	1.70	21.59	±	6.30	Molecular Sieve	Y
BLACKFOOT	10/22/2008	11/12/2008	13.83	±	1.88	51.19	±	6.96	Molecular Sieve	Y
BLACKFOOT	11/12/2008	12/10/2008	4.62	±	1.43	17.11	±	5.29	Molecular Sieve	Y
IDAHO FALLS	09/10/2008	10/10/2008	2.83	±	1.30	10.47	±	4.82	Molecular Sieve	
IDAHO FALLS	10/10/2008	11/03/2008	5.51	±	1.57	20.39	±	5.82	Molecular Sieve	Y
IDAHO FALLS	11/03/2008	12/04/2008	4.13	±	1.53	15.28	±	5.65	Molecular Sieve	
REXBURG CMS	09/11/2008	10/07/2008	4.72	±	1.77	17.47	±	6.54	Molecular Sieve	
REXBURG CMS	10/07/2008	11/12/2008	3.08	±	1.73	11.39	±	6.42	Molecular Sieve	
REXBURG CMS	11/12/2008	12/31/2008	5.09	±	1.38	18.84	±	5.10	Molecular Sieve	Y

TABLE C-5. Monthly and Weekly Tritium Concentrations in Precipitation.

Location	Start Date	End Date	Result \pm 1s Uncertainty			Result \pm 1s Uncertainty			Result > 3s
			(pCi/L)			(Bq/L)			
IDAHO FALLS	10/1/2008	11/10/2008	33.10	\pm	31.70	1.22	\pm	1.17	
	11/10/2008	12/1/2008	4.15	\pm	32.70	0.15	\pm	1.21	
CFA	9/2/2008	10/1/2008	77.00	\pm	32.60	2.85	\pm	1.21	
	10/1/2008	11/3/2008	100.00	\pm	32.40	3.70	\pm	1.20	Y
	11/3/2008	12/10/2008	36.00	\pm	31.70	1.33	\pm	1.17	
EFS	10/1/2008	10/8/2008	60.00	\pm	33.40	2.22	\pm	1.24	
	10/8/2008	10/15/2008	139.00	\pm	33.80	5.14	\pm	1.25	Y
	10/29/2008	11/5/2008	156.00	\pm	34.20	5.77	\pm	1.27	Y
	11/5/2008	11/12/2008	67.20	\pm	32.80	2.49	\pm	1.21	
	12/3/2008	12/10/2008	44.70	\pm	31.90	1.65	\pm	1.18	
	12/10/2008	12/17/2008	44.10	\pm	32.30	1.63	\pm	1.20	

Table C-6. Weekly and Monthly Iodine-131 and Cesium-137 Concentrations in Milk

Location	Sampling Date	Iodine-131			Cesium-137		
		Result ± 1s Uncertainty (pCi [†] /L)		Result ± 1s Uncertainty (Bq [†] /L)	Result > 3s	Result ± 1s Uncertainty (pCi/L)	
BLACKFOOT	10/9/2008	-0.95 ± 1.17	-0.035 ± 0.043		0.09 ± 0.83	0.003 ± 0.031	
DIETRICH	10/07/2008	0.19 ± 1.05	0.007 ± 0.039		-1.30 ± 0.92	-0.048 ± 0.034	
	11/04/2008	-0.38 ± 1.06	-0.014 ± 0.039		0.28 ± 0.89	0.010 ± 0.033	
	12/02/2008	-0.90 ± 1.59	-0.033 ± 0.059		-1.84 ± 1.39	-0.068 ± 0.051	
Duplicate	12/02/2008	-0.42 ± 2.55	-0.015 ± 0.094		-0.67 ± 2.66	-0.025 ± 0.099	
FORT HALL	10/13/2008	-0.55 ± 0.87	-0.020 ± 0.032		-0.28 ± 0.77	-0.010 ± 0.028	
	11/10/2008	-0.84 ± 1.24	-0.031 ± 0.046		0.23 ± 0.86	0.008 ± 0.032	
	12/08/2008	-0.59 ± 1.22	-0.022 ± 0.045		-1.12 ± 0.96	-0.041 ± 0.035	
HOWE	10/07/2008	-0.95 ± 2.40	-0.035 ± 0.089		2.46 ± 2.70	0.091 ± 0.100	
	11/04/2008	-2.13 ± 1.18	-0.079 ± 0.044		0.29 ± 0.90	0.011 ± 0.033	
	12/02/2008	-1.85 ± 0.82	-0.069 ± 0.030		-0.71 ± 0.81	-0.026 ± 0.030	
IDAHO FALLS	10/07/2008	-0.67 ± 0.66	-0.025 ± 0.025		1.38 ± 0.77	0.051 ± 0.028	
	10/14/2008	-1.63 ± 1.49	-0.060 ± 0.055		1.50 ± 1.29	0.056 ± 0.048	
	10/21/2008	0.21 ± 1.50	0.008 ± 0.056		1.39 ± 1.28	0.051 ± 0.047	
	10/28/2008	0.31 ± 0.67	0.012 ± 0.025		-1.25 ± 0.80	-0.046 ± 0.030	
	11/04/2008	0.65 ± 0.73	0.024 ± 0.027		-0.88 ± 0.76	-0.033 ± 0.028	
	11/11/2008	0.88 ± 2.76	0.033 ± 0.102		-1.67 ± 1.47	-0.062 ± 0.054	
	11/18/2008	0.29 ± 0.76	0.011 ± 0.028		0.75 ± 0.82	0.028 ± 0.030	
	11/25/2008	0.24 ± 0.74	0.009 ± 0.027		0.29 ± 0.80	0.011 ± 0.029	
	12/02/2008	-1.48 ± 0.73	-0.055 ± 0.027		-0.22 ± 0.80	-0.008 ± 0.030	
	12/09/2008	-0.88 ± 2.40	-0.033 ± 0.089		0.25 ± 2.69	0.009 ± 0.100	
	12/16/2008	0.37 ± 0.78	0.014 ± 0.029		0.09 ± 0.82	0.003 ± 0.030	
	12/23/2008	-0.75 ± 0.75	-0.028 ± 0.028		-0.31 ± 0.81	-0.011 ± 0.030	
	12/30/2008	1.33 ± 0.74	0.049 ± 0.027		0.21 ± 0.81	0.008 ± 0.030	
MORELAND	10/07/2008	-1.34 ± 2.59	-0.050 ± 0.096		-1.97 ± 2.68	-0.073 ± 0.099	
	11/04/2008	0.29 ± 0.83	0.011 ± 0.031		-1.09 ± 0.80	-0.040 ± 0.030	

Table C-6. Weekly and Monthly Iodine-131 and Cesium-137 Concentrations in Milk

	12/02/2008	0.04	±	2.18	0.002	±	0.081	1.71	±	2.63	0.063	±	0.097
RUPERT													
	10/07/2008	0.37	±	2.13	0.014	±	0.079	-1.80	±	2.63	-0.067	±	0.097
	11/04/2008	-1.13	±	2.15	-0.042	±	0.080	0.84	±	2.68	0.031	±	0.099
	12/02/2008	-0.09	±	1.11	-0.003	±	0.041	0.35	±	0.91	0.013	±	0.034
TERRETON													
	10/07/2008	1.11	±	1.18	0.041	±	0.044	-0.66	±	0.93	-0.024	±	0.034
	11/04/2008	3.59	±	2.33	0.133	±	0.086	-0.12	±	2.68	-0.005	±	0.099
	12/02/2008	0.33	±	0.89	0.012	±	0.033	-0.83	±	0.81	-0.031	±	0.030
Duplicate	12/02/2008	0.13	±	1.32	0.005	±	0.049	-1.61	±	0.88	-0.060	±	0.033

Table C-7. Strontium-90 and Tritium Concentrations in Milk

Strontium-90								
Location	Sampling Date	Result \pm 1s Uncertainty (pCi/L)			Result \pm 1s Uncertainty (Bq/L)			Result > 3s
FORT HALL	11/04/2008	0.24	\pm	0.05	0.009	\pm	0.002	Y
HOWE	11/04/2008	0.16	\pm	0.05	0.006	\pm	0.002	
IDAHO FALLS	11/04/2008	0.67	\pm	0.07	0.025	\pm	0.003	Y
MORELAND	11/04/2008	0.64	\pm	0.07	0.024	\pm	0.003	Y
Tritium								
		Concentration \pm 1s			Concentration \pm 1s			Result > 3s
DIETRICH	11/04/2008	43.20	\pm	33.80	1.600	\pm	1.252	
FORT HALL	11/10/2008	102.00	\pm	33.70	3.778	\pm	1.248	Y
RUPERT	11/04/2008	62.90	\pm	34.20	2.330	\pm	1.267	
TERRETON	11/04/2008	52.69	\pm	33.50	1.952	\pm	1.241	

Table C-8. Cesium-137 and Strontium-90 Concentrations in Potatoes

Sampling Date	Location	Cesium-137						
		Result ± 1s Uncertainty pCi/kg			Result ± 1s Uncertainty Bq/kg			Result > 3s
10/1/2008	ARCO	1.26	±	0.71	0.05	±	0.03	
10/1/2008	ARCO (DUPLICATE)	-0.53	±	0.74	-0.02	±	0.03	
10/9/2008	BLACKFOOT	-0.38	±	0.47	-0.01	±	0.02	
11/14/2008	COLORADO	-0.44	±	0.73	-0.02	±	0.03	
9/29/2008	IDAHO FALLS	-0.98	±	0.48	-0.04	±	0.02	
10/7/2008	MINIDOKA	0.04	±	1.77	0.00	±	0.07	
10/8/2008	MONTEVIEW	2.51	±	1.76	0.09	±	0.07	
11/11/2008	OREGON	-1.65	±	1.75	-0.06	±	0.06	
		Strontium-90						
Sampling Date	Location	Result ± 1s Uncertainty pCi/kg			Result ± 1s Uncertainty Bq/kg			Result > 3s
10/1/2008	ARCO	3.73	±	1.68	0.14	±	0.06	
10/1/2008	ARCO (DUPLICATE)	5.49	±	1.90	0.20	±	0.07	
10/9/2008	BLACKFOOT	-2.90	±	1.34	-0.11	±	0.05	
11/14/2008	COLORADO	2.01	±	1.08	0.07	±	0.04	
9/29/2008	IDAHO FALLS	4.53	±	1.20	0.17	±	0.04	Y
10/7/2008	MINIDOKA	4.89	±	1.43	0.18	±	0.05	Y
10/8/2008	MONTEVIEW	1.15	±	1.03	0.04	±	0.04	
11/11/2008	OREGON	-1.55	±	1.41	-0.06	±	0.05	

Table C-9. Cesium-137 and Cobalt-60 Concentrations in Soil

Location	Sampling Date	Cesium-137				Result > 3s
		Concentration ± 1s (pCi/Kg)		Concentration ± 1s (Bq/Kg)		
BOUNDARY						
ATOMIC CITY	08/25/08	524.94	± 15.81	19.44	± 0.59	Y
BUTTE CITY	08/25/08	69.76	± 5.86	2.58	± 0.22	Y
FAA TOWER	08/25/08	457.65	± 11.79	16.95	± 0.44	Y
HOWE	08/18/08	307.67	± 7.20	11.40	± 0.27	Y
MONTEVIEW	08/18/08	360.02	± 8.96	13.33	± 0.33	Y
MUD LAKE #1	08/18/08	341.19	± 7.76	12.64	± 0.29	Y
MUD LAKE #1 (DUP)	08/18/08	401.03	± 9.02	14.85	± 0.33	Y
MUD LAKE #2	08/18/08	316.78	± 10.69	11.73	± 0.40	Y
RENO RANCH	08/18/08	664.29	± 19.46	24.60	± 0.72	Y
DISTANT						
BLACKFOOT	08/25/08	602.05	± 14.10	22.30	± 0.52	Y
CAREY	08/25/08	445.92	± 11.67	16.52	± 0.43	Y
CRYSTAL ICE CAVES	08/26/08	500.00	± 11.17	18.52	± 0.41	Y
ST. ANTHONY	08/18/08	487.50	± 10.87	18.06	± 0.40	Y
Cobalt-60						
Location	Sampling Date	Cobalt-60				Result > 3s
		Concentration ± 1s (pCi/Kg)		Concentration ± 1s (Bq/Kg)		
BOUNDARY						
ATOMIC CITY	08/25/08	-4.67	± 584.51	-0.17	± 21.65	
BUTTE CITY	08/25/08	-2.02	± 419.96	-0.07	± 15.55	
FAA TOWER	08/25/08	1.41	± 2.69	0.05	± 0.10	
HOWE	08/18/08	0.90	± 1.60	0.03	± 0.06	
MONTEVIEW	08/18/08	-4.52	± 176.21	-0.17	± 6.53	
MUD LAKE #1	08/18/08	-0.47	± 139.21	-0.02	± 5.16	
MUD LAKE #1 (DUP)	08/18/08	1.49	± 1.48	0.06	± 0.05	
MUD LAKE #2	08/18/08	-6.81	± 420.17	-0.25	± 15.56	
RENO RANCH	08/18/08	-8.54	± 504.40	-0.32	± 18.68	
DISTANT						
BLACKFOOT	08/25/08	0.40	± 2.87	0.01	± 0.11	
CAREY	08/25/08	-1.70	± 286.48	-0.06	± 10.61	
CRYSTAL ICE CAVES	08/26/08	-3.09	± 165.51	-0.11	± 6.13	
ST. ANTHONY	08/18/08	3.37	± 1.63	0.12	± 0.06	

TABLE C-10. Americium-241, Plutonium, and Strontium-90 Concentrations in Soil

Location	Sampling Date	Americium-241						
		Concentration ± 1s (pCi/Kg)			Concentration ± 1s (Bq/Kg)			Result > 3s
BOUNDARY								
ATOMIC CITY	08/25/08	4.65	±	16.94	0.17	±	0.63	
BUTTE CITY	08/25/08	94.15	±	44.33	3.49	±	1.64	
FAA TOWER	08/25/08	13.78	±	12.91	0.51	±	0.48	
HOWE	08/18/08	4.20	±	6.75	0.16	±	0.25	
MONTEVIEW	08/18/08	8.51	±	11.20	0.32	±	0.41	
MUD LAKE #1	08/18/08	87.49	±	42.55	3.24	±	1.58	
MUD LAKE #1 (DUP)	08/18/08	-23.12	±	39.81	-0.86	±	1.47	
MUD LAKE #2	08/18/08	9.00	±	23.20	0.33	±	0.86	
RENO RANCH	08/18/08	26.82	±	16.76	0.99	±	0.62	
DISTANT								
BLACKFOOT SOIL	08/25/08	-6.24	±	4.42	-0.23	±	0.16	
CAREY SOIL	08/25/08	28.59	±	16.75	1.06	±	0.62	
CRYSTAL ICE CAVES	08/26/08	-4.09	±	16.01	-0.15	±	0.59	
ST. ANTHONY SOIL	08/18/08	13.20	±	10.84	0.49	±	0.40	

Location	Sampling Date	Plutonium-238						
		Concentration ± 1s (pCi/Kg)			Concentration ± 1s (Bq/Kg)			Result > 3s
BOUNDARY								
ATOMIC CITY	08/25/08	1.64	±	6.00	0.06	±	0.22	
BUTTE CITY	08/25/08	16.23	±	13.17	0.60	±	0.49	
FAA TOWER	08/25/08	20.17	±	14.65	0.75	±	0.54	
HOWE	08/18/08	-5.16	±	8.89	-0.19	±	0.33	
MONTEVIEW	08/18/08	22.70	±	21.04	0.84	±	0.78	
MUD LAKE #1	08/18/08	20.54	±	16.68	0.76	±	0.62	
MUD LAKE #1 (DUP)	08/18/08	-0.36	±	10.20	-0.01	±	0.38	
MUD LAKE #2	08/18/08	-26.14	±	8.80	-0.97	±	0.33	
RENO RANCH	08/18/08	4.46	±	11.51	0.17	±	0.43	
DISTANT								
BLACKFOOT SOIL	08/25/08	-5.63	±	8.80	-0.21	±	0.33	
CAREY SOIL	08/25/08	33.64	±	16.84	1.25	±	0.62	
CRYSTAL ICE CAVES	08/26/08	7.20	±	11.55	0.27	±	0.43	
ST. ANTHONY SOIL	08/18/08	18.99	±	15.57	0.70	±	0.58	

TABLE C-10. Americium-241, Plutonium, and Strontium-90 Concentrations in Soil

Location	Sampling Date	Plutonium-239/240						Result > 3s
		Concentration ± 1s (pCi/Kg)			Concentration ± 1s (Bq/Kg)			
BOUNDARY								
ATOMIC CITY	08/25/08	40.95	±	16.64	1.52	±	0.62	
BUTTE CITY	08/25/08	8.11	±	9.31	0.30	±	0.34	
FAA TOWER	08/25/08	14.25	±	11.14	0.53	±	0.41	
HOWE	08/18/08	-4.87	±	3.45	-0.18	±	0.13	
MONTEVIEW	08/18/08	2.41	±	8.80	0.09	±	0.33	
MUD LAKE #1	08/18/08	41.73	±	19.74	1.55	±	0.73	
MUD LAKE #1 (DUP)	08/18/08	37.26	±	22.52	1.38	±	0.83	
MUD LAKE #2	08/18/08	10.92	±	12.53	0.40	±	0.46	
RENO RANCH	08/18/08	16.54	±	15.69	0.61	±	0.58	
DISTANT								
BLACKFOOT	08/25/08	18.29	±	10.19	0.68	±	0.38	
CAREY	08/25/08	-3.95	±	2.80	-0.15	±	0.10	
CRYSTAL ICE CAVES	08/26/08	24.50	±	17.81	0.91	±	0.66	
ST. ANTHONY	08/18/08	11.79	±	8.35	0.44	±	0.31	
Location	Sampling Date	Strontium-90						Result > 3s
		Concentration ± 1s (pCi/Kg)			Concentration ± 1s (Bq/Kg)			
BOUNDARY								
ATOMIC CITY	08/25/08	241.00	±	23.45	8.93	±	0.87	Y
BUTTE CITY	08/25/08	91.00	±	12.55	3.37	±	0.46	Y
FAA TOWER	08/25/08	199.00	±	20.95	7.37	±	0.78	Y
HOWE	08/18/08	123.00	±	13.90	4.56	±	0.51	Y
MONTEVIEW	08/18/08	56.60	±	10.00	2.10	±	0.37	Y
MUD LAKE #1	08/18/08	24.30	±	11.25	0.90	±	0.42	
MUD LAKE #1 (DUP)	08/18/08	24.50	±	12.75	0.91	±	0.47	
MUD LAKE #2	08/18/08	81.30	±	14.65	3.01	±	0.54	Y
RENO RANCH	08/18/08	233.00	±	14.30	8.63	±	0.53	Y
DISTANT								
BLACKFOOT	08/25/08	0.72	±	6.35	0.03	±	0.24	
CAREY	08/25/08	44.70	±	19.20	1.66	±	0.71	
CRYSTAL ICE CAVES	08/26/08	197.00	±	14.75	7.30	±	0.55	Y
ST. ANTHONY	08/18/08	174.00	±	11.40	6.44	±	0.42	Y

Table C-11. Cesium-137 and Iodine-131 Concentrations in Large Game Animals

Species	Collection		Analyte	Result ± 1s Uncertainty			Result ± 1s Uncertainty			Result > 3s
	Date	Tissue		(pCi/kg wet weight)			(x 10 ⁻² Bq/kg wet weight)			
MULE DEER	10/14/2008	Muscle	¹³¹ I	-2.36	±	27.60	-8.73	±	102.12	
			¹³⁷ Cs	3.39	±	3.77	12.54	±	13.95	
ELK	11/17/2008	Muscle	¹³¹ I	17.60	±	25.20	65.12	±	93.24	
			¹³⁷ Cs	1.04	±	4.84	3.85	±	17.91	
MULE DEER	10/14/2008	Liver	¹³¹ I	-26.50	±	36.70	-98.05	±	135.79	
			¹³⁷ Cs	0.48	±	2.86	1.76	±	10.58	
MULE DEER	10/14/2008	Thyroid	¹³¹ I	-67.80	±	68.10	-250.86	±	251.97	
			¹³⁷ Cs	196.00	±	61.80	725.20	±	228.66	Y
ELK	11/17/2008	Thyroid	¹³¹ I	23.30	±	32.30	86.21	±	119.51	
			¹³⁷ Cs	27.80	±	30.30	102.86	±	112.11	

**Table C-12. Gamma-emitting Radionuclides.
Strontium-90 and Actinides in Waterfowl**

Location	Sampling	Result ± Uncertainty(1s)			Result ± Uncertainty(1s)			Result > 3s
Species	Date	Analyte	(x 10 ⁻³) pCi/g		(x 10 ⁻⁵) Bq/g			
RTC	9/12/2008							
Eared Grebe		AMERICIUM-241	104.80	± 10.77	388.15	± 39.87	Y	
		CESIUM-137	158.78	± 7.23	588.09	± 26.78	Y	
		CHROMIUM-51	-149.80	± 177.37	-554.80	± 656.93		
		COBALT-60	12.20	± 3.23	45.18	± 11.98	Y	
		PLUTONIUM-238	-9.20	± 9.24	-34.07	± 34.22		
		PLUTONIUM-239/240	46.00	± 24.73	170.37	± 91.59		
		STRONTIUM-90	5.05	± 1.17	18.70	± 4.33	Y	
		ZINC-65	14.38	± 10.82	53.26	± 40.09		
RTC	9/12/2008							
Eared Grebe		CESIUM-137	116.96	± 17.08	433.17	± 63.25	Y	
		CHROMIUM-51	-371.34	± 456.43	-1375.35	± 1690.48		
		COBALT-60	-15.57	± 1453.92	-57.65	± 5384.89		
		PLUTONIUM-238	31.63	± 14.45	117.15	± 53.50		
		PLUTONIUM-239/240	12.65	± 12.71	46.85	± 47.06		
		STRONTIUM-90	7.23	± 1.05	26.78	± 3.89	Y	
		ZINC-65	16.81	± 27.27	62.26	± 101.01		
RTC	9/12/2008							
Eared Grebe		AMERICIUM-241	108.50	± 15.50	401.85	± 57.41	Y	
		CESIUM-137	181.50	± 22.97	672.22	± 85.08	Y	
		CHROMIUM-51	968.97	± 987.97	3588.78	± 3659.14		
		COBALT-60	-27.06	± 1975.11	-100.22	± 7315.21		
		PLUTONIUM-238	0.00	± 0.00	0.00	± 0.00		
		PLUTONIUM-239/240	330.00	± 137.65	1222.22	± 509.81		
		STRONTIUM-90	8.60	± 1.27	31.85	± 4.69	Y	
		ZINC-65	-11.32	± 39.87	-41.93	± 147.65		
RTC	9/12/2008							
Eared Grebe		AMERICIUM-241	267.00	± 23.90	988.89	± 88.52	Y	
		CESIUM-137	520.23	± 13.36	1926.78	± 49.48	Y	
		CHROMIUM-51	470.76	± 250.26	1743.55	± 926.87		

**Table C-12. Gamma-emitting Radionuclides.
Strontium-90 and Actinides in Waterfowl**

	COBALT-60	80.75	±	4.02	299.09	±	14.88	Y
	PLUTONIUM-238	0.00	±	0.00	0.00	±	0.00	
	PLUTONIUM-239/240	8.04	±	13.95	29.77	±	51.65	
	STRONTIUM-90	3.08	±	0.78	11.41	±	2.87	Y
	ZINC-65	275.46	±	14.93	1020.24	±	55.30	Y
MFC	9/20/2008							
Shoveler	AMERICIUM-241	19.56	±	8.54	72.44	±	31.63	
	CESIUM-137	46.17	±	19.37	171.00	±	71.75	
	CHROMIUM-51	-231.07	±	457.60	-855.83	±	1694.82	
	COBALT-60	-45.56	±	1792.56	-168.75	±	6639.10	
	PLUTONIUM-238	3.44	±	5.97	12.74	±	22.09	
	PLUTONIUM-239/240	17.20	±	9.21	63.70	±	34.11	
	STRONTIUM-90	2.24	±	0.88	8.30	±	3.24	
	ZINC-65	-56.00	±	33.47	-207.39	±	123.94	
MFC	9/20/2008							
Gadwall	AMERICIUM-241	12.64	±	7.00	46.81	±	25.91	
	CESIUM-137	24.55	±	15.17	90.92	±	56.20	
	CHROMIUM-51	-16.33	±	517.14	-60.47	±	1915.35	
	COBALT-60	-40.13	±	1452.51	-148.62	±	5379.67	
	PLUTONIUM-238	9.90	±	5.78	36.66	±	21.39	
	PLUTONIUM-239/240	13.20	±	6.69	48.89	±	24.78	
	STRONTIUM-90	0.74	±	0.59	2.73	±	2.19	
	ZINC-65	-14.83	±	27.28	-54.93	±	101.03	
MFC	9/26/2008							
Wigeon	AMERICIUM-241	17.99	±	7.96	66.63	±	29.48	
	CESIUM-137	-2.34	±	18.19	-8.65	±	67.38	
	CHROMIUM-51	-738.22	±	551.07	-2734.16	±	2040.99	
	COBALT-60	-42.42	±	1765.52	-157.11	±	6538.95	
	PLUTONIUM-238	0.00	±	0.00	0.00	±	0.00	
	PLUTONIUM-239/240	5.80	±	4.13	21.49	±	15.30	
	STRONTIUM-90	-1.13	±	0.52	-4.19	±	1.91	
	ZINC-65	-11.94	±	30.26	-44.23	±	112.06	
CONTROL	10/5/2008							

**Table C-12. Gamma-emitting Radionuclides.
Strontium-90 and Actinides in Waterfowl**

Green-winged Teal	AMERICIUM-241	15.25	±	5.40	56.48	±	20.00		
	CESIUM-137	13.34	±	14.52	49.40	±	53.78		
	CHROMIUM-51	-218.45	±	314.89	-809.09	±	1166.26		
	COBALT-60	5.01	±	10.22	18.57	±	37.84		
	STRONTIUM-90	0.19	±	1.73	0.69	±	6.39		
	ZINC-65	-28.92	±	30.27	-107.10	±	112.10		
<hr/>									
CONTROL	10/5/2008								
Green-winged Teal	AMERICIUM-241	-24.01	±	93.05	-88.93	±	344.63		
	CESIUM-137	3.95	±	8.58	14.65	±	31.79		
	CHROMIUM-51	-220.34	±	209.97	-816.08	±	777.68		
	COBALT-60	0.40	±	6.07	1.48	±	22.48		
	PLUTONIUM-238	121.60	±	21.17	450.37	±	78.41	Y	
	PLUTONIUM-239/240	55.51	±	13.47	205.59	±	49.87	Y	
	STRONTIUM-90	1.81	±	0.67	6.70	±	2.46		
ZINC-65	6.22	±	20.33	23.02	±	75.31			
<hr/>									

Table C-13. Environmental Radiation Results

Location	Start Date	End Date	Radiation Measurement \pm 2s Uncertainty mR	Exposure mR/day
BOUNDARY				
ARCO	5/7/2008	11/5/2008	63.5 \pm 12.4	0.35
ATOMIC CITY	5/7/2008	11/5/2008	64.3 \pm 12.6	0.35
BIRCH CREEK	5/7/2008	11/5/2008	55.1 \pm 10.8	0.30
BLUE DOME	5/7/2008	11/5/2008	54.5 \pm 10.7	0.30
HOWE	5/7/2008	11/5/2008	60.8 \pm 11.9	0.33
MONTEVIEW	5/7/2008	11/5/2008	59.3 \pm 11.6	0.33
MUD LAKE	5/7/2008	11/5/2008	65.5 \pm 12.8	0.36
Boundary Average				0.33
DISTANT				
ABERDEEN	5/6/2008	11/4/2008	64.4 \pm 12.6	0.35
BLACKFOOT	5/7/2008	11/5/2008	58.8 \pm 11.5	0.32
BLACKFOOT CMS	5/7/2008	11/5/2008	54.6 \pm 10.7	0.30
CRATERS	5/7/2008	11/5/2008	60.6 \pm 11.9	0.33
DUBOIS	5/7/2008	11/5/2008	50.8 \pm 10.0	0.28
IDAHO FALLS	5/5/2008	11/6/2008	62.3 \pm 12.2	0.34
MINIDOKA	5/6/2008	11/4/2008	55.6 \pm 10.9	0.31
REXBURG	5/7/2008	11/5/2008	73.0 \pm 14.3	0.40
ROBERTS	5/6/2008	11/4/2008	67.3 \pm 13.2	0.37
Distant Average				0.33
OUT-OF-STATE				
JACKSON	5/5/2008	11/6/2008	50.7 \pm 9.9	0.27

APPENDIX D
STATISTICAL ANALYSIS RESULTS

Table D-1. Results of the Kruskal-Wallace statistical test between INL Site, Boundary and Distant sample groups by month.

Parameter	P^a
Gross Alpha	
Quarter	0.79
October	0.69
November	0.68
December	0.76
Gross Beta	
Quarter	0.03
October	0.52
November	0.08
December	0.08
a. A 'p' value greater than 0.05 signifies no statistical difference between data groups.	

Table D-2. Statistical difference in weekly gross alpha and gross beta concentrations measured at Boundary and Distant locations.

Parameter	Mann-Whitney U test	
	Week	P ^a
Gross Alpha		
	October 1	0.39
	October 8	0.94
	October 15	0.26
	October 22	0.87
	October 29	0.78
	November 5	0.02
	November 12	0.72
	November 19	0.43
	November 26	0.78
	December 3	0.57
	December 10	0.20
	December 17	0.67
	December 24	0.68
	December 31	0.52
Gross Beta		
	October 1	0.67
	October 8	0.94
	October 15	0.94
	October 22	0.52
	October 29	0.25
	November 5	0.48
	November 12	0.00
	November 19	0.05
	November 26	0.35
	December 3	0.37
	December 10	0.05
	December 17	0.28
	December 24	0.42
	December 31	0.05

a. A 'p' value greater than 0.05 signifies no statistical difference between data groups.