

S.M. Stoller Corporation  
Environmental Surveillance, Education, and Research Program  
ISSN NUMBER 1089-5469

# Idaho National Laboratory Site Offsite Environmental Surveillance Program Report: Second Quarter 2010

October 2010



*Contributors:  
Marilyn Case, Russ Mitchell*

Program conducted for the U.S. Department of Energy, Idaho Operations Office  
Under Contract DE-AC07-06ID14680

By the S.M. Stoller Corporation,  
Environmental Surveillance, Education and Research Program  
Douglas K. Halford, Program Manager  
120 Technology Dr., Idaho Falls, Idaho 83402  
[www.stoller-eser.com](http://www.stoller-eser.com)

## **EXECUTIVE SUMMARY**

None of the radionuclides detected in samples collected during the second quarter of 2010 could be directly linked with INL Site activities. Levels of detected radionuclides were no different than values measured at other locations across the United States or were consistent with levels measured historically at the INL Site. All detected radionuclide concentrations were well below guidelines set by the U.S. Department of Energy (DOE) and regulatory standards established by the U.S. Environmental Protection Agency (EPA) for protection of the public.

This report for the second quarter of 2010 contains results from the Environmental Surveillance, Education and Research (ESER) Program's monitoring of the Department of Energy's Idaho National Laboratory (INL) Site's offsite environment, April 1 through June 30, 2010. All sample types (media) and the sampling schedule followed during 2010 are listed in Appendix A. Specifically, this report contains the results for the following:

- Air sampling, including particulate air filters, charcoal cartridges, and atmospheric moisture
- Precipitation sampling
- Surface water sampling
- Milk sampling
- Game animal sampling
- Environmental radiation measurements

Table E-1 Summary of results for the Second Quarter of 2010.

Media	Sample Type	Analysis	Results
Air	Filters	Gross alpha, gross beta	No statistical differences were noted in either gross alpha or gross beta data on quarterly and monthly comparisons. No statistical differences were observed between Distant locations than Boundary locations for either gross alpha or gross beta concentrations measured during any week of the second quarter. No result exceeded the DCG for gross alpha or gross beta activity in air.
		Gamma-emitting radionuclides, select actinides ( $^{241}\text{Am}$ , $^{238}\text{Pu}$ , and $^{239,240}\text{Pu}$ ), $^{90}\text{Sr}$	No man-made radionuclides were detected.
	Charcoal Cartridge	Iodine-131	Iodine-131 was initially reported at just above the detection level on one charcoal cartridge batch during the week of May 19. Two subsequent recounts failed to confirm the presence of $^{131}\text{I}$ .
Atmospheric Moisture	Liquid	Tritium	A total of 17 samples were collected. Nine of these samples had tritium results greater than the 3s uncertainty. No sample result exceeded the DCG for tritium in air. Results are consistent with historical measurements.
Precipitation	Liquid	Tritium	Thirteen samples were collected. Five of the results were greater than the 3s uncertainty. The concentration was consistent with those reported across the region by the Environmental Protection Agency and with previous results.
Surface Water (Big Lost River) <sup>a</sup>	Liquid	Gross alpha, gross beta, tritium	Five samples were collected from the BLR at the offsite Rest Area, near INTEC, at EFS (duplicates), and just north of NRF. All samples had detectable concentrations of gross alpha activity. Gross beta activity was detected in all samples except in the one from near NRF. Tritium was detected in all samples but the one collected near INTEC.
Milk	Liquid	Iodine-131, gamma-emitting radionuclides	Thirty-six samples, including two duplicates, were collected. No Iodine-131 or other manmade gamma-emitting radionuclides were detected in any sample. Strontium-90 was detected all eight samples selected for this analysis.
Large Game Animals	Tissue	Gamma-emitting radionuclides	Five large game animals (three pronghorn and two mule deer) were sampled. No man-made radionuclides were detected in animal tissues.
Thermo-	Environmental	External radioactivity	The average measurements over the six-

luminescent  
Dosimeters

radiation

month period were 0.34 mrem/day at boundary and 0.35 mrem/day at distant locations.

- 
- a. The Big Lost River only flows occasionally, usually after a large spring runoff event. Samples are thus collected opportunistically, not routinely. This quarter samples were collected at the Rest Area before entering the INL Site, near the INTEC, near the EFS, and just north of NRF. There was no water in the Big Lost River Sinks.
- 
-

### LIST OF ABBREVIATIONS

AEC	Atomic Energy Commission
CFA	Central Facilities Area
DCG	Derived Concentration Guide
DOE	Department of Energy
DOE – ID	Department of Energy Idaho Operations Office
EAL	Environmental Assessment Laboratory
EFS	Experimental Field Station
EPA	Environmental Protection Agency
ERAMS	Environmental Radiation Ambient Monitoring System
ESER	Environmental Surveillance, Education, and Research
ICP	Idaho Cleanup Project
INL	Idaho National Laboratory
INEL	Idaho National Engineering Laboratory
INEEL	Idaho National Engineering and Environmental Laboratory
ISU	Idaho State University
MDC	minimum detectable concentration
NRTS	National Reactor Testing Station

## **LIST OF UNITS**

Bq	becquerel
Ci	curie
g	gram
L	liter
$\mu$ Ci	microcurie
mL	milliliter
pCi	picocurie

## 1. ESEER PROGRAM DESCRIPTION

Operations at the Idaho National Laboratory (INL) Site are conducted under requirements imposed by the U.S. Department of Energy (DOE) under authority of the Atomic Energy Act and the U.S. Environmental Protection Agency (EPA) under a number of acts (e.g. the Clean Air Act and Safe Drinking Water Act). The requirements imposed by DOE are specified in DOE Orders. These requirements include those to monitor the effects of DOE activities both inside and outside the boundaries of DOE facilities (DOE 2003). During calendar year 2010, environmental monitoring within the INL Site boundaries was primarily the responsibility of the INL and Idaho Cleanup Project (ICP) contractors, while monitoring outside the INL Site boundaries was conducted under the Environmental Surveillance, Education and Research (ESEER) Program. The ESEER Program is led by the S.M. Stoller Corporation in cooperation with its team members, including the University of Idaho, Idaho State University (ISU), the Wildlife Conservation Society, and Teledyne Brown Engineering. This report contains monitoring results from the ESEER Program for samples collected during the second quarter of 2010 (April 1-June 30, 2010).

The surveillance portion of the ESEER Program is designed to satisfy the following program objectives:

- Verify compliance with applicable environmental laws, regulations, and DOE Orders
- Characterize and define trends in the physical, chemical, and biological condition of environmental media on and around the INL Site
- Assess the potential radiation dose to members of the public from INL Site effluents
- Present program results clearly and concisely through the use of reports, presentations, newsletter articles and press releases.

The goal of the surveillance program is to monitor different media at a number of potential exposure points within the various exposure pathways, including air, water, agricultural products, wildlife, and soil that could possibly contribute to the radiation dose received by the public.

Environmental samples collected include:

- air at 16 locations on and around the INL Site
- moisture in air at four locations around the INL Site
- precipitation from three locations on and around the INL Site
- agricultural products, including milk at six dairies around the INL Site, potatoes from at least five local producers, wheat from approximately 10 local producers, and lettuce from approximately nine home-owned and portable gardens on and around the INL
- soil from 12 locations around the INL Site biennially
- environmental dosimeters from 15 locations semi-annually
- various numbers of wildlife including big game (pronghorn, mule deer, and elk) and waterfowl sampled on and near the INL Site.

Table A-1 in Appendix A lists samples, sampling locations and collection frequency for the ESEER Program.

The ESEER Program used two laboratories to perform analyses on routine environmental samples collected during the quarter reported here. The ISU Environmental Assessment Laboratory (EAL) performed routine gross alpha, gross beta, tritium, and gamma spectrometry analyses. Analyses requiring radiochemistry including strontium-90 (<sup>90</sup>Sr), plutonium-238

(<sup>238</sup>Pu), plutonium-239/240 (<sup>239/240</sup>Pu), and americium-241 (<sup>241</sup>Am) were performed by Teledyne Brown Engineering, Inc. of Knoxville, Tennessee.

In the event of non-routine occurrences, such as suspected releases of radioactive material, the ESER Program may increase the frequency of sampling and/or the number of sampling locations based on the nature of the release and wind distribution patterns. Any data found to be outside historical norms in the ESER Program is thoroughly investigated to determine if an INL Site origin is likely. Investigation may include re-sampling and/or re-analysis of prior samples.

In the event of any suspected worldwide nuclear incidents, like the 1986 Chernobyl accident, the EPA may request additional sampling be performed through RadNet [previously known as the Environmental Radiation Ambient Monitoring System (ERAMS) network] (EPA 2009). The EPA established the ERAMS network in 1973 with an emphasis on identifying trends in the accumulation of long-lived radionuclides in the environment. ERAMS was renamed RadNet in 2005 to reflect a new mission. RadNet is comprised of a nationwide network of sampling stations that provide air, precipitation, drinking water, and milk samples. The ESER Program currently operates a high-volume air sampler and collects precipitation and drinking water in Idaho Falls for this national program and routinely sends samples to EPA's Eastern Environmental Radiation Facility for analyses. The RadNet data collected at Idaho Falls are not reported by the ESER Program but are available through the EPA RadNet website (<http://www.epa.gov/narel/radnet/>).


Once samples have been collected and analyzed, the ESER Program has the responsibility for quality control of the data and for preparing quarterly reports on results from the environmental surveillance program. The quarterly reports are then consolidated into the INL Site Environmental Report for each calendar year. These annual reports also include data collected by other INL Site contractors.

The results reported in the quarterly and annual reports are assessed in terms of data quality and statistical significance with respect to laboratory analytical uncertainties, sample locations, reported INL Site releases, meteorological data, and worldwide events that might conceivably have an effect on the INL Site environment. First, field collection and laboratory information are reviewed to determine identifiable errors that would invalidate or limit use of the data. Examples of such limitations include insufficient sample volume, torn filters, evidence of laboratory cross-contamination or quality control issues. Data that pass initial screening are further evaluated using statistical methods. Statistical tools are necessary for data evaluation particularly since environmental measurements typically involve the determination of minute concentrations, which are difficult to detect and even more difficult to distinguish from other measurements.

Results are presented in this report with an analytical uncertainty term, *s*, where "s" is the estimated sample standard deviation ( $\sigma$ ), assuming a Gaussian or normal distribution. All results are reported in this document, even those that do not necessarily represent detections. The term "detected", as used for the discussion of results in this report, does not imply any degree of risk to the public or environment, but rather indicates that the radionuclide was measured at a concentration sufficient for the analytical instrument to record a value that is statistically different from background. The ESER has adopted guidelines developed by the United States Geological Survey (Bartholomay, et al. 2003), based on an extension of a method proposed by Currie (1984), to interpret analytical results and make decisions concerning detection. Most of the following discussion is taken from Bartholomay et al (2003).



Laboratory measurements involve the analysis of a target sample and the analysis of a prepared laboratory blank (i.e., a sample which is identical to the sample collected in the environment, except that the radionuclide of interest is absent). Instrument signals for the target and blank vary randomly about the true signals and may overlap making it difficult to distinguish between radionuclide activities in blank and in environmental samples (Figure 1). That is, the variability around the sample result may substantially overlap the variability around a net activity of zero for samples with no radioactivity. In order to conclude that a radionuclide has been detected, it is essential to consider two fundamental aspects of the problem of detection: (1) the instrument signal for the sample must be greater than that observed for the blank before the decision can be made that the radionuclide has been detected; and (2) an estimate must be made of the minimum radionuclide concentration that will yield a sufficiently large observed signal before the correct decision can be made for detection or non-detection.



**Example of overlap of blank and sample measurement distributions.**

In the laboratory, instrument signals must exceed a critical level of  $1.6s$  before the qualitative decision can be made as to whether the radionuclide was detected in a sample. At  $1.6s$  there is about a 95-percent probability that the correct conclusion—not detected—will be made. Given a large number of samples, approximately 5 percent of the samples with measured concentrations greater than or equal to  $1.6s$ , which were concluded as being detected, might not contain the radionuclide. These are referred to as false positives. For purposes of simplicity and consistency with past reporting, the ESER has rounded the  $1.6s$  critical level estimate to  $2s$ .

Once the critical level has been defined, the minimum detectable concentration may be determined. Concentrations that equal  $3s$  represent a measurement at the detection level or minimum detectable concentration. For true concentrations of  $3s$  or greater, there is a 95-percent probability that the radionuclide was detected in the target sample. In a large number of samples, the conclusion—not detected—will be made in 5 percent of the samples with true concentrations at the minimum detectable concentration of  $3s$ . These measurements are known as false negatives. The ESER reports measured radionuclide concentrations greater than or equal to their respective  $3s$  uncertainties as being “detected with confidence.”

Concentrations between  $2s$  and  $3s$  are reported as “questionably detected”. That is, the radionuclide may be present in the sample; however, the detection may not be reliable.

Measurements made between 2s and 3s are examined further to determine if they are a part of a pattern (temporal or spatial) that might warrant further investigation or recounting. For example, if a particular radionuclide is typically detected at > 3s at a specific location, a sample result between 2s and 3s might be considered detected.

If a result is less than or equal to 2s there is little confidence that the radionuclide is present in the sample. Analytical results in this report are presented as the result value  $\pm$  one standard deviation (1s) for reporting consistency with the annual report. To obtain the 2s or 3s values simply multiply the uncertainty term by 2 or 3.

For more information concerning the ESER Program, contact the S.M. Stoller Corporation at (208) 525-9358, or visit the Program's web page (<http://www.stoller-eser.com>).

## 2. THE INL SITE

The INL Site is a nuclear energy and homeland security research and environmental management facility. It is owned and administered by the U.S. Department of Energy, Idaho Operations Office (DOE-ID) and occupies about 890 mi<sup>2</sup> (2300 km<sup>2</sup>) of the upper Snake River Plain in Southeastern Idaho. The history of the INL Site began during World War II when the U.S. Naval Ordnance Station was located in Pocatello, Idaho. This station, one of two such installations in the U.S., retooled large guns from U.S. Navy warships. The retooled guns were tested on the nearby, uninhabited plain, known as the Naval Proving Ground. In the years following the war, as the nation worked to develop nuclear power, the Atomic Energy Commission (AEC), predecessor to the DOE, became interested in the Naval Proving Ground and made plans for a facility to build, test, and perfect nuclear power reactors.

The Naval Proving Ground became the National Reactor Testing Station (NRTS) in 1949, under the AEC. By the end of 1951, a reactor at the NRTS became the first to produce useful amounts of electricity. Over time the site has operated 52 various types of reactors, associated research centers, and waste handling areas. The NRTS was renamed the Idaho National Engineering Laboratory (INEL) in 1974, and the Idaho National Engineering and Environmental Laboratory (INEEL) in January 1997. With renewed interest in nuclear power the DOE announced in 2003 that Argonne National Laboratory and the INEEL would be the lead laboratories for development of the next generation of power reactors. On February 1, 2005 the INEEL and Argonne National Laboratory-West became the INL. The INL is committed to providing international nuclear leadership for the 21st Century, developing and demonstrating compelling national security technologies, and delivering excellence in science and technology as one of the Department of Energy's multiprogram national laboratories.

The cleanup operation, the ICP, is now a separately managed effort. The ICP is charged with safely and cost-effectively completing the majority of cleanup work from past laboratory missions by 2012.




### 3. AIR SAMPLING

The primary pathway by which radionuclides can move off the INL Site is through the air and for this reason the air pathway is the primary focus of monitoring on and around the INL Site. Samples for particulates and iodine-131 ( $^{131}\text{I}$ ) gas in air were collected weekly for the duration of the quarter at 16 locations using low-volume air samplers. Moisture in the atmosphere was sampled at four locations around the INL Site and analyzed for tritium. Air sampling activities and results for the second quarter of 2010 are discussed below. A summary of approximate minimum detectable concentrations (MDCs) for radiological analyses and DOE Derived Concentration Guide (DCG) (DOE 1993) values is provided in Appendix B.

#### LOW-VOLUME AIR SAMPLING

Radioactivity associated with airborne particulates was monitored continuously by 18 low-volume air samplers (two of which are used as replicate samplers) at 16 locations during the second quarter of 2010 (Figure 2). Four of these samplers are located on the INL Site, seven are situated off the INL Site near the boundary, and seven have been placed at locations distant to the INL Site. Samplers are divided into INL Site, Boundary, and Distant groups to determine if there is a gradient of radionuclide concentrations, increasing towards the INL Site. Each replicate sampler is relocated every other year to a new location. During 2010, one replicate sampler was operating in Dubois (a Distant location) and one was operating at the Van Buren gate (an INL Site location). An average of 18,308 ft<sup>3</sup> (518 m<sup>3</sup>) of air was sampled at each location, each week, at an average flow rate of 1.82 ft<sup>3</sup>/min (0.05 m<sup>3</sup>/min). Particulates in air were collected on membrane particulate filters (1.2- $\mu\text{m}$  pore size). Gases passing through the filter were collected with an activated charcoal cartridge.



Low-volume air sampler locations.

Filters and charcoal cartridges were changed weekly at each station during the quarter. Each particulate filter was analyzed for gross alpha and gross beta radioactivity using thin-window gas flow proportional counting systems after waiting about four days for naturally-occurring daughter products of radon and thorium to decay.

The weekly particulate filters collected during the quarter for each location were composited and analyzed for gamma-emitting radionuclides. Selected composites were also analyzed by location for  $^{90}\text{Sr}$ ,  $^{238}\text{Pu}$ ,  $^{239/240}\text{Pu}$ , and  $^{241}\text{Am}$  as determined by a rotating quarterly schedule.

Charcoal cartridges were analyzed for gamma-emitting radionuclides, specifically for iodine-131 ( $^{131}\text{I}$ ). Iodine-131 is of particular interest because it is produced in relatively large quantities by nuclear fission, is readily accumulated in human and animal thyroids, and has a half-life of eight days. This means that any elevated level of  $^{131}\text{I}$  in the environment could be from a recent release of fission products.

Gross alpha results are reported in Table C-1. Median gross alpha concentrations in air for INL Site, Boundary, and Distant locations for the second quarter of 2010 are shown in Figure 3. Gross alpha data are tested for normality prior to statistical analyses, and generally show no consistent discernable distribution. Box and whisker plots are commonly used when there is no assumed distribution. Each data group in Figure 3 is presented as a box and whisker plot, with a median (small red square), a box enclosing values between the 25<sup>th</sup> and 75<sup>th</sup> percentiles, and whiskers representing the non-outlier range. Outliers and extreme values are identified separately from the box and whiskers. Outliers and extreme values are atypical, infrequent data points that are far from the middle of the data distribution. For this report, outliers are defined as values that are greater than 1.5 times the height of the box, above, or below the box. Extreme values are greater than 2 times the height of the box, above or below the box. Outliers and extreme values may reflect inherent variability, may be due to errors associated with transcription or measurement, or may be related to other anomalies. A careful review of the data collected during the second quarter indicates that the outlier values were not due to mistakes in collection, analysis, or reporting procedures, but rather reflect natural variability in the measurements. Thus, rather than dismissing the outliers, they were included in the subsequent statistical analyses.

Because there is no discernable distribution of the data, the nonparametric Kruskal-Wallis test of multiple independent groups was used to test for statistical differences between INL Site, Boundary, and Distant locations. The use of nonparametric tests, such as Kruskal-Wallis, gives less weight to outlier and extreme values thus allowing a more appropriate comparison of data groups. A statistically significant difference exists between data groups if the (p) value is less than 0.05. Values greater than 0.05 translate into a 95 percent confidence that the medians are statistically the same. The p-value for each comparison is shown in Table D-1. For the quarter, there was no statistical difference noted. Figure 3 graphically shows that the gross alpha measurements made at INL Site, Boundary, and Distant locations are similar for the second quarter. If the INL Site were a significant source of offsite contamination, concentrations of contaminants could be statistically greater at Boundary locations than at Distant locations.

Comparisons of gross alpha concentrations were made for each month of the quarter (Figures 4 – 6). Again the Kruskal-Wallis test of multiple independent groups was used to determine if statistical differences exist between INL Site, Boundary, and Distant data groups. No statistical differences in gross alpha concentrations between groups were noted during any month of the quarter (Table D-1).




Figure 3. Gross alpha concentrations in air at ESER INL Site, Boundary and Distant locations for the second quarter of 2010.




Figure 4. April gross alpha concentrations in air at ESER INL Site, Boundary, and Distant locations. Number of samples (N) = 4 at each location.






Figure 5. May gross alpha concentrations in air at ESER INL Site, Boundary, and Distant locations. Number of samples (N) = 4 at each location, except Blackfoot (N = 3).




Figure 6. June gross alpha concentrations in air at ESER INL Site, Boundary, and Distant locations. Number of samples (N) = 5 at each location, except Blackfoot (N = 4).

As an additional check, comparisons between gross alpha concentrations measured at Boundary and Distant locations were made on a weekly basis. The Mann-Whitney U test was used to compare the Boundary and Distant data because it is the most powerful nonparametric alternative to the t-test for independent samples. INL Site sample results were not included in this analysis because the onsite data, collected at only three locations, are not representative of the entire INL Site and would not aid in determining offsite impacts. In the second quarter, there were no weeks where a statistical difference existed between the two sample groups (Table D-2).

Gross beta results are presented in Table C-1. Gross beta concentrations in air for INL Site, Boundary, and Distant locations for the second quarter of 2010 are shown in Figure 7. The data were tested and found to be neither normally nor log-normally distributed. Box and whiskers plots were used for presentation of the data. Outliers and extreme values were retained in subsequent statistical analyses because they are within the range of measurements made in the past five years, and because these values could not be attributed to mistakes in collection, analysis, or reporting procedures. No statistical differences were noted in the quarterly data using the Kruskal-Wallis test (Table D-1).

Monthly median gross beta concentrations in air for each sampling group are shown in Figures 8 – 10. Statistical data are presented in Table D-1. No statistical differences were noted during any month.

Comparison of weekly Boundary and Distant gross beta data sets, using the Mann Whitney U test, showed no statistical differences between Boundary and Distant measurements during any week of the quarter (Table D-1).

Iodine-131 was initially reported at just above the detection level on one charcoal cartridge batch during the week of May 19. Two subsequent recounts failed to confirm the presence of  $^{131}\text{I}$ . Weekly  $^{131}\text{I}$  results for each location are listed in Table C-2 of Appendix C. Gamma spectrographic analysis is also done with the  $^{131}\text{I}$  analysis. Cesium-137 was not detected in any of the 26 measured batches of cartridges this quarter.

Weekly filters for the second quarter of 2010 were composited by location. All samples were analyzed for gamma-emitting radionuclides, including  $^{137}\text{Cs}$ . Selected composites were also analyzed for  $^{90}\text{Sr}$ ,  $^{238}\text{Pu}$ ,  $^{239/240}\text{Pu}$ , and  $^{241}\text{Am}$  (see Table C-3, Appendix C). None of these radionuclides were found on any of the composites.

#### **ATMOSPHERIC MOISTURE SAMPLING**

Seventeen atmospheric moisture samples were obtained during the second quarter of 2010 from Atomic City, Blackfoot, Idaho Falls, and Rexburg. Atmospheric moisture is collected by pulling air through a column of absorbent material (molecular sieve material) to absorb water vapor. The water is then extracted from the absorbent material by heat distillation. The resulting water samples are then analyzed for tritium using liquid scintillation.

Nine of the 17 samples exceeded the 3s uncertainty level for tritium, with similar results to those reported previously. All samples were significantly below the DOE DCG for tritium in air of  $1 \times 10^{-7} \mu\text{Ci}/\text{mL}_{\text{air}}$  with a maximum reported value of  $15.2 \times 10^{-13} \mu\text{Ci}/\text{mL}_{\text{air}}$  at Rexburg. All results are shown in Table C-4, Appendix C.




Figure 7. Gross beta concentrations in air at ESER INL Site, Boundary, and Distant locations for the second quarter of 2010.




Figure 8. April gross beta concentrations in air at ESER INL Site, Boundary, and Distant locations. Number of samples (N) = 4 at each location.




Figure 9. May gross beta concentrations in air at ESER INL Site, Boundary and Distant locations. Number of samples (N) = 4 at each location, except Blackfoot (N = 3).




Figure 10. June gross beta concentrations in air at ESER INL Site, Boundary, and Distant locations. Number of samples (N) = 5 at each location, except Blackfoot (N = 4).





## 4. PRECIPITATION AND SURFACE WATER SAMPLING

### ***PRECIPITATION SAMPLING***

Precipitation samples are gathered when sufficient precipitation occurs to allow for the collection of the minimum sample volume of approximately 50 mL. Samples are taken of monthly composites from Idaho Falls and CFA, and weekly from the EFS. Precipitation samples are analyzed for tritium. Storm events in the second quarter of 2010 produced sufficient precipitation to yield 16 samples; results were available for 13 of the samples.

Tritium was measured above the 3s values in 5 of the 13 samples for which results were available. Low levels of tritium exist in the environment at all times as a result of cosmic ray reactions with water molecules in the upper atmosphere. The EPA's RadNet program collects precipitation samples from across the United States. From 1980 to 2008, tritium measured in samples from Region 10 (which includes Idaho) averaged 117 pCi/L (EPA 2009). Data for all second quarter precipitation samples collected by the ESER Program were in the range of this value (averaging 84 pCi/L) and are listed in Table C-5 (Appendix C).

### ***SURFACE WATER SAMPLING***

The Big Lost River contained water on the INL Site for a short period during the second quarter of 2010. Samples were collected at four locations and analyzed for gross alpha activity, gross beta activity, and tritium. Collection locations included the public rest area on US Highway 20/26, on Lincoln Boulevard near the Idaho Nuclear Technology and Engineering Center, at EFS (plus a duplicate sample), and on Lincoln Boulevard near the Naval Reactors Facility.

Gross alpha was detected in all of the samples and gross beta was detected in four of the five samples analyzed. It is not unusual to detect these constituents in water of the Snake River Plain, related to natural production from the basalts that make up the aquifer. In 2006, the last year in which the ESER Program sampled offsite surface water, gross alpha concentrations ranged up to 3.77 pCi/L and gross beta concentrations ranged up to 8.82 pCi/L. All results from the Big Lost River sampling were well within these ranges (see Appendix C, Table C-6).

Tritium was detected in four of the five samples. As described in the Precipitation section above, an average of 117 pCi/L was detected in water taken across the region. The samples from the Big Lost River averaged 116 pCi/L, or nearly exactly the same as the background results (Appendix C, Table C-6).



## 5. AGRICULTURAL PRODUCT, WILDLIFE AND SOIL SAMPLING

Another potential pathway for contaminants to reach humans is through the food chain. The ESER Program samples multiple agricultural products and game animals from around the INL Site and Southeast Idaho. Specifically, milk, wheat, potatoes, garden lettuce, large game animals, and waterfowl are sampled. Milk is sampled throughout the year and large game animals are sampled whenever large game animals are killed onsite from vehicle collisions. Lettuce and wheat are sampled during the third quarter, while potatoes are collected during the fourth quarter. Waterfowl are collected in either the third or fourth quarter. See Table A-1, Appendix A, for more details on agricultural product and wildlife sampling. This section discusses results from milk and a special waterfowl sample collected during the second quarter of 2010.

### MILK SAMPLING

Milk samples were collected weekly in Idaho Falls. Monthly samples were collected at six other locations around the INL Site (Figure 11) during the second quarter of 2010. In addition, commercially-available organic milk was purchased as a control sample. All samples were analyzed for gamma emitting radionuclides. No Iodine-131 or other gamma-emitting radionuclides were detected in any sample. Data for  $^{131}\text{I}$  and  $^{137}\text{Cs}$  in milk samples are listed in Appendix C, Table C-7.

Strontium-90 was detected in all eight samples analyzed (including the control sample) at levels within historical measurements, ranging from 0.36 pCi/L (Dietrich) to 1.84 pCi/L (Rupert). Tritium was not detected in any of the samples analyzed (see Table C-8 in Appendix C).





Figure 11. ESER milk sampling locations.

### LARGE GAME ANIMAL SAMPLING

Five large game animals (three pronghorn and two mule deer) were available for sampling during the second quarter. Man-made radionuclides were not detected in any of the tissue samples collected. Results are presented in Appendix C, Table C-9.

## 6. ENVIRONMENTAL RADIATION

An array of thermoluminescent dosimeters (TLDs) is distributed throughout the Eastern Snake River Plain to monitor for environmental radiation (Figure 12). TLDs are changed out in May and again in November after six months in the field. The results of the TLDs exposed from November 2009 to May 2010 are discussed below.



**Figure 1. TLD locations.**

Similar to the low-volume air results the environmental dosimeter locations are also divided into Boundary and Distant groupings. Boundary average exposure rates ranged from a low of 0.30 mR/day at Blue Dome to a high of 0.38 mR/day at Mud Lake. The overall Boundary average was 0.34 mR/day. The Distant group had a high of 0.40 mR/day at Rexburg and a low of 0.29 mR/day at the Dubois location. The overall average Distant value was 0.35 mR/day. There was no statistical difference between Boundary and Distant locations and all values are consistent with past readings. All results are listed in Appendix C, Table C-10.

## 7. QUALITY ASSURANCE

The ESER Quality Assurance Program consists of five ongoing tasks which measure:

1. method uncertainty
2. data completeness
3. data accuracy, using spike, performance evaluation and laboratory control samples
4. data precision, using split samples, duplicate samples and recounts
5. presence of contamination in samples, using blanks.

Data completeness for sample collection and delivery was 100 percent during the second quarter. There were two air samples that had volumes below the 7000 ft<sup>3</sup> (200 m<sup>3</sup>) threshold for a valid sample. Both were at the Blackfoot location. One involved the station manager unplugging the sampler and the second resulted when the school sprinkler system caused the sampler GFCI to trip repeatedly.

Sample results are compared to criteria described in the Quality Assurance Project Plan for the INL Site Offsite Environmental Surveillance Program (Stoller 2007). The following table summarizes the results of the quality assurance program for the second quarter of 2010.

<b>QA Sample Type</b>	<b>Number of Sample Results</b>	<b>Number of Results Meeting Criteria</b>	<b>Percentage Meeting Criteria</b>
Spikes/Laboratory Control Samples	169	169	100.0
Field Duplicates	66	63	95.5
Laboratory Splits	29	28	96.6
Recounts	185	185	100
Field Blanks	70	70	100
Method Uncertainty	1830	1801	98.4



---

## 8. REFERENCES

- Bartholomay, R.C., Knobel, L.L., and Rousseau, J.P., 2003, *Field Methods and Quality Plan for Quality-of-Water Activities, U.S. Geological Survey, Idaho National Engineering and Environmental Laboratory, Idaho*, DOE/ID-22182, January 2003.
- Currie, L.A., 1984, *Lower Limit of Detection: Definition and Elaboration of a Proposed Position for Radiological Effluent and Environmental Measurements*, NUREG/CR-4007, U.S. Nuclear Regulatory Commission, Washington, D.C., September 1984.
- DOE, 2003, "Environmental Management System," U.S. Department of Energy Order 450.1, January 2003.
- DOE, 1993, "Radiation Protection of the Public and the Environment," U.S. Department of Energy Order 5400.5, January 1993.
- EPA, 2009, RadNet—Tracking Environmental Radiation Nationwide, Web-page: <http://www.epa.gov/narel/radnet/>
- Stoller, 2007, *Quality Assurance Project Plan for the INL Site Offsite Environmental Surveillance Program*, Environmental Surveillance, Education and Research Program, February 2007.





**APPENDIX A**  
***SUMMARY OF SAMPLING SCHEDULE***



**Table A-1. Summary of the ESER Program's Sampling Schedule**

Sample Type Analysis	Collection Frequency	LOCATIONS		
		Distant	Boundary	INL Site
<b>AIR SAMPLING</b>				
<i>LOW-VOLUME AIR</i>				
Gross Alpha, Gross Beta, <sup>131</sup> I	weekly	Blackfoot, Craters of the Moon, Dubois, Idaho Falls, Jackson WY, Rexburg	Arco, Atomic City, FAA Tower, Howe, Montevieu, Mud Lake, Blue Dome	Main Gate, EFS, Van Buren
Gamma Spec	quarterly	Blackfoot, Craters of the Moon, Dubois, Idaho Falls, Jackson WY, Rexburg	Arco, Atomic City, FAA Tower, Howe, Montevieu, Mud Lake, Blue Dome	Main Gate, EFS, Van Buren
<sup>90</sup> Sr, Transuranics	quarterly	Rotating schedule	Rotating schedule	Rotating schedule
<i>ATMOSPHERIC MOISTURE</i>				
Tritium	2 to 13 weeks	Blackfoot, Idaho Falls, Rexburg	Atomic City	None
<i>PRECIPITATION</i>				
Tritium	monthly	Idaho Falls	None	CFA
Tritium	weekly	None	None	EFS
<b>ENVIRONMENTAL RADIATION SAMPLING</b>				
<i>TLDs</i>				
Gamma Radiation	semiannual	Aberdeen, Blackfoot (2), Craters of the Moon, Dubois, Idaho Falls, Jackson WY, Minidoka, Rexburg, Roberts	Arco, Atomic City, Birch Creek, Blue Dome, Howe, Montevieu, Mud Lake	None
<b>SOIL SAMPLING</b>				
<i>SOIL</i>				
Gamma Spec, <sup>90</sup> Sr, Transuranics	biennially	Carey, Crystal Ice Caves (Aberdeen), Blackfoot, St. Anthony	Butte City, Montevieu, Atomic City, FAA Tower, Howe, Mud Lake (2), Birch Creek	None

**Table A-1. Summary of the ESER Program's Sampling Schedule (continued)**

Sample Type Analysis	Collection Frequency	LOCATIONS		
		Distant	Boundary	INL Site
<b>FOODSTUFF SAMPLING</b>				
<i>MILK</i>				
Gamma Spec ( <sup>131</sup> I)	weekly	Idaho Falls	None	None
Gamma Spec ( <sup>131</sup> I)	monthly	Blackfoot, Dietrich, Fort Hall, Idaho Falls, Minidoka	Howe, Terreton	None
Tritium, <sup>90</sup> Sr	Semi-annually	Blackfoot, Dietrich, Fort Hall, Idaho Falls, Minidoka	Howe, Terreton	None
<i>POTATOES</i>				
Gamma Spec, <sup>90</sup> Sr	annually	Aberdeen, Blackfoot, Fort Hall, Idaho Falls, Rupert, Taber, occasional samples across the U.S.	Arco, Monteview, Mud Lake, Terreton	None
<i>WHEAT</i>				
Gamma Spec, <sup>90</sup> Sr	annually	American Falls, Blackfoot, Dietrich, Idaho Falls, Minidoka, Carey	Arco, Monteview, Mud Lake, Taber, Terreton	None
<i>LETTUCE</i>				
Gamma Spec, <sup>90</sup> Sr	annually	Blackfoot, Carey, Idaho Falls	Arco, Atomic City, FAA Tower, Howe, Monteview	EFS
<i>BIG GAME</i>				
Gamma Spec	varies	Occasional samples across the U.S.	Public Highways	INL Site roads
<i>WATERFOWL</i>				
Gamma Spec, <sup>90</sup> Sr, Transuranics	annually	Varies among: Heise, Firth, Fort Hall, Mud Lake, Market Lake, and American Falls	None	Wastewater disposal ponds

**APPENDIX B**  
***SUMMARY OF MDCs AND DCGs***



**Table B-1. Summary of Approximate Minimum Detectable Concentrations for Radiological Analyses Performed During Second Quarter 2010**

Sample Type	Analysis	Approximate Minimum Detectable Concentration <sup>a</sup> (MDC)	Derived Concentration Guide <sup>b</sup> (DCG)
<b>Air</b> (particulate filter) <sup>e</sup>	Gross alpha <sup>c</sup>	$5.97 \times 10^{-16}$ $\mu\text{Ci/mL}$	$2 \times 10^{-14}$ $\mu\text{Ci/mL}$
	Gross beta <sup>d</sup>	$1.78 \times 10^{-15}$ $\mu\text{Ci/mL}$	$3 \times 10^{-12}$ $\mu\text{Ci/mL}$
	Specific gamma ( <sup>137</sup> Cs)	$1.75 \times 10^{-16}$ $\mu\text{Ci/mL}$	$4 \times 10^{-10}$ $\mu\text{Ci/mL}$
	<sup>238</sup> Pu	$1.33 \times 10^{-18}$ $\mu\text{Ci/mL}$	$3 \times 10^{-14}$ $\mu\text{Ci/mL}$
	<sup>239/240</sup> Pu	$7.51 \times 10^{-19}$ $\mu\text{Ci/mL}$	$2 \times 10^{-14}$ $\mu\text{Ci/mL}$
	<sup>241</sup> Am	$3.46 \times 10^{-18}$ $\mu\text{Ci/mL}$	$2 \times 10^{-14}$ $\mu\text{Ci/mL}$
	<sup>90</sup> Sr	$2.80 \times 10^{-17}$ $\mu\text{Ci/mL}$	$9 \times 10^{-12}$ $\mu\text{Ci/mL}$
<b>Air</b> (charcoal cartridge) <sup>e</sup>	<sup>131</sup> I	$1.38 \times 10^{-15}$ $\mu\text{Ci/mL}$	$4 \times 10^{-10}$ $\mu\text{Ci/mL}$
<b>Air</b> (atmospheric moisture)	<sup>3</sup> H	122.6 pCi/L <sub>water</sub>	$1 \times 10^{-7}$ $\mu\text{Ci/mL}_{\text{air}}$
<b>Air</b> (precipitation)	<sup>3</sup> H	122.7 pCi/L	$2 \times 10^{-3}$ $\mu\text{Ci/mL}$
<b>Milk</b>	<sup>131</sup> I	0.63 pCi/L	--
	<sup>137</sup> Cs	1.07 pCi/L	--
	<sup>90</sup> Sr	0.43 pCi/L	--
<p>a The MDC is an estimate of the concentration of radioactivity in a given sample type that can be identified with a 95 percent level of confidence and precision of plus or minus 100 percent under a specified set of typical laboratory measurement conditions.</p> <p>b DCGs, set by the DOE, represent reference values for radiation exposure. They are based on a radiation dose of 100 mrem/yr for exposure through a particular exposure mode such as direct exposure, inhalation, or ingestion of water.</p> <p>c The DCG for gross alpha is equivalent to the DCGs for <sup>239,240</sup>Pu and <sup>241</sup>Am.</p> <p>d The DCG for gross beta is equivalent to the DCGs for <sup>228</sup>Ra</p> <p>e The approximate MDC is based on an average filtered air volume (pressure corrected) of 445 m<sup>3</sup>/week.</p>			





**APPENDIX C**  
***SAMPLE ANALYSIS RESULTS***



TABLE C-1. Weekly Gross Alpha and Gross Beta Concentrations in Air

Sampling Group and Location	Sampling Date	GROSS ALPHA							GROSS BETA				
		Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)		Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)			Result > 3s	Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)		Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)		Result > 3s	
ARCO	4/7/2010	0.55	± 0.18	2.02	± 0.65	Y	10.70	± 0.59	39.59	± 2.17	Y		
	4/14/2010	1.46	± 0.23	5.40	± 0.85	Y	20.80	± 0.76	76.96	± 2.81	Y		
	4/21/2010	1.51	± 0.20	5.59	± 0.74	Y	29.90	± 0.74	110.63	± 2.73	Y		
	4/28/2010	0.96	± 0.20	3.56	± 0.74	Y	19.60	± 0.73	72.52	± 2.71	Y		
	5/5/2010	0.79	± 0.18	2.91	± 0.67	Y	9.21	± 0.59	34.08	± 2.17	Y		
	5/12/2010	0.73	± 0.19	2.68	± 0.72	Y	14.90	± 0.67	55.13	± 2.49	Y		
	5/19/2010	1.20	± 0.22	4.44	± 0.80	Y	26.90	± 0.82	99.53	± 3.04	Y		
	5/26/2010	0.64	± 0.18	2.36	± 0.66	Y	13.10	± 0.64	48.47	± 2.37	Y		
	6/2/2010	0.54	± 0.20	1.99	± 0.74		11.80	± 0.65	43.66	± 2.39	Y		
	6/9/2010	0.68	± 0.19	2.52	± 0.70	Y	16.80	± 0.70	62.16	± 2.59	Y		
	6/16/2010	0.48	± 0.19	1.76	± 0.71		19.80	± 0.74	73.26	± 2.73	Y		
	6/23/2010	0.81	± 0.21	2.99	± 0.76	Y	19.90	± 0.78	73.63	± 2.88	Y		
	6/30/2010	0.95	± 0.23	3.52	± 0.84	Y	29.60	± 0.91	109.52	± 3.35	Y		
	ATOMIC CITY	4/7/2010	0.56	± 0.19	2.07	± 0.71		11.40	± 0.64	42.18	± 2.36	Y	
4/14/2010		1.28	± 0.22	4.74	± 0.81	Y	19.50	± 0.73	72.15	± 2.72	Y		
4/21/2010		1.43	± 0.21	5.29	± 0.78	Y	30.70	± 0.80	113.59	± 2.95	Y		
4/28/2010		1.02	± 0.20	3.77	± 0.73	Y	18.90	± 0.71	69.93	± 2.63	Y		
5/5/2010		0.28	± 0.16	1.05	± 0.58		9.69	± 0.61	35.85	± 2.25	Y		
5/12/2010		0.58	± 0.17	2.15	± 0.63	Y	15.70	± 0.63	58.09	± 2.35	Y		
5/19/2010		1.48	± 0.26	5.48	± 0.95	Y	26.70	± 0.91	98.79	± 3.37	Y		
5/26/2010		0.63	± 0.19	2.32	± 0.70	Y	13.00	± 0.67	48.10	± 2.49	Y		
6/2/2010		0.36	± 0.23	1.33	± 0.83		14.00	± 0.77	51.80	± 2.84	Y		
6/9/2010		0.68	± 0.19	2.50	± 0.68	Y	19.50	± 0.72	72.15	± 2.65	Y		
6/16/2010		0.66	± 0.22	2.45	± 0.83		18.20	± 0.79	67.34	± 2.91	Y		
6/23/2010		0.42	± 0.16	1.55	± 0.58		20.40	± 0.68	75.48	± 2.52	Y		
6/30/2010		1.28	± 0.27	4.74	± 0.99	Y	31.20	± 1.01	115.44	± 3.74	Y		
BLUE DOME		4/7/2010	0.36	± 0.16	1.32	± 0.58		12.30	± 0.58	45.51	± 2.16	Y	
	4/14/2010	1.04	± 0.20	3.85	± 0.75	Y	18.80	± 0.71	69.56	± 2.63	Y		
	4/21/2010	1.53	± 0.19	5.66	± 0.70	Y	27.80	± 0.68	102.86	± 2.50	Y		
	4/28/2010	0.83	± 0.18	3.08	± 0.68	Y	19.10	± 0.69	70.67	± 2.56	Y		
	5/5/2010	0.43	± 0.16	1.59	± 0.58		7.63	± 0.55	28.23	± 2.04	Y		
	5/12/2010	0.45	± 0.17	1.68	± 0.62		13.70	± 0.62	50.69	± 2.31	Y		
	5/19/2010	1.36	± 0.21	5.03	± 0.79	Y	24.50	± 0.77	90.65	± 2.83	Y		
	5/26/2010	0.69	± 0.18	2.56	± 0.65	Y	12.10	± 0.61	44.77	± 2.26	Y		
	6/2/2010	0.07	± 0.17	0.26	± 0.63		12.60	± 0.63	46.62	± 2.35	Y		
	6/9/2010	0.65	± 0.20	2.42	± 0.75	Y	16.10	± 0.74	59.57	± 2.73	Y		
	6/16/2010	0.38	± 0.17	1.42	± 0.64		17.40	± 0.67	64.38	± 2.47	Y		
	6/23/2010	0.46	± 0.17	1.69	± 0.62		16.20	± 0.67	59.94	± 2.48	Y		
	6/30/2010	0.96	± 0.21	3.54	± 0.78	Y	28.10	± 0.84	103.97	± 3.09	Y		
	FAA TOWER	4/7/2010	0.63	± 0.19	2.32	± 0.72	Y	10.50	± 0.62	38.85	± 2.31	Y	
4/14/2010		1.14	± 0.22	4.22	± 0.83	Y	20.40	± 0.78	75.48	± 2.90	Y		
4/21/2010		1.51	± 0.20	5.59	± 0.73	Y	28.60	± 0.71	105.82	± 2.63	Y		
4/28/2010		0.75	± 0.25	2.78	± 0.94		20.60	± 0.97	76.22	± 3.58	Y		
5/5/2010		0.49	± 0.17	1.82	± 0.64		8.62	± 0.61	31.89	± 2.25	Y		
5/12/2010		0.39	± 0.19	1.45	± 0.68		13.20	± 0.68	48.84	± 2.52	Y		
5/19/2010		0.97	± 0.22	3.60	± 0.80	Y	27.70	± 0.86	102.49	± 3.20	Y		
5/26/2010		0.50	± 0.21	1.84	± 0.78		12.20	± 0.76	45.14	± 2.82	Y		
6/2/2010		0.23	± 0.19	0.84	± 0.69		14.20	± 0.68	52.54	± 2.52	Y		
6/9/2010		0.78	± 0.19	2.89	± 0.71	Y	17.40	± 0.70	64.38	± 2.58	Y		
6/16/2010		0.72	± 0.20	2.66	± 0.73	Y	16.00	± 0.68	59.20	± 2.50	Y		
6/23/2010		0.85	± 0.19	3.13	± 0.71	Y	18.60	± 0.71	68.82	± 2.64	Y		
6/30/2010		1.45	± 0.25	5.37	± 0.94	Y	28.60	± 0.91	105.82	± 3.37	Y		
HOWE		4/7/2010	0.43	± 0.16	1.60	± 0.57		10.70	± 0.54	39.59	± 2.00	Y	

TABLE C-1. Weekly Gross Alpha and Gross Beta Concentrations in Air

Sampling Group and Location	Sampling Date	GROSS ALPHA				GROSS BETA			
		Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)		Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)		Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)		Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)	
	4/14/2010	0.97 ± 0.18	3.58 ± 0.67	Y	19.70 ± 0.66	72.89 ± 2.45	Y		
	4/21/2010	1.31 ± 0.18	4.85 ± 0.65	Y	29.60 ± 0.68	109.52 ± 2.50	Y		
	4/28/2010	1.16 ± 0.19	4.29 ± 0.70	Y	18.70 ± 0.66	69.19 ± 2.43	Y		
	5/5/2010	0.52 ± 0.16	1.92 ± 0.57	Y	8.89 ± 0.55	32.89 ± 2.02	Y		
	5/12/2010	0.60 ± 0.17	2.21 ± 0.63	Y	15.10 ± 0.62	55.87 ± 2.29	Y		
	5/19/2010	0.90 ± 0.18	3.32 ± 0.67	Y	26.20 ± 0.74	96.94 ± 2.75	Y		
	5/26/2010	0.80 ± 0.17	2.97 ± 0.62	Y	13.60 ± 0.58	50.32 ± 2.15	Y		
	6/2/2010	0.56 ± 0.18	2.08 ± 0.66	Y	13.00 ± 0.59	48.10 ± 2.19	Y		
	6/9/2010	0.59 ± 0.16	2.18 ± 0.59	Y	17.20 ± 0.62	63.64 ± 2.29	Y		
	6/16/2010	0.60 ± 0.19	2.23 ± 0.69	Y	17.50 ± 0.68	64.75 ± 2.51	Y		
	6/23/2010	0.83 ± 0.17	3.09 ± 0.62	Y	20.20 ± 0.65	74.74 ± 2.39	Y		
	6/30/2010	1.43 ± 0.23	5.29 ± 0.85	Y	32.00 ± 0.86	118.40 ± 3.19	Y		
MONTEVIEW	4/7/2010	0.88 ± 0.19	3.24 ± 0.70	Y	13.00 ± 0.60	48.10 ± 2.22	Y		
	4/14/2010	1.60 ± 0.23	5.92 ± 0.85	Y	23.40 ± 0.77	86.58 ± 2.85	Y		
	4/21/2010	1.43 ± 0.19	5.29 ± 0.70	Y	28.60 ± 0.70	105.82 ± 2.60	Y		
	4/28/2010	0.99 ± 0.19	3.66 ± 0.71	Y	18.20 ± 0.69	67.34 ± 2.55	Y		
	5/5/2010	0.91 ± 0.19	3.36 ± 0.68	Y	8.77 ± 0.57	32.45 ± 2.12	Y		
	5/12/2010	1.02 ± 0.20	3.77 ± 0.73	Y	14.10 ± 0.63	52.17 ± 2.32	Y		
	5/19/2010	1.07 ± 0.21	3.96 ± 0.76	Y	25.10 ± 0.79	92.87 ± 2.90	Y		
	5/26/2010	0.48 ± 0.16	1.77 ± 0.60	Y	14.10 ± 0.63	52.17 ± 2.33	Y		
	6/2/2010	0.83 ± 0.23	3.09 ± 0.84	Y	15.10 ± 0.72	55.87 ± 2.68	Y		
	6/9/2010	0.84 ± 0.19	3.12 ± 0.71	Y	16.80 ± 0.68	62.16 ± 2.51	Y		
	6/16/2010	0.66 ± 0.24	2.43 ± 0.87	Y	16.60 ± 0.81	61.42 ± 2.99	Y		
	6/23/2010	1.07 ± 0.21	3.96 ± 0.78	Y	18.90 ± 0.75	69.93 ± 2.76	Y		
	6/30/2010	1.06 ± 0.22	3.92 ± 0.82	Y	30.80 ± 0.88	113.96 ± 3.26	Y		
MUD LAKE	4/7/2010	0.53 ± 0.17	1.95 ± 0.64	Y	12.10 ± 0.60	44.77 ± 2.21	Y		
	4/14/2010	1.47 ± 0.22	5.44 ± 0.82	Y	22.30 ± 0.75	82.51 ± 2.76	Y		
	4/21/2010	1.35 ± 0.20	5.00 ± 0.73	Y	31.70 ± 0.77	117.29 ± 2.83	Y		
	4/28/2010	1.07 ± 0.18	3.96 ± 0.68	Y	20.60 ± 0.67	76.22 ± 2.48	Y		
	5/5/2010	0.48 ± 0.17	1.78 ± 0.61	Y	8.54 ± 0.58	31.60 ± 2.15	Y		
	5/12/2010	0.65 ± 0.18	2.39 ± 0.66	Y	15.40 ± 0.65	56.98 ± 2.39	Y		
	5/19/2010	0.94 ± 0.20	3.49 ± 0.75	Y	27.80 ± 0.83	102.86 ± 3.07	Y		
	5/26/2010	0.73 ± 0.17	2.69 ± 0.63	Y	14.00 ± 0.60	51.80 ± 2.23	Y		
	6/2/2010	0.58 ± 0.20	2.14 ± 0.74	Y	15.30 ± 0.68	56.61 ± 2.52	Y		
	6/9/2010	0.94 ± 0.18	3.49 ± 0.68	Y	17.20 ± 0.64	63.64 ± 2.36	Y		
	6/16/2010	0.23 ± 0.19	0.85 ± 0.71	Y	18.50 ± 0.77	68.45 ± 2.84	Y		
	6/23/2010	0.92 ± 0.18	3.41 ± 0.65	Y	19.20 ± 0.65	71.04 ± 2.40	Y		
	6/30/2010	1.66 ± 0.28	6.14 ± 1.05	Y	33.30 ± 1.02	123.21 ± 3.77	Y		
<b>DISTANT</b>									
BLACKFOOT CMS	4/7/2010	0.63 ± 0.18	2.33 ± 0.67	Y	12.90 ± 0.62	47.73 ± 2.28	Y		
	4/14/2010	1.41 ± 0.22	5.22 ± 0.80	Y	21.60 ± 0.73	79.92 ± 2.70	Y		
	4/21/2010	1.93 ± 0.20	7.14 ± 0.75	Y	30.50 ± 0.69	112.85 ± 2.57	Y		
	4/28/2010	0.92 ± 0.19	3.39 ± 0.70	Y	18.90 ± 0.70	69.93 ± 2.59	Y		
	5/5/2010	0.47 ± 0.16	1.72 ± 0.60	Y	9.51 ± 0.59	35.19 ± 2.18	Y		
a	5/12/2010	0.88 ± 0.78	3.26 ± 2.88		11.30 ± 2.38	41.81 ± 8.81	Y		
	5/19/2010	1.17 ± 0.21	4.33 ± 0.76	Y	24.30 ± 0.76	89.91 ± 2.83	Y		
	5/26/2010	0.88 ± 0.19	3.26 ± 0.70	Y	14.80 ± 0.65	54.76 ± 2.42	Y		
	6/2/2010	0.61 ± 0.20	2.27 ± 0.75	Y	15.60 ± 0.68	57.72 ± 2.53	Y		
	6/9/2010	0.39 ± 0.22	1.44 ± 0.81	Y	14.80 ± 0.82	54.76 ± 3.05	Y		
	6/16/2010	0.67 ± 0.21	2.48 ± 0.76	Y	16.40 ± 0.72	60.68 ± 2.65	Y		
	6/23/2010	0.72 ± 0.27	2.67 ± 0.98	Y	19.30 ± 0.99	71.41 ± 3.67	Y		
a	6/30/2010	1.43 ± 0.54	5.29 ± 2.01		33.50 ± 1.91	123.95 ± 7.07	Y		
CRATERS OF THE MOON	4/7/2010	0.68 ± 0.18	2.52 ± 0.67	Y	10.40 ± 0.58	38.48 ± 2.14	Y		
	4/14/2010	0.70 ± 0.19	2.58 ± 0.69	Y	18.80 ± 0.72	69.56 ± 2.65	Y		
	4/21/2010	1.37 ± 0.19	5.07 ± 0.69	Y	27.10 ± 0.69	100.27 ± 2.54	Y		

TABLE C-1. Weekly Gross Alpha and Gross Beta Concentrations in Air

Sampling Group and Location	Sampling Date	GROSS ALPHA				GROSS BETA			
		Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)		Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)		Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)		Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)	
	4/28/2010	0.71 ± 0.19	2.64 ± 0.69	Y	17.80 ± 0.71	65.86 ± 2.63	Y		
	5/5/2010	0.59 ± 0.17	2.17 ± 0.62	Y	8.35 ± 0.57	30.90 ± 2.10	Y		
	5/12/2010	0.62 ± 0.19	2.29 ± 0.68	Y	13.80 ± 0.65	51.06 ± 2.40	Y		
	5/19/2010	0.81 ± 0.19	3.00 ± 0.71	Y	23.30 ± 0.77	86.21 ± 2.85	Y		
	5/26/2010	0.48 ± 0.18	1.78 ± 0.65		12.80 ± 0.66	47.36 ± 2.42	Y		
	6/2/2010	0.90 ± 0.21	3.32 ± 0.79	Y	11.30 ± 0.63	41.81 ± 2.33	Y		
	6/9/2010	0.48 ± 0.18	1.76 ± 0.65		16.10 ± 0.69	59.57 ± 2.53	Y		
	6/16/2010	0.81 ± 0.20	3.00 ± 0.74	Y	18.10 ± 0.70	66.97 ± 2.58	Y		
	6/23/2010	0.49 ± 0.17	1.82 ± 0.64		16.80 ± 0.70	62.16 ± 2.57	Y		
	6/30/2010	0.89 ± 0.21	3.30 ± 0.78	Y	29.40 ± 0.86	108.78 ± 3.18	Y		
DUBOIS	4/7/2010	0.98 ± 0.21	3.61 ± 0.78	Y	12.70 ± 0.65	46.99 ± 2.40	Y		
	4/14/2010	1.79 ± 0.24	6.62 ± 0.87	Y	21.40 ± 0.74	79.18 ± 2.73	Y		
	4/21/2010	1.06 ± 0.18	3.92 ± 0.65	Y	29.30 ± 0.72	108.41 ± 2.66	Y		
	4/28/2010	0.96 ± 0.21	3.57 ± 0.79	Y	20.90 ± 0.79	77.33 ± 2.93	Y		
	5/5/2010	0.19 ± 0.16	0.71 ± 0.60		7.75 ± 0.62	28.68 ± 2.29	Y		
	5/12/2010	0.79 ± 0.19	2.93 ± 0.69	Y	15.60 ± 0.65	57.72 ± 2.40	Y		
	5/19/2010	1.79 ± 0.25	6.62 ± 0.92	Y	28.90 ± 0.86	106.93 ± 3.18	Y		
	5/26/2010	0.73 ± 0.18	2.70 ± 0.67	Y	13.50 ± 0.63	49.95 ± 2.35	Y		
	6/2/2010	0.38 ± 0.22	1.41 ± 0.82		15.40 ± 0.77	56.98 ± 2.85	Y		
	6/9/2010	1.00 ± 0.22	3.70 ± 0.80	Y	18.50 ± 0.75	68.45 ± 2.79	Y		
	6/16/2010	0.56 ± 0.21	2.08 ± 0.76		16.10 ± 0.72	59.57 ± 2.68	Y		
	6/23/2010	1.12 ± 0.21	4.14 ± 0.76	Y	21.50 ± 0.74	79.55 ± 2.74	Y		
	6/30/2010	1.29 ± 0.24	4.77 ± 0.88	Y	31.20 ± 0.91	115.44 ± 3.35	Y		
QA-2 (DUBOIS)	4/7/2010	0.73 ± 0.18	2.70 ± 0.67	Y	11.50 ± 0.59	42.55 ± 2.17	Y		
	4/14/2010	1.45 ± 0.22	5.37 ± 0.83	Y	19.70 ± 0.73	72.89 ± 2.69	Y		
	4/21/2010	1.54 ± 0.19	5.70 ± 0.71	Y	28.60 ± 0.69	105.82 ± 2.55	Y		
	4/28/2010	0.67 ± 0.20	2.47 ± 0.73	Y	18.50 ± 0.76	68.45 ± 2.81	Y		
	5/5/2010	0.36 ± 0.15	1.32 ± 0.56		6.73 ± 0.53	24.90 ± 1.97	Y		
	5/12/2010	0.73 ± 0.19	2.70 ± 0.70	Y	15.10 ± 0.66	55.87 ± 2.43	Y		
	5/19/2010	1.17 ± 0.20	4.33 ± 0.73	Y	23.90 ± 0.73	88.43 ± 2.71	Y		
	5/26/2010	0.82 ± 0.22	3.02 ± 0.80	Y	12.40 ± 0.73	45.88 ± 2.69	Y		
	6/2/2010	0.36 ± 0.19	1.34 ± 0.72		14.60 ± 0.69	54.02 ± 2.54	Y		
	6/9/2010	0.82 ± 0.21	3.03 ± 0.79	Y	16.70 ± 0.75	61.79 ± 2.79	Y		
	6/16/2010	0.46 ± 0.19	1.69 ± 0.72		16.20 ± 0.71	59.94 ± 2.62	Y		
	6/23/2010	0.50 ± 0.19	1.86 ± 0.70		19.60 ± 0.77	72.52 ± 2.85	Y		
	6/30/2010	0.99 ± 0.21	3.66 ± 0.76	Y	28.60 ± 0.81	105.82 ± 3.01	Y		
IDAHO FALLS	4/7/2010	0.31 ± 0.22	1.14 ± 0.80		13.70 ± 0.79	50.69 ± 2.92	Y		
	4/14/2010	2.53 ± 0.33	9.36 ± 1.21	Y	24.60 ± 0.97	91.02 ± 3.59	Y		
	4/21/2010	1.87 ± 0.24	6.92 ± 0.89	Y	31.70 ± 0.84	117.29 ± 3.11	Y		
	4/28/2010	1.20 ± 0.23	4.44 ± 0.87	Y	21.60 ± 0.83	79.92 ± 3.07	Y		
	5/5/2010	0.53 ± 0.19	1.98 ± 0.70		9.39 ± 0.66	34.74 ± 2.45	Y		
	5/12/2010	1.13 ± 0.23	4.18 ± 0.87	Y	14.30 ± 0.73	52.91 ± 2.70	Y		
	5/19/2010	0.95 ± 0.23	3.50 ± 0.84	Y	26.00 ± 0.89	96.20 ± 3.30	Y		
	5/26/2010	1.16 ± 0.22	4.29 ± 0.82	Y	15.40 ± 0.72	56.98 ± 2.67	Y		
	6/2/2010	0.63 ± 0.30	2.33 ± 1.10		15.50 ± 0.95	57.35 ± 3.53	Y		
	6/9/2010	1.41 ± 0.25	5.22 ± 0.94	Y	20.10 ± 0.83	74.37 ± 3.05	Y		
	6/16/2010	0.34 ± 0.22	1.26 ± 0.82		17.30 ± 0.83	64.01 ± 3.06	Y		
	6/23/2010	1.54 ± 0.25	5.70 ± 0.91	Y	21.50 ± 0.82	79.55 ± 3.02	Y		
	6/30/2010	1.67 ± 0.27	6.18 ± 0.99	Y	33.60 ± 0.97	124.32 ± 3.60	Y		
JACKSON	4/7/2010	0.66 ± 0.22	2.42 ± 0.81		11.70 ± 0.72	43.29 ± 2.65	Y		
	4/14/2010	1.15 ± 0.25	4.26 ± 0.93	Y	21.50 ± 0.89	79.55 ± 3.29	Y		
	4/21/2010	1.57 ± 0.22	5.81 ± 0.82	Y	30.20 ± 0.81	111.74 ± 2.99	Y		
	4/28/2010	0.97 ± 0.23	3.59 ± 0.87	Y	17.30 ± 0.82	64.01 ± 3.04	Y		
	5/5/2010	0.54 ± 0.21	2.00 ± 0.76		10.00 ± 0.72	37.00 ± 2.68	Y		
	5/12/2010	0.84 ± 0.24	3.12 ± 0.88	Y	14.10 ± 0.78	52.17 ± 2.89	Y		

TABLE C-1. Weekly Gross Alpha and Gross Beta Concentrations in Air

Sampling Group and Location	Sampling Date	GROSS ALPHA					GROSS BETA						
		Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)		Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)		Result > 3s	Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)		Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)		Result > 3s		
REXBURG CMS	5/19/2010	1.29 ± 0.25	4.77 ± 0.93	Y	23.40 ± 0.89	86.58 ± 3.29	Y						
	5/26/2010	1.14 ± 0.25	4.22 ± 0.93	Y	14.30 ± 0.80	52.91 ± 2.97	Y						
	6/2/2010	0.52 ± 0.25	1.92 ± 0.91		14.00 ± 0.81	51.80 ± 3.00	Y						
	6/9/2010	0.79 ± 0.23	2.90 ± 0.85	Y	15.30 ± 0.80	56.61 ± 2.94	Y						
	6/16/2010	0.66 ± 0.24	2.46 ± 0.90		16.10 ± 0.82	59.57 ± 3.04	Y						
	6/23/2010	1.49 ± 0.28	5.51 ± 1.02	Y	22.10 ± 0.92	81.77 ± 3.41	Y						
	6/30/2010	1.45 ± 0.28	5.37 ± 1.02	Y	30.60 ± 1.00	113.22 ± 3.70	Y						
	4/7/2010	0.54 ± 0.17	1.99 ± 0.64	Y	13.00 ± 0.61	48.10 ± 2.25	Y						
	4/14/2010	1.31 ± 0.22	4.85 ± 0.81	Y	21.50 ± 0.75	79.55 ± 2.79	Y						
	4/21/2010	1.78 ± 0.21	6.59 ± 0.77	Y	31.40 ± 0.74	116.18 ± 2.73	Y						
4/28/2010	0.78 ± 0.19	2.87 ± 0.71	Y	21.50 ± 0.76	79.55 ± 2.81	Y							
5/5/2010	0.41 ± 0.16	1.50 ± 0.57		8.08 ± 0.56	29.90 ± 2.06	Y							
5/12/2010	0.78 ± 0.20	2.89 ± 0.73	Y	15.00 ± 0.67	55.50 ± 2.49	Y							
5/19/2010	1.71 ± 0.24	6.33 ± 0.90	Y	27.70 ± 0.85	102.49 ± 3.13	Y							
5/26/2010	0.31 ± 0.16	1.14 ± 0.60		13.10 ± 0.65	48.47 ± 2.41	Y							
6/2/2010	0.69 ± 0.21	2.55 ± 0.78	Y	15.20 ± 0.70	56.24 ± 2.59	Y							
6/9/2010	0.48 ± 0.21	1.78 ± 0.76		18.90 ± 0.82	69.93 ± 3.03	Y							
6/16/2010	0.57 ± 0.21	2.10 ± 0.77		16.00 ± 0.73	59.20 ± 2.69	Y							
6/23/2010	0.69 ± 0.20	2.56 ± 0.75	Y	21.10 ± 0.81	78.07 ± 2.99	Y							
6/30/2010	2.13 ± 0.31	7.88 ± 1.16	Y	32.80 ± 1.05	121.36 ± 3.89	Y							
<b>INL SITE</b>													
EFS	4/7/2010	0.31 ± 0.17	1.15 ± 0.64		11.90 ± 0.63	44.03 ± 2.33	Y						
	4/14/2010	1.19 ± 0.22	4.40 ± 0.83	Y	21.50 ± 0.79	79.55 ± 2.91	Y						
	4/21/2010	1.71 ± 0.22	6.33 ± 0.83	Y	30.20 ± 0.79	111.74 ± 2.93	Y						
	4/28/2010	0.97 ± 0.20	3.57 ± 0.73	Y	17.80 ± 0.71	65.86 ± 2.61	Y						
	5/5/2010	1.11 ± 0.21	4.11 ± 0.77	Y	9.80 ± 0.63	36.26 ± 2.32	Y						
	5/12/2010	0.43 ± 0.18	1.57 ± 0.65		15.00 ± 0.67	55.50 ± 2.48	Y						
	5/19/2010	0.72 ± 0.19	2.65 ± 0.70	Y	26.60 ± 0.81	98.42 ± 3.00	Y						
	5/26/2010	0.69 ± 0.21	2.54 ± 0.78	Y	13.50 ± 0.74	49.95 ± 2.74	Y						
	6/2/2010	0.67 ± 0.23	2.46 ± 0.85		14.10 ± 0.74	52.17 ± 2.73	Y						
	6/9/2010	0.63 ± 0.21	2.33 ± 0.76	Y	18.90 ± 0.79	69.93 ± 2.91	Y						
	6/16/2010	0.36 ± 0.20	1.33 ± 0.74		16.90 ± 0.75	62.53 ± 2.79	Y						
	6/23/2010	0.87 ± 0.21	3.23 ± 0.77	Y	19.80 ± 0.78	73.26 ± 2.87	Y						
	6/30/2010	1.43 ± 0.25	5.29 ± 0.91	Y	30.60 ± 0.91	113.22 ± 3.36	Y						
MAIN GATE	4/7/2010	0.83 ± 0.19	3.06 ± 0.68	Y	11.50 ± 0.58	42.55 ± 2.14	Y						
	4/14/2010	1.44 ± 0.22	5.33 ± 0.83	Y	20.10 ± 0.74	74.37 ± 2.72	Y						
	4/21/2010	1.25 ± 0.20	4.63 ± 0.72	Y	30.10 ± 0.76	111.37 ± 2.83	Y						
	4/28/2010	0.59 ± 0.17	2.18 ± 0.64	Y	20.10 ± 0.71	74.37 ± 2.62	Y						
	5/5/2010	0.36 ± 0.15	1.31 ± 0.56		8.93 ± 0.56	33.04 ± 2.08	Y						
	5/12/2010	0.59 ± 0.18	2.20 ± 0.67	Y	14.60 ± 0.65	54.02 ± 2.39	Y						
	5/19/2010	1.00 ± 0.21	3.70 ± 0.77	Y	26.10 ± 0.81	96.57 ± 3.01	Y						
	5/26/2010	0.27 ± 0.16	0.98 ± 0.61		12.10 ± 0.65	44.77 ± 2.42	Y						
	6/2/2010	0.19 ± 0.20	0.70 ± 0.74		14.40 ± 0.72	53.28 ± 2.68	Y						
	6/9/2010	0.64 ± 0.18	2.38 ± 0.68	Y	16.70 ± 0.69	61.79 ± 2.54	Y						
	6/16/2010	0.58 ± 0.19	2.13 ± 0.71	Y	17.20 ± 0.69	63.64 ± 2.57	Y						
	6/23/2010	0.60 ± 0.18	2.23 ± 0.65	Y	17.90 ± 0.70	66.23 ± 2.57	Y						
	6/30/2010	1.33 ± 0.24	4.92 ± 0.90	Y	30.10 ± 0.91	111.37 ± 3.37	Y						
VAN BUREN GATE	4/7/2010	0.48 ± 0.17	1.79 ± 0.62		10.80 ± 0.57	39.96 ± 2.11	Y						
	4/14/2010	1.44 ± 0.22	5.33 ± 0.80	Y	20.00 ± 0.71	74.00 ± 2.63	Y						
	4/21/2010	1.49 ± 0.21	5.51 ± 0.78	Y	31.00 ± 0.79	114.70 ± 2.92	Y						
	4/28/2010	1.16 ± 0.21	4.29 ± 0.78	Y	18.90 ± 0.73	69.93 ± 2.72	Y						
	5/5/2010	0.44 ± 0.16	1.61 ± 0.58		9.13 ± 0.57	33.78 ± 2.11	Y						
	5/12/2010	0.70 ± 0.27	2.58 ± 1.01		16.90 ± 0.95	62.53 ± 3.52	Y						
	5/19/2010	1.01 ± 0.24	3.74 ± 0.87	Y	25.60 ± 0.91	94.72 ± 3.36	Y						
	5/26/2010	0.53 ± 0.18	1.96 ± 0.65		11.80 ± 0.64	43.66 ± 2.37	Y						

TABLE C-1. Weekly Gross Alpha and Gross Beta Concentrations in Air

Sampling Group and Location	Sampling Date	GROSS ALPHA				GROSS BETA			
		Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)		Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)		Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)		Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)	
	6/2/2010	0.43 ± 0.20	1.60 ± 0.74		14.70 ± 0.70	54.39 ± 2.58	Y		
	6/9/2010	0.36 ± 0.21	1.34 ± 0.78		19.60 ± 0.87	72.52 ± 3.20	Y		
	6/16/2010	0.84 ± 0.21	3.09 ± 0.76	Y	18.10 ± 0.71	66.97 ± 2.63	Y		
	6/23/2010	0.70 ± 0.21	2.60 ± 0.79	Y	20.30 ± 0.83	75.11 ± 3.07	Y		
	6/30/2010	1.25 ± 0.26	4.63 ± 0.96	Y	29.40 ± 0.97	108.78 ± 3.57	Y		
QA-1 (VAN BUREN)	4/7/2010	0.54 ± 0.20	1.98 ± 0.74		11.40 ± 0.67	42.18 ± 2.48	Y		
	4/14/2010	1.28 ± 0.25	4.74 ± 0.94	Y	22.10 ± 0.88	81.77 ± 3.24	Y		
	4/21/2010	1.44 ± 0.23	5.33 ± 0.84	Y	28.40 ± 0.83	105.08 ± 3.08	Y		
	4/28/2010	1.06 ± 0.22	3.92 ± 0.83	Y	17.70 ± 0.78	65.49 ± 2.87	Y		
	5/5/2010	0.64 ± 0.21	2.36 ± 0.78	Y	7.96 ± 0.70	29.45 ± 2.59	Y		
	5/12/2010	0.84 ± 0.23	3.12 ± 0.84	Y	15.50 ± 0.76	57.35 ± 2.83	Y		
	5/19/2010	0.98 ± 0.23	3.64 ± 0.86	Y	27.00 ± 0.92	99.90 ± 3.39	Y		
	5/26/2010	0.44 ± 0.17	1.63 ± 0.61		12.00 ± 0.62	44.40 ± 2.30	Y		
	6/2/2010	0.29 ± 0.19	1.07 ± 0.71		14.90 ± 0.70	55.13 ± 2.58	Y		
	6/9/2010	0.92 ± 0.18	3.40 ± 0.65	Y	16.40 ± 0.61	60.68 ± 2.26	Y		
	6/16/2010	0.59 ± 0.21	2.17 ± 0.76		18.30 ± 0.75	67.71 ± 2.78	Y		
	6/23/2010	0.75 ± 0.18	2.79 ± 0.65	Y	19.40 ± 0.68	71.78 ± 2.50	Y		
	6/30/2010	1.91 ± 0.30	7.07 ± 1.10	Y	31.40 ± 1.02	116.18 ± 3.77	Y		
a. Invalid Sample Result									

TABLE C-2. Weekly Iodine-131 Activity in Air.

Sampling Group and Location	Sampling Date	Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)			Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)			Result > 3s
<b>BOUNDARY</b>								
ARCO	04/07/2010	-0.10	±	1.11	-0.38	±	4.12	
	04/14/2010	-1.72	±	1.18	-6.35	±	4.36	
	04/21/2010	-3.22	±	1.21	-11.92	±	4.49	
	04/28/2010	-1.31	±	1.15	-4.85	±	4.24	
	05/05/2010	-1.33	±	1.12	-4.91	±	4.16	
	05/12/2010	0.88	±	1.16	3.25	±	4.27	
	05/19/2010	4.27	±	1.24	15.78	±	4.60	Y
	05/26/2010	-0.26	±	1.17	-0.94	±	4.33	
	06/02/2010	-1.19	±	1.04	-4.42	±	3.84	
	06/09/2010	0.10	±	1.20	0.38	±	4.46	
	06/16/2010	-1.16	±	1.08	-4.28	±	3.99	
	06/23/2010	0.28	±	1.19	1.03	±	4.40	
	06/30/2010	1.36	±	1.28	5.03	±	4.74	
ATOMIC CITY	04/07/2010	-0.11	±	1.22	-0.42	±	4.52	
	04/14/2010	-1.69	±	1.16	-6.24	±	4.28	
	04/21/2010	-3.59	±	1.35	-13.29	±	5.00	
	04/28/2010	-1.28	±	1.12	-4.72	±	4.13	
	05/05/2010	-1.38	±	1.17	-5.09	±	4.32	
	05/12/2010	0.79	±	1.04	2.92	±	3.85	
	05/19/2010	5.00	±	1.46	18.51	±	5.39	Y
	05/26/2010	-0.27	±	1.26	-1.02	±	4.65	
	06/02/2010	-1.43	±	1.24	-5.28	±	4.59	
	06/09/2010	0.10	±	1.16	0.36	±	4.31	
	06/16/2010	-1.32	±	1.23	-4.87	±	4.54	
	06/23/2010	0.23	±	0.97	0.84	±	3.59	
	06/30/2010	1.55	±	1.46	5.75	±	5.41	
BLUE DOME	04/07/2010	-0.67	±	0.97	-2.49	±	3.59	
	04/14/2010	-0.07	±	1.03	-0.26	±	3.80	
	04/21/2010	1.18	±	1.05	4.35	±	3.90	
	04/28/2010	-0.03	±	1.00	-0.13	±	3.71	
	05/05/2010	-0.20	±	1.06	-0.74	±	3.90	
	05/12/2010	-1.44	±	0.95	-5.34	±	3.51	
	05/19/2010	0.33	±	1.02	1.23	±	3.78	
	05/26/2010	0.12	±	1.03	0.44	±	3.80	
	06/02/2010	0.11	±	0.93	0.40	±	3.43	
	06/09/2010	-0.84	±	1.10	-3.10	±	4.06	
	06/16/2010	0.02	±	0.91	0.08	±	3.37	
	06/23/2010	-0.42	±	0.94	-1.56	±	3.47	
	06/30/2010	1.40	±	1.00	5.18	±	3.71	
FAA TOWER	04/07/2010	-0.78	±	1.12	-2.88	±	4.16	
	04/14/2010	-0.08	±	1.14	-0.28	±	4.21	
	04/21/2010	1.26	±	1.13	4.66	±	4.18	
	04/28/2010	-0.05	±	1.58	-0.20	±	5.84	
	05/05/2010	-0.22	±	1.16	-0.81	±	4.28	
	05/12/2010	-1.66	±	1.09	-6.14	±	4.04	
	05/19/2010	0.38	±	1.15	1.39	±	4.26	
	05/26/2010	0.16	±	1.39	0.59	±	5.13	
	06/02/2010	0.11	±	0.98	0.42	±	3.61	
	06/09/2010	-0.75	±	0.98	-2.76	±	3.62	
	06/16/2010	0.02	±	0.97	0.08	±	3.57	



TABLE C-2. Weekly Iodine-131 Activity in Air.

Sampling Group and Location	Sampling Date	Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)			Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)			Result > 3s
<b>BOUNDARY</b>								
	06/23/2010	-0.43	±	0.97	-1.60	±	3.57	
	06/30/2010	1.58	±	1.13	5.85	±	4.19	
HOWE	04/07/2010	-0.64	±	0.92	-2.37	±	3.42	
	04/14/2010	-0.06	±	0.90	-0.23	±	3.34	
	04/21/2010	1.13	±	1.01	4.18	±	3.75	
	04/28/2010	-0.03	±	0.94	-0.12	±	3.46	
	05/05/2010	-0.19	±	0.99	-0.70	±	3.68	
	05/12/2010	-1.38	±	0.91	-5.11	±	3.36	
	05/19/2010	0.31	±	0.94	1.14	±	3.49	
	05/26/2010	0.11	±	0.92	0.39	±	3.40	
	06/02/2010	0.10	±	0.83	0.36	±	3.09	
	06/09/2010	-0.63	±	0.82	-2.33	±	3.05	
	06/16/2010	0.02	±	0.93	0.08	±	3.44	
	06/23/2010	-0.36	±	0.80	-1.33	±	2.97	
	06/30/2010	1.37	±	0.98	5.05	±	3.62	
MONTEVIEW	04/07/2010	-0.68	±	0.99	-2.53	±	3.65	
	04/14/2010	-0.07	±	1.03	-0.26	±	3.82	
	04/21/2010	1.24	±	1.11	4.58	±	4.11	
	04/28/2010	-0.03	±	1.02	-0.13	±	3.76	
	05/05/2010	-0.20	±	1.06	-0.74	±	3.92	
	05/12/2010	-1.44	±	0.95	-5.33	±	3.51	
	05/19/2010	0.34	±	1.05	1.27	±	3.88	
	05/26/2010	0.12	±	1.01	0.43	±	3.75	
	06/02/2010	0.12	±	1.04	0.45	±	3.84	
	06/09/2010	-0.73	±	0.96	-2.70	±	3.54	
	06/16/2010	0.03	±	1.22	0.10	±	4.51	
	06/23/2010	-0.46	±	1.02	-1.69	±	3.78	
	06/30/2010	1.45	±	1.04	5.35	±	3.83	
MUD LAKE	04/07/2010	-0.70	±	1.01	-2.60	±	3.74	
	04/14/2010	-0.07	±	1.01	-0.25	±	3.74	
	04/21/2010	1.33	±	1.20	4.93	±	4.42	
	04/28/2010	-0.03	±	0.92	-0.12	±	3.40	
	05/05/2010	-0.21	±	1.09	-0.77	±	4.04	
	05/12/2010	-1.45	±	0.95	-5.35	±	3.52	
	05/19/2010	0.35	±	1.08	1.31	±	4.01	
	05/26/2010	0.11	±	0.95	0.41	±	3.53	
	06/02/2010	0.11	±	0.95	0.41	±	3.52	
	06/09/2010	-0.66	±	0.87	-2.44	±	3.20	
	06/16/2010	0.02	±	1.09	0.09	±	4.02	
	06/23/2010	-0.37	±	0.83	-1.37	±	3.06	
	06/30/2010	1.74	±	1.25	6.45	±	4.61	
<b>DISTANT</b>								
BLACKFOOT CMS	04/07/2010	-0.10	±	1.12	-0.38	±	4.13	
	04/14/2010	-1.59	±	1.09	-5.87	±	4.02	
	04/21/2010	-2.85	±	1.07	-10.53	±	3.96	
	04/28/2010	-1.25	±	1.09	-4.63	±	4.04	
	05/05/2010	-1.32	±	1.12	-4.89	±	4.14	
a	05/12/2010	4.33	±	5.70	16.03	±	21.08	
	05/19/2010	4.03	±	1.17	14.90	±	4.34	Y
	05/26/2010	-0.25	±	1.14	-0.93	±	4.24	

TABLE C-2. Weekly Iodine-131 Activity in Air.

Sampling Group and Location	Sampling Date	Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)			Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)			Result > 3s
<b>BOUNDARY</b>								
	06/02/2010	-1.17	±	1.02	-4.33	±	3.76	
	06/09/2010	0.13	±	1.59	0.50	±	5.89	
	06/16/2010	-1.20	±	1.12	-4.44	±	4.14	
	06/23/2010	0.40	±	1.69	1.47	±	6.25	
<b>a</b>	<b>06/30/2010</b>	<b>3.74</b>	<b>±</b>	<b>3.52</b>	<b>13.83</b>	<b>±</b>	<b>13.03</b>	
<b>CRATERS</b>								
	04/07/2010	-0.10	±	1.11	-0.38	±	4.10	
	04/14/2010	-1.66	±	1.14	-6.14	±	4.21	
	04/21/2010	-3.05	±	1.15	-11.27	±	4.24	
	04/28/2010	-1.32	±	1.15	-4.87	±	4.26	
	05/05/2010	-1.31	±	1.11	-4.85	±	4.11	
	05/12/2010	0.86	±	1.13	3.18	±	4.18	
	05/19/2010	4.17	±	1.22	15.44	±	4.50	Y
	05/26/2010	-0.27	±	1.21	-0.98	±	4.49	
	06/02/2010	-1.17	±	1.02	-4.34	±	3.77	
	06/09/2010	0.10	±	1.19	0.37	±	4.40	
	06/16/2010	-1.11	±	1.03	-4.10	±	3.82	
	06/23/2010	0.25	±	1.09	0.94	±	4.02	
	06/30/2010	1.26	±	1.19	4.67	±	4.40	
<b>DUBOIS</b>								
	04/07/2010	-0.77	±	1.11	-2.84	±	4.10	
	04/14/2010	-0.07	±	1.01	-0.25	±	3.75	
	04/21/2010	1.26	±	1.13	4.67	±	4.19	
	04/28/2010	-0.04	±	1.17	-0.15	±	4.35	
	05/05/2010	-0.23	±	1.21	-0.85	±	4.49	
	05/12/2010	-1.45	±	0.95	-5.35	±	3.52	
	05/19/2010	0.36	±	1.12	1.35	±	4.13	
	05/26/2010	0.12	±	1.04	0.44	±	3.86	
	06/02/2010	0.13	±	1.12	0.49	±	4.15	
	06/09/2010	-0.81	±	1.06	-3.00	±	3.94	
	06/16/2010	0.02	±	1.06	0.09	±	3.93	
	06/23/2010	-0.43	±	0.95	-1.58	±	3.52	
	06/30/2010	1.49	±	1.07	5.53	±	3.96	
<b>QA-2</b>								
	04/07/2010	-0.69	±	1.00	-2.57	±	3.70	
	04/14/2010	-0.07	±	1.03	-0.26	±	3.83	
	04/21/2010	1.19	±	1.07	4.42	±	3.97	
	04/28/2010	-0.04	±	1.17	-0.15	±	4.33	
	05/05/2010	-0.20	±	1.04	-0.73	±	3.86	
	05/12/2010	-1.50	±	0.99	-5.54	±	3.65	
	05/19/2010	0.31	±	0.97	1.16	±	3.57	
	05/26/2010	0.15	±	1.29	0.55	±	4.78	
	06/02/2010	0.11	±	0.98	0.42	±	3.61	
	06/09/2010	-0.85	±	1.11	-3.14	±	4.12	
	06/16/2010	0.02	±	1.03	0.09	±	3.80	
	06/23/2010	-0.47	±	1.05	-1.75	±	3.90	
	06/30/2010	1.33	±	0.95	4.91	±	3.51	
<b>IDAHO FALLS</b>								
	04/07/2010	-0.98	±	1.41	-3.62	±	5.22	
	04/14/2010	-0.10	±	1.42	-0.36	±	5.26	
	04/21/2010	1.56	±	1.40	5.76	±	5.17	
	04/28/2010	-0.04	±	1.23	-0.16	±	4.57	
	05/05/2010	-0.24	±	1.26	-0.88	±	4.65	
	05/12/2010	-1.76	±	1.16	-6.51	±	4.29	

TABLE C-2. Weekly Iodine-131 Activity in Air.

Sampling Group and Location	Sampling Date	Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)			Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)			Result > 3s
<b>BOUNDARY</b>								
	05/19/2010	0.41	±	1.26	1.51	±	4.64	
	05/26/2010	0.14	±	1.18	0.50	±	4.38	
	06/02/2010	0.17	±	1.49	0.64	±	5.50	
	06/09/2010	-0.89	±	1.17	-3.30	±	4.33	
	06/16/2010	0.03	±	1.24	0.10	±	4.59	
	06/23/2010	-0.49	±	1.10	-1.82	±	4.06	
	06/30/2010	1.61	±	1.15	5.94	±	4.25	
<b>JACKSON</b>								
	04/07/2010	-0.13	±	1.41	-0.48	±	5.21	
	04/14/2010	-2.14	±	1.47	-7.91	±	5.42	
	04/21/2010	-3.72	±	1.40	-13.76	±	5.18	
	04/28/2010	-1.63	±	1.43	-6.05	±	5.29	
	05/05/2010	-1.70	±	1.44	-6.29	±	5.33	
	05/12/2010	1.10	±	1.45	4.08	±	5.36	
	05/19/2010	5.16	±	1.50	19.11	±	5.56	Y
	05/26/2010	-0.34	±	1.53	-1.24	±	5.68	
	06/02/2010	-1.53	±	1.33	-5.66	±	4.92	
	06/09/2010	0.13	±	1.50	0.47	±	5.53	
	06/16/2010	-1.46	±	1.37	-5.42	±	5.05	
	06/23/2010	0.34	±	1.44	1.25	±	5.35	
	06/30/2010	1.56	±	1.47	5.78	±	5.44	
<b>REXBURG CMS</b>								
	04/07/2010	-0.70	±	1.01	-2.58	±	3.73	
	04/14/2010	-0.07	±	1.04	-0.26	±	3.86	
	04/21/2010	1.26	±	1.13	4.65	±	4.17	
	04/28/2010	-0.04	±	1.09	-0.14	±	4.02	
	05/05/2010	-0.20	±	1.05	-0.74	±	3.88	
	05/12/2010	-1.55	±	1.02	-5.74	±	3.78	
	05/19/2010	0.36	±	1.11	1.34	±	4.12	
	05/26/2010	0.13	±	1.09	0.47	±	4.04	
	06/02/2010	0.12	±	0.99	0.43	±	3.66	
	06/09/2010	-0.91	±	1.19	-3.36	±	4.40	
	06/16/2010	0.02	±	1.07	0.09	±	3.96	
	06/23/2010	-0.49	±	1.09	-1.81	±	4.04	
	06/30/2010	1.83	±	1.31	6.79	±	4.86	
<b>INL SITE</b>								
<b>EFS</b>								
	04/07/2010	-0.11	±	1.18	-0.41	±	4.37	
	04/14/2010	-1.78	±	1.22	-6.59	±	4.52	
	04/21/2010	-3.59	±	1.35	-13.28	±	5.00	
	04/28/2010	-1.30	±	1.14	-4.81	±	4.21	
	05/05/2010	-1.42	±	1.20	-5.25	±	4.45	
	05/12/2010	0.88	±	1.15	3.24	±	4.26	
	05/19/2010	4.21	±	1.22	15.56	±	4.53	Y
	05/26/2010	-0.31	±	1.41	-1.14	±	5.20	
	06/02/2010	-1.35	±	1.17	-4.98	±	4.33	
	06/09/2010	0.11	±	1.36	0.42	±	5.02	
	06/16/2010	-1.28	±	1.19	-4.74	±	4.42	
	06/23/2010	0.28	±	1.19	1.03	±	4.39	
	06/30/2010	1.34	±	1.26	4.96	±	4.67	
<b>MAIN GATE</b>								
	04/07/2010	-0.10	±	1.07	-0.37	±	3.94	
	04/14/2010	-1.67	±	1.14	-6.16	±	4.23	
	04/21/2010	-3.40	±	1.28	-12.56	±	4.73	

TABLE C-2. Weekly Iodine-131 Activity in Air.

Sampling Group and Location	Sampling Date	Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)			Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)			Result > 3s
<b>BOUNDARY</b>								
	04/28/2010	-1.24	±	1.08	-4.58	±	4.01	
	05/05/2010	-1.27	±	1.08	-4.70	±	3.98	
	05/12/2010	0.84	±	1.10	3.10	±	4.08	
	05/19/2010	4.27	±	1.24	15.80	±	4.60	Y
	05/26/2010	-0.27	±	1.24	-1.00	±	4.57	
	06/02/2010	-1.30	±	1.13	-4.83	±	4.20	
	06/09/2010	0.10	±	1.17	0.37	±	4.35	
	06/16/2010	-1.13	±	1.05	-4.17	±	3.89	
	06/23/2010	0.25	±	1.06	0.92	±	3.91	
	06/30/2010	1.36	±	1.28	5.03	±	4.74	
<b>VAN BUREN GATE</b>								
	04/07/2010	-0.10	±	1.07	-0.37	±	3.95	
	04/14/2010	-1.59	±	1.09	-5.88	±	4.03	
	04/21/2010	-3.52	±	1.33	-13.03	±	4.90	
	04/28/2010	-1.34	±	1.17	-4.96	±	4.34	
	05/05/2010	-1.29	±	1.09	-4.77	±	4.04	
	05/12/2010	1.35	±	1.77	4.98	±	6.55	
	05/19/2010	5.10	±	1.48	18.87	±	5.49	Y
	05/26/2010	-0.26	±	1.21	-0.98	±	4.49	
	06/02/2010	-1.23	±	1.07	-4.55	±	3.95	
	06/09/2010	0.13	±	1.53	0.48	±	5.67	
	06/16/2010	-1.14	±	1.06	-4.21	±	3.92	
	06/23/2010	0.30	±	1.29	1.12	±	4.78	
	06/30/2010	1.50	±	1.41	5.56	±	5.23	
<b>QA-1</b>								
	04/07/2010	-0.12	±	1.30	-0.45	±	4.83	
	04/14/2010	-2.07	±	1.42	-7.65	±	5.24	
	04/21/2010	-4.07	±	1.53	-15.05	±	5.67	
	04/28/2010	-1.50	±	1.31	-5.55	±	4.85	
	05/05/2010	-1.72	±	1.46	-6.38	±	5.40	
	05/12/2010	1.03	±	1.36	3.82	±	5.03	
	05/19/2010	5.03	±	1.46	18.59	±	5.41	Y
	05/26/2010	-0.25	±	1.16	-0.94	±	4.29	
	06/02/2010	-1.23	±	1.07	-4.54	±	3.94	
	06/09/2010	0.08	±	1.00	0.31	±	3.69	
	06/16/2010	-1.23	±	1.14	-4.54	±	4.23	
	06/23/2010	0.23	±	0.98	0.85	±	3.63	
	06/30/2010	1.57	±	1.48	5.81	±	5.47	
<b>a. Invalid Sample Result</b>								

TABLE C-3. Quarterly Americium-241, Cesium-137, Plutonium-238, Plutonium-239/240, and Strontium-90 Concentrations in Composite Air Filters.

Sampling Group and Location	Sampling Date	Analyte	Result ± 1s Uncertainty (x 10 <sup>-18</sup> µCi/mL)			Result ± 1s Uncertainty (x 10 <sup>-13</sup> Bq/mL)			Result > 3s
<b>BOUNDARY</b>									
ARCO	6/30/2010	CESIUM-137	-82.40	±	141.00	-304.88	±	521.70	
ATOMIC CITY	6/30/2010	CESIUM-137	-100.00	±	125.00	-370.00	±	462.50	
		STRONTIUM-90	13.85	±	27.70	51.25	±	102.49	
BLUE DOME	6/30/2010	CESIUM-137	29.70	±	130.00	109.89	±	481.00	
FAA TOWER	6/30/2010	AMERICIUM-241	-0.49	±	0.85	-1.82	±	3.16	
		CESIUM-137	83.30	±	140.00	308.21	±	518.00	
		PLUTONIUM-238	0.00	±	0.13	0.00	±	0.47	
		PLUTONIUM-239/40	0.00	±	0.00	0.00	±	0.00	
HOWE	6/30/2010	CESIUM-137	-11.20	±	122.00	-41.44	±	451.40	
MONTEVIEW	6/30/2010	CESIUM-137	-146.00	±	137.00	-540.20	±	506.90	
		STRONTIUM-90	11.00	±	22.00	40.70	±	81.40	
		AMERICIUM-241	1.86	±	1.64	6.86	±	6.08	
MUD LAKE	6/30/2010	CESIUM-137	46.50	±	90.40	172.05	±	334.48	
		PLUTONIUM-238	0.31	±	0.44	1.16	±	1.64	
		PLUTONIUM-239/40	0.47	±	0.27	1.74	±	1.01	

TABLE C-3. Quarterly Americium-241, Cesium-137, Plutonium-238, Plutonium-239/240, and Strontium-90 Concentrations in Composite Air Filters.

Sampling Group and Location	Sampling Date	Analyte	Result ± 1s Uncertainty (x 10 <sup>-18</sup> µCi/mL)			Result ± 1s Uncertainty (x 10 <sup>-13</sup> Bq/mL)			Result > 3s
<b>DISTANT</b>									
BLACKFOOT	6/30/2010	CESIUM-137	25.10	±	134.00	92.87	±	495.80	
		STRONTIUM-90	13.35	±	26.70	49.40	±	98.79	
CRATERS	6/30/2010	CESIUM-137	-90.40	±	89.10	-334.48	±	329.67	
DUBOIS	6/30/2010	AMERICIUM-241	2.39	±	1.70	8.84	±	6.30	
		CESIUM-137	-23.00	±	256.00	-85.10	±	947.20	
		PLUTONIUM-238	0.72	±	0.64	2.68	±	2.37	
		PLUTONIUM-239/40	0.48	±	0.34	1.78	±	1.27	
DUBOIS (QA-2)	6/30/2010	AMERICIUM-241	0.00	±	0.00	0.00	±	0.00	
		CESIUM-137	52.70	±	95.60	194.99	±	353.72	
		PLUTONIUM-238	0.00	±	0.00	0.00	±	0.00	
		PLUTONIUM-239/40	0.00	±	0.00	0.00	±	0.00	
IDAHO FALLS	6/30/2010	AMERICIUM-241	1.35	±	1.35	4.99	±	5.01	
		CESIUM-137	-42.40	±	140.00	-156.88	±	518.00	
		PLUTONIUM-238	0.18	±	0.31	0.65	±	1.13	
		PLUTONIUM-239/40	0.71	±	0.36	2.61	±	1.32	
JACKSON	6/30/2010	CESIUM-137	39.50	±	106.00	146.15	±	392.20	
REXBURG CMS	6/30/2010	CESIUM-137	-62.90	±	146.00	-232.73	±	540.20	
		STRONTIUM-90	12.45	±	24.90	46.07	±	92.13	
<b>INL SITE</b>									
EFS	6/30/2010	AMERICIUM-241	0.92	±	1.60	3.41	±	5.90	
		CESIUM-137	-82.30	±	85.70	-304.51	±	317.09	
		PLUTONIUM-238	-0.77	±	0.56	-2.86	±	2.06	
		PLUTONIUM-239/40	-0.15	±	0.41	-0.57	±	1.51	
MAIN GATE	6/30/2010	AMERICIUM-241	-0.58	±	1.01	-2.16	±	3.74	
		CESIUM-137	89.70	±	90.20	331.89	±	333.74	
		PLUTONIUM-238	0.74	±	0.39	2.73	±	1.45	
		PLUTONIUM-239/40	0.49	±	0.35	1.82	±	1.29	
VAN BUREN GATE	6/30/2010	CESIUM-137	-48.80	±	88.00	-180.56	±	325.60	
		STRONTIUM-90	16.50	±	33.00	61.05	±	122.10	
VAN BUREN GATE (QA-1)	6/30/2010	CESIUM-137	34.10	±	133.00	0.00	±	0.00	
		STRONTIUM-90	14.85	±	29.70	54.95	±	109.89	

**TABLE C-4. Tritium Concentrations in Atmospheric Moisture.**

Sampling Group and Location	Start Date	Sampling Date	Result ± 1s Uncertainty			Result ± 1s Uncertainty			Collection Medium	Result > 3s
			(x 10 <sup>-13</sup> μCi/mL <sub>air</sub> )			(x 10 <sup>-9</sup> Bq/mL <sub>air</sub> )				
<b>BOUNDARY</b>										
ATOMIC CITY	03/24/2010	04/28/2010	3.06	±	1.58	11.32	±	5.84	Molecular Sieve	
ATOMIC CITY	04/28/2010	05/26/2010	6.50	±	1.91	24.06	±	7.07	Molecular Sieve	Y
ATOMIC CITY	05/26/2010	06/14/2010	8.97	±	3.07	33.20	±	11.35	Molecular Sieve	
ATOMIC CITY	06/14/2010	06/30/2010	10.09	±	2.86	37.32	±	10.60	Molecular Sieve	Y
<b>DISTANT</b>										
BLACKFOOT	03/31/2010	04/28/2010	2.76	±	1.43	10.20	±	5.30	Molecular Sieve	
BLACKFOOT	04/28/2010	05/19/2010	5.05	±	1.74	18.68	±	6.43	Molecular Sieve	
BLACKFOOT	05/19/2010	06/09/2010	10.53	±	2.33	38.97	±	8.63	Molecular Sieve	Y
BLACKFOOT	06/09/2010	06/30/2010	3.25	±	2.18	12.02	±	8.05	Molecular Sieve	
IDAHO FALLS	03/21/2010	04/21/2010	1.37	±	1.26	5.08	±	4.65	Molecular Sieve	
IDAHO FALLS	04/21/2010	05/12/2010	3.29	±	1.48	12.19	±	5.48	Molecular Sieve	
IDAHO FALLS	05/12/2010	06/02/2010	7.54	±	1.89	27.88	±	6.99	Molecular Sieve	Y
IDAHO FALLS	06/02/2010	06/14/2010	14.39	±	2.97	53.25	±	10.97	Molecular Sieve	Y
IDAHO FALLS	06/14/2010	06/30/2010	7.31	±	2.34	27.06	±	8.67	Molecular Sieve	Y
REXBURG	03/24/2010	04/21/2010	2.77	±	1.59	10.23	±	5.88	Molecular Sieve	
REXBURG	04/21/2010	05/19/2010	8.15	±	1.88	30.16	±	6.96	Molecular Sieve	Y
REXBURG	05/19/2010	06/09/2010	15.19	±	2.58	56.19	±	9.54	Molecular Sieve	Y
REXBURG	06/09/2010	06/30/2010	10.84	±	2.71	40.11	±	10.04	Molecular Sieve	Y

**TABLE C-5. Monthly and Weekly Tritium Concentrations in Precipitation.**

Location	Start Date	End Date	Result $\pm$ 1s Uncertainty			Result $\pm$ 1s Uncertainty			Result > 3s
			(pCi/L)			(Bq/L)			
IDAHO FALLS	3/2/2010	4/1/2010	-14.80	$\pm$	31.80	-0.55	$\pm$	1.18	
	4/1/2010	5/3/2010	76.50	$\pm$	34.20	2.83	$\pm$	1.27	
	5/3/2010	6/1/2010	107.00	$\pm$	32.80	3.96	$\pm$	1.21	Y
CFA	3/1/2010	4/1/2010	9.00	$\pm$	31.80	0.33	$\pm$	1.18	
	4/1/2010	5/3/2010	69.70	$\pm$	33.20	2.58	$\pm$	1.23	
	5/3/2010	6/1/2010	21.50	$\pm$	31.60	0.80	$\pm$	1.17	
EFS	3/31/2010	4/7/2010	173.00	$\pm$	36.10	6.40	$\pm$	1.34	Y
	4/14/2010	4/21/2010	18.60	$\pm$	33.00	0.69	$\pm$	1.22	
	4/21/2010	4/28/2010	68.30	$\pm$	33.50	2.53	$\pm$	1.24	
	5/5/2010	5/12/2010	45.70	$\pm$	33.00	1.69	$\pm$	1.22	
	5/12/2010	5/19/2010	103.00	$\pm$	32.80	3.81	$\pm$	1.21	Y
	5/19/2010	5/26/2010	251.00	$\pm$	35.70	9.29	$\pm$	1.32	Y
	5/26/2010	6/2/2010	157.00	$\pm$	33.90	5.81	$\pm$	1.25	Y



Table C-6. Gross Alpha, Gross Beta and Tritium Concentrations in Surface Water

Location	Analyte	Sampling Date	Result ± 1s Uncertainty			Result ± 1s Uncertainty			Result > 3s
			(pCi/L)			(Bq/L)			
<b>DRINKING WATER</b>									
Big Lost River at Rest Area	GROSS ALPHA	6/14/10	1.11	±	0.27	0.04	±	0.01	Y
	GROSS BETA		1.68	±	0.51	0.06	±	0.02	Y
	TRITIUM		114.00	±	32.20	4.22	±	1.19	Y
Big Lost River near INTEC	GROSS ALPHA	6/14/10	1.46	±	0.28	0.05	±	0.01	Y
	GROSS BETA		2.18	±	0.51	0.08	±	0.02	Y
	TRITIUM		40.80	±	31.90	1.51	±	1.18	Y
Big Lost River at EFS	GROSS ALPHA	6/14/10	1.39	±	0.29	0.05	±	0.01	Y
	GROSS BETA		2.07	±	0.51	0.08	±	0.02	Y
	TRITIUM		163.00	±	33.20	6.04	±	1.23	Y
Big Lost River at EFS (Duplicate)	GROSS ALPHA	6/14/10	1.62	±	0.29	0.06	±	0.01	Y
	GROSS BETA		2.42	±	0.52	0.09	±	0.02	Y
	TRITIUM		147.00	±	32.90	5.44	±	1.22	Y
Big Lost River north of NRF	GROSS ALPHA	6/14/10	1.71	±	0.31	0.06	±	0.01	Y
	GROSS BETA		0.90	±	0.50	0.03	±	0.02	Y
	TRITIUM		115.00	±	32.20	4.26	±	1.19	Y

Table C-7. Weekly and Monthly Iodine-131 and Cesium-137 Concentrations in Milk

Location	Sampling Date	Iodine-131				Cesium-137			
		Result ± 1s Uncertainty (pCi/L)		Result ± 1s Uncertainty (Bq/L)		Result > 3s	Result ± 1s Uncertainty (pCi/L)		Result ± 1s Uncertainty (Bq/L)
BLACKFOOT									
	04/06/10	1.43 ± 2.75	0.053 ± 0.102		2.70 ± 1.25	0.100 ± 0.046			
	05/03/10	0.81 ± 1.87	0.030 ± 0.069		1.34 ± 1.39	0.050 ± 0.051			
	06/01/10	0.45 ± 2.73	0.017 ± 0.101		-1.13 ± 1.38	-0.042 ± 0.051			
CONTROL									
	04/06/10	-2.86 ± 2.22	-0.106 ± 0.082		1.95 ± 1.46	0.072 ± 0.054			
	05/04/10	3.83 ± 1.94	0.142 ± 0.072		-1.11 ± 1.39	-0.041 ± 0.051			
	06/01/10	-0.63 ± 2.39	-0.023 ± 0.089		2.28 ± 1.44	0.084 ± 0.053			
DIETRICH									
	04/06/10	-2.25 ± 1.72	-0.083 ± 0.064		-0.62 ± 1.45	-0.023 ± 0.054			
	05/04/10	-1.10 ± 0.86	-0.041 ± 0.032		-0.08 ± 0.76	-0.003 ± 0.028			
	06/01/10	0.58 ± 1.73	0.022 ± 0.064		0.38 ± 1.41	0.014 ± 0.052			
Duplicate	06/01/10	-4.04 ± 1.94	-0.150 ± 0.072		0.25 ± 1.39	0.009 ± 0.051			
FORT HALL									
	04/06/10	-0.74 ± 0.90	-0.027 ± 0.033		1.47 ± 0.78	0.054 ± 0.029			
	05/05/10	-0.39 ± 1.71	-0.014 ± 0.063		3.15 ± 1.39	0.117 ± 0.051			
	06/01/10	1.14 ± 1.15	0.042 ± 0.043		0.47 ± 0.91	0.017 ± 0.034			
HOWE									
	04/06/10	-0.64 ± 1.71	-0.024 ± 0.063		2.15 ± 1.32	0.080 ± 0.049			
	05/04/10	-2.19 ± 1.84	-0.081 ± 0.068		0.84 ± 1.35	0.031 ± 0.050			
	06/01/10	0.49 ± 1.89	0.018 ± 0.070		1.09 ± 1.35	0.040 ± 0.050			
IDAHO FALLS									
	04/06/10	1.22 ± 1.54	0.045 ± 0.057		2.20 ± 1.26	0.081 ± 0.047			
	04/13/10	1.24 ± 0.83	0.046 ± 0.031		-0.11 ± 0.76	-0.004 ± 0.028			
	04/20/10	-0.47 ± 0.83	-0.017 ± 0.031		-0.13 ± 0.75	-0.005 ± 0.028			
	04/27/10	-0.92 ± 0.84	-0.034 ± 0.031		0.60 ± 0.78	0.022 ± 0.029			
	05/04/10	-0.67 ± 0.82	-0.025 ± 0.030		-0.06 ± 0.77	-0.002 ± 0.029			
	05/11/10	1.19 ± 0.84	0.044 ± 0.031		0.91 ± 0.75	0.034 ± 0.028			
	05/18/10	-1.00 ± 0.84	-0.037 ± 0.031		-1.33 ± 0.75	-0.049 ± 0.028			
	05/25/10	0.01 ± 0.81	0.000 ± 0.030		1.48 ± 0.76	0.055 ± 0.028			
	06/01/10	0.45 ± 1.53	0.017 ± 0.057		0.60 ± 1.33	0.022 ± 0.049			
	06/08/10	0.36 ± 0.92	0.013 ± 0.034		-0.36 ± 0.78	-0.013 ± 0.029			
	06/15/10	0.73 ± 0.87	0.027 ± 0.032		1.30 ± 0.77	0.048 ± 0.029			
	06/22/10	1.33 ± 0.91	0.049 ± 0.034		0.92 ± 0.75	0.034 ± 0.028			
	06/29/10	-1.32 ± 1.07	-0.049 ± 0.040		-1.04 ± 0.85	-0.039 ± 0.032			
RUPERT									
	04/06/10	-1.50 ± 0.83	-0.056 ± 0.031		0.25 ± 0.73	0.009 ± 0.027			
	05/04/10	-0.09 ± 1.62	-0.003 ± 0.060		-0.36 ± 1.38	-0.013 ± 0.051			
	06/01/10	1.69 ± 1.88	0.063 ± 0.070		-1.06 ± 1.42	-0.039 ± 0.053			
TERRETON									
	04/06/10	-3.43 ± 1.93	-0.127 ± 0.071		0.89 ± 1.42	0.033 ± 0.053			
Duplicate	04/06/10	-0.76 ± 1.77	-0.028 ± 0.066		0.14 ± 1.37	0.005 ± 0.051			
	05/04/10	1.52 ± 2.09	0.056 ± 0.077		-1.17 ± 1.46	-0.043 ± 0.054			
	06/01/10	1.03 ± 2.24	0.038 ± 0.083		0.51 ± 1.42	0.019 ± 0.053			

Table C-8. Strontium-90 and Tritium Concentrations in Milk

Strontium-90								
Location	Sampling Date	Result ± 1s Uncertainty			Result ± 1s Uncertainty			Result > 3s
		(pCi/L)			(Bq/L)			
BLACKFOOT	05/03/10	1.01	±	0.15	0.037	±	0.006	Y
CONTROL	05/04/10	1.37	±	0.16	0.051	±	0.006	Y
DIETRICH	05/04/10	0.36	±	0.11	0.013	±	0.004	Y
FORT HALL	05/05/10	0.45	±	0.13	0.017	±	0.005	Y
HOWE	05/04/10	0.69	±	0.16	0.025	±	0.006	Y
IDAHO FALLS	05/04/10	1.74	±	0.18	0.064	±	0.007	Y
RUPERT	05/04/10	1.84	±	0.15	0.068	±	0.006	Y
TERRETON	05/04/10	0.94	±	0.17	0.035	±	0.006	Y
Tritium								
Location	Sampling Date	Concentration ± 1s			Concentration ± 1s			Result > 3s
		(pCi/L)			(Bq/L)			
BLACKFOOT	05/03/10	47.35	±	32.92	1.754	±	1.219	
CONTROL	05/04/10	82.56	±	33.04	3.058	±	1.224	
DIETRICH	05/04/10	-2.08	±	32.97	-0.077	±	1.221	
FORT HALL	05/05/10	33.45	±	32.31	1.239	±	1.197	
HOWE	05/04/10	71.44	±	33.43	2.646	±	1.238	
IDAHO FALLS	05/04/10	29.30	±	33.64	1.085	±	1.246	
RUPERT	05/04/10	50.79	±	32.68	1.881	±	1.210	
TERRETON	05/04/10	-183.83	±	132.89	-6.809	±	4.922	

Table C-9. Cesium-137 and Iodine-131 Concentrations in Large Game Animals

Species	Collection		Analyte	Result ± 1s Uncertainty			Result ± 1s Uncertainty			Result > 3s
	Date	Tissue		(pCi/kg wet weight)			(x 10 <sup>-2</sup> Bq/kg wet weight)			
PRONGHORN	5/4/2010	Liver	<sup>131</sup> I	-35.70	±	20.80	-132.09	±	76.96	
			<sup>137</sup> Cs	5.02	±	2.80	18.57	±	10.36	
PRONGHORN	5/4/2010	Muscle	<sup>131</sup> I	3.21	±	10.80	11.88	±	39.96	
			<sup>137</sup> Cs	1.71	±	1.58	6.33	±	5.85	
PRONGHORN	5/4/2010	Thyroid	<sup>131</sup> I	0.97	±	85.40	3.60	±	315.98	
			<sup>137</sup> Cs	118.00	±	84.20	436.60	±	311.54	
PRONGHORN	6/1/2010	Liver	<sup>131</sup> I	-2.10	±	4.52	-7.77	±	16.72	
			<sup>137</sup> Cs	2.77	±	2.03	10.25	±	7.51	
PRONGHORN	6/1/2010	Muscle	<sup>131</sup> I	4.02	±	2.35	14.87	±	8.70	
			<sup>137</sup> Cs	1.61	±	1.08	5.96	±	4.00	
PRONGHORN	6/1/2010	Thyroid	<sup>131</sup> I	-141.00	±	101.00	-521.70	±	373.70	
			<sup>137</sup> Cs	-7.87	±	109.00	-29.12	±	403.30	
PRONGHORN	6/2/2010	Muscle	<sup>131</sup> I	3.99	±	2.04	14.76	±	7.55	
			<sup>137</sup> Cs	2.50	±	0.95	9.25	±	3.53	
PRONGHORN	6/2/2010	Thyroid	<sup>131</sup> I	55.80	±	162.00	206.46	±	599.40	
			<sup>137</sup> Cs	95.30	±	156.00	352.61	±	577.20	
MULE DEER	6/9/2010	Muscle	<sup>131</sup> I	6.29	±	3.09	23.27	±	11.43	
			<sup>137</sup> Cs	1.15	±	1.59	4.26	±	5.88	
MULE DEER	6/24/2010	Liver	<sup>131</sup> I	-288.00	±	1390.00	-1065.60	±	5143.00	
			<sup>137</sup> Cs	45.40	±	111.00	167.98	±	410.70	
MULE DEER	6/24/2010	Muscle	<sup>131</sup> I	11.60	±	14.60	42.92	±	54.02	
			<sup>137</sup> Cs	-0.53	±	1.29	-1.96	±	4.77	
MULE DEER	6/24/2010	Thyroid	<sup>131</sup> I	21.60	±	49.80	79.92	±	184.26	
			<sup>137</sup> Cs	-10.60	±	27.20	-39.22	±	100.64	

Table C-10. Environmental Radiation Measurements

Location	Start Date	End Date	Radiation Measurement $\pm$ 2s Uncertainty mR	Exposure mR/day
<b>BOUNDARY</b>				
ARCO	5/6/2009	11/4/2009	65.0 $\pm$ 12.7	0.36
ATOMIC CITY	5/6/2009	11/4/2009	65.2 $\pm$ 12.8	0.36
BIRCH CREEK	5/6/2009	11/4/2009	58.6 $\pm$ 11.5	0.32
BLUE DOME	5/6/2009	11/4/2009	54.4 $\pm$ 10.7	0.30
HOWE	5/6/2009	11/4/2009	58.9 $\pm$ 11.5	0.32
MONTEVIEW	5/6/2009	11/4/2009	61.5 $\pm$ 12.1	0.34
MUD LAKE	5/6/2009	11/4/2009	70.0 $\pm$ 13.7	0.38
<b>Boundary Average</b>				<b>0.34</b>
<b>DISTANT</b>				
ABERDEEN	5/5/2009	11/3/2009	68.4 $\pm$ 13.4	0.38
BLACKFOOT	5/6/2009	11/4/2009	65.3 $\pm$ 12.8	0.36
BLACKFOOT CMS	5/6/2009	11/4/2009	56.2 $\pm$ 11.0	0.31
CRATERS	5/6/2009	11/4/2009	65.3 $\pm$ 12.8	0.36
DUBOIS	5/6/2009	11/4/2009	53.5 $\pm$ 10.5	0.29
IDAHO FALLS	5/4/2009	11/5/2009	64.2 $\pm$ 12.6	0.35
MINIDOKA	5/5/2009	11/3/2009	58.0 $\pm$ 11.4	0.32
REXBURG	5/6/2009	11/4/2009	73.6 $\pm$ 14.4	0.40
ROBERTS	5/5/2009	11/3/2009	69.0 $\pm$ 13.5	0.38
<b>Distant Average</b>				<b>0.35</b>
<b>OUT-OF-STATE</b>				
JACKSON	5/11/2009	11/2/2009	52.5 $\pm$ 10.3	0.30

**APPENDIX D**  
***STATISTICAL ANALYSIS RESULTS***



**Table D-1. Results of the Kruskal-Wallis statistical test between INL Site, Boundary and Distant sample groups by month.**

<b>Parameter</b>	<b>P<sup>a</sup></b>
<b>Gross Alpha</b>	
Quarter	0.30
April	0.95
May	0.35
June	0.56
<b>Gross Beta</b>	
Quarter	0.94
April	0.79
May	0.99
June	0.94
a. A 'p' value greater than 0.05 signifies no statistical difference between data groups.	



**Table D-2. Statistical difference in weekly gross alpha and gross beta concentrations measured at Boundary and Distant locations.**

Parameter	Mann-Whitney U test	
	Week	P <sup>a</sup>
<b>Gross Alpha</b>		
	April 7	0.32
	April 14	0.57
	April 21	0.20
	April 28	0.48
	May 5	0.67
	May 12	0.06
	May 19	0.57
	May 27	0.32
	June 2	0.15
	June 9	0.78
	June 16	0.48
	June 23	0.39
	June 30	0.33
<b>Gross Beta</b>		
	April 7	0.22
	April 14	0.35
	April 21	0.52
	April 28	0.72
	May 5	0.89
	May 12	1.00
	May 19	0.35
	May 27	0.17
	June 2	0.17
	June 9	0.83
	June 16	0.10
	June 23	0.09
	June 30	0.42

a. A 'p' value greater than 0.05 signifies no statistical difference between data groups.