

Gonzales Stoller Surveillance, LLC  
Environmental Surveillance, Education, and Research Program  
ISSN NUMBER 1089-5469

# Idaho National Laboratory Site Offsite Environmental Surveillance Program Report: Fourth Quarter 2010

April 2011



*Contributors:  
Marilyn Case, Russ Mitchell*

Program conducted for the U.S. Department of Energy, Idaho Operations Office  
Under Contract DE-NE0000300

By Gonzales Stoller Surveillance, LLC  
Environmental Surveillance, Education, and Research Program  
Douglas K. Halford, Program Manager  
120 Technology Dr., Idaho Falls, Idaho 83402  
[www.gsseser.com](http://www.gsseser.com)

## **EXECUTIVE SUMMARY**

Most of the radionuclides detected in samples collected during the fourth quarter of 2010 could be directly linked with INL Site activities. One exception was samples of waterfowl taken directly from wastewater ponds at the INL Site. Except for waterfowl, levels of detected radionuclides were no different than values measured at other locations across the United States or were consistent with levels measured historically at the INL Site. All detected radionuclide concentrations were well below guidelines set by the U.S. Department of Energy (DOE) and regulatory standards established by the U.S. Environmental Protection Agency (EPA) for protection of the public.

This report for the fourth quarter of 2010 contains results from the Environmental Surveillance, Education, and Research (ESER) Program's monitoring of the Department of Energy's Idaho National Laboratory (INL) Site's offsite environment, October 1 through December 31, 2010. All sample types (media) and the sampling schedule followed during 2010 are listed in Appendix A. Specifically, this report contains the results for the following:

- Air sampling, including particulate air filters, charcoal cartridges, and atmospheric moisture
- Precipitation sampling
- Milk sampling
- Lettuce and potato sampling
- Game animal sampling
- Waterfowl sampling
- Soil sampling
- Environmental radiation measurements

**Table E-1 Summary of results for the Fourth Quarter of 2010.**

Media	Sample Type	Analysis	Results
Air	Filters	Gross alpha, gross beta	No statistical differences were noted in either gross alpha or gross beta data on quarterly and monthly comparisons. During one week, gross alpha concentrations were statistically higher at distant locations than boundary locations. No result exceeded the DCG for gross alpha or gross beta activity in air.
		Gamma-emitting radionuclides, select actinides ( $^{241}\text{Am}$ , $^{238}\text{Pu}$ , and $^{239,240}\text{Pu}$ ), $^{90}\text{Sr}$	No man-made gamma-emitting radionuclides, actinides or $^{90}\text{Sr}$ were detected.
	Charcoal Cartridge	Iodine-131	Iodine-131 was not reported on any charcoal cartridge batches during any week of the quarter.
Atmospheric Moisture	Liquid	Tritium	A total of 12 samples were collected. Eight of these samples had tritium results greater than the 3s uncertainty. No sample result exceeded the DCG for tritium in air. Results were consistent with historical measurements.
Precipitation	Liquid	Tritium	Twelve results for samples collected were available. Six of the results were greater than the 3s uncertainty. The concentrations were consistent with those reported across the region by the Environmental Protection Agency and with previous results.
Milk	Liquid	Iodine-131, gamma-emitting radionuclides, $^{90}\text{Sr}$ , tritium	Thirty-five samples, including two duplicates, were collected. No Iodine-131 was detected in any sample. One sample was initially reported with a detectable Cesium-137 result but this was not confirmed by recount or analysis of a duplicate sample. Strontium-90 was detected in four of seven samples and tritium was found in three of seven samples. All results were consistent with historical measurements.
Lettuce	Vegetation	Gamma-emitting radionuclides, $^{90}\text{Sr}$	Ten lettuce samples were collected. Cesium-137 was detected in samples collected from Arco and FAA tower. Strontium-90 was detected in two samples at levels consistent with fallout from weapons testing.
Potatoes	Vegetation	Gamma-emitting radionuclides, $^{90}\text{Sr}$	Ten samples were collected, including one duplicate. No man-made gamma-emitting radionuclides were detected. Strontium-90 was found in one sample from Rupert at just above the detection level.
Large Game Animals	Tissue	Gamma-emitting radionuclides	Four large game animals (two mule deer, one pronghorn, and one elk) were sampled. Cesium-137 was detected in muscle tissue of one deer at a level within background concentrations.
Waterfowl	Tissue	Gamma-emitting radionuclides, select actinides ( $^{241}\text{Am}$ , $^{238}\text{Pu}$ , and $^{239,240}\text{Pu}$ ), $^{90}\text{Sr}$	Seven radionuclides were found in tissues from ducks collected at the Advanced Test Reactor Complex, including edible tissue. Only $^{90}\text{Sr}$ was found in birds from the Materials and Fuels

			Complex and control samples. Concentrations in edible tissues were somewhat higher than those found in the previous few years. The estimated dose from eating the entire edible mass of the duck with the highest concentrations was estimated at 0.059 mrem.
Soil	Solid	Gamma emitting radionuclides, <sup>90</sup> Sr, <sup>241</sup> Am, and plutonium	Cesium-137 was detected in all samples collected. Strontium-90 was detected in half of the samples. The origin of these nuclides is probably deposition of fallout from atmospheric nuclear weapons testing. Americium and Plutonium were not detected in any samples.
Thermo-luminescent Dosimeters	Environmental radiation	External radioactivity	The average measurements over the six-month period were 0.33 mrem/day at boundary and 0.35 mrem/day at distant locations.

### LIST OF ABBREVIATIONS

AEC	Atomic Energy Commission
CFA	Central Facilities Area
DCG	Derived Concentration Guide
DOE	Department of Energy
DOE – ID	Department of Energy Idaho Operations Office
EAL	Environmental Assessment Laboratory
EFS	Experimental Field Station
EPA	Environmental Protection Agency
ERAMS	Environmental Radiation Ambient Monitoring System
ESER	Environmental Surveillance, Education, and Research
GSS	Gonzales Stoller Surveillance, LLC
ICP	Idaho Cleanup Project
INL	Idaho National Laboratory
INEL	Idaho National Engineering Laboratory
INEEL	Idaho National Engineering and Environmental Laboratory
ISU	Idaho State University
MDC	minimum detectable concentration
NRTS	National Reactor Testing Station

## **LIST OF UNITS**

Bq	becquerel
Ci	curie
g	gram
L	liter
$\mu$ Ci	microcurie
mL	milliliter
pCi	picocurie

## **1. ESEER PROGRAM DESCRIPTION**

Operations at the Idaho National Laboratory (INL) Site are conducted under requirements imposed by the U.S. Department of Energy (DOE) under authority of the Atomic Energy Act and the U.S. Environmental Protection Agency (EPA) under a number of acts (e.g. the Clean Air Act and Safe Drinking Water Act). The requirements imposed by DOE are specified in DOE Orders. These requirements include those to monitor the effects of DOE activities both inside and outside the boundaries of DOE facilities (DOE 2003). During calendar year 2010, environmental monitoring within the INL Site boundaries was primarily the responsibility of the INL and Idaho Cleanup Project (ICP) contractors, while monitoring outside the INL Site boundaries was conducted under the Environmental Surveillance, Education, and Research (ESER) Program. The ESER Program was led by the S.M. Stoller Corporation in cooperation with its team members, including the University of Idaho, Idaho State University (ISU), the Wildlife Conservation Society, and Teledyne Brown Engineering. At the end of the fourth quarter of 2010, the ESER Program became led by a new partnership between S.M. Stoller and Jerome Gonzales Management Systems, Inc. with the support of the previous team members. This partnership is named Gonzales Stoller Surveillance, LLC (GSS).

This report contains monitoring results from the ESER Program for samples collected during the fourth quarter of 2010 (October 1-December 31, 2010).

The surveillance portion of the ESER Program is designed to satisfy the following program objectives:

- Verify compliance with applicable environmental laws, regulations, and DOE Orders
- Characterize and define trends in the physical, chemical, and biological condition of environmental media on and around the INL Site
- Assess the potential radiation dose to members of the public from INL Site effluents
- Present program results clearly and concisely through the use of reports, presentations, newsletter articles and press releases.

The goal of the surveillance program is to monitor different media at a number of potential exposure points within the various exposure pathways, including air, water, agricultural products, wildlife, and soil that could possibly contribute to the radiation dose received by the public.

Environmental samples collected include:

- air at 16 locations on and around the INL Site
- moisture in air at four locations around the INL Site
- precipitation from three locations on and around the INL Site
- agricultural products, including milk at six dairies around the INL Site, potatoes from at least five local producers, wheat from approximately 10 local producers, and lettuce from approximately nine home-owned and portable gardens on and around the INL
- soil from 12 locations around the INL Site biennially
- environmental dosimeters from 15 locations semi-annually
- various numbers of wildlife including big game (pronghorn, mule deer, and elk) and waterfowl sampled on and near the INL Site.

Table A-1 in Appendix A lists samples, sampling locations and collection frequency for the ESER Program.

The ESER Program used two laboratories to perform analyses on routine environmental samples collected during the quarter reported here. The ISU Environmental Assessment Laboratory (EAL) performed routine gross alpha, gross beta, tritium, and gamma spectrometry analyses. Analyses requiring radiochemistry including strontium-90 ( $^{90}\text{Sr}$ ), plutonium-238 ( $^{238}\text{Pu}$ ), plutonium-239/240 ( $^{239/240}\text{Pu}$ ), and americium-241 ( $^{241}\text{Am}$ ) were performed by Teledyne Brown Engineering, Inc. of Knoxville, Tennessee.

In the event of non-routine occurrences, such as suspected releases of radioactive material, the ESER Program may increase the frequency of sampling and/or the number of sampling locations based on the nature of the release and wind distribution patterns. Any data found to be outside historical norms in the ESER Program is thoroughly investigated to determine if an INL Site origin is likely. Investigation may include re-sampling and/or re-analysis of prior samples.

In the event of any suspected worldwide nuclear incidents, like the 1986 Chernobyl accident, the EPA may request additional sampling be performed through RadNet [previously known as the Environmental Radiation Ambient Monitoring System (ERAMS) network] (EPA 2009). The EPA established the ERAMS network in 1973 with an emphasis on identifying trends in the accumulation of long-lived radionuclides in the environment. ERAMS was renamed RadNet in 2005 to reflect a new mission. RadNet is comprised of a nationwide network of sampling stations that provide air, precipitation, drinking water, and milk samples. The ESER Program currently operates a high-volume air sampler and collects precipitation and drinking water in Idaho Falls for this national program and routinely sends samples to EPA's Eastern Environmental Radiation Facility for analyses. The RadNet data collected at Idaho Falls are not reported by the ESER Program but are available through the EPA RadNet website (<http://www.epa.gov/narel/radnet/>).

Once samples have been collected and analyzed, the ESER Program has the responsibility for quality control of the data and for preparing quarterly reports on results from the environmental surveillance program. The quarterly reports are then consolidated into the INL Site Environmental Report for each calendar year. These annual reports also include data collected by other INL Site contractors.


The results reported in the quarterly and annual reports are assessed in terms of data quality and statistical significance with respect to laboratory analytical uncertainties, sample locations, reported INL Site releases, meteorological data, and worldwide events that might conceivably have an effect on the INL Site environment. First, field collection and laboratory information are reviewed to determine identifiable errors that would invalidate or limit use of the data. Examples of such limitations include insufficient sample volume, torn filters, evidence of laboratory cross-contamination or quality control issues. Data that pass initial screening are further evaluated using statistical methods. Statistical tools are necessary for data evaluation particularly since environmental measurements typically involve the determination of minute concentrations, which are difficult to detect and even more difficult to distinguish from other measurements.

Results are presented in this report with an analytical uncertainty term,  $s$ , where " $s$ " is the estimated sample standard deviation ( $\sigma$ ), assuming a Gaussian or normal distribution. All results are reported in this document, even those that do not necessarily represent detections. The term "detected", as used for the discussion of results in this report, does not imply any degree of risk to the public or environment, but rather indicates that the radionuclide was measured at a concentration sufficient for the analytical instrument to record a value that is statistically different from background. The ESER has adopted guidelines developed by the United States Geological Survey (Bartholomay, et al. 2003), based on an extension of a method



proposed by Currie (1984), to interpret analytical results and make decisions concerning detection. Most of the following discussion is taken from Bartholomay et al (2003).

Laboratory measurements involve the analysis of a target sample and the analysis of a prepared laboratory blank (i.e., a sample which is identical to the sample collected in the environment, except that the radionuclide of interest is absent). Instrument signals for the target and blank vary randomly about the true signals and may overlap making it difficult to distinguish between radionuclide activities in blank and in environmental samples (Figure 1). That is, the variability around the sample result may substantially overlap the variability around a net activity of zero for samples with no radioactivity. In order to conclude that a radionuclide has been detected, it is essential to consider two fundamental aspects of the problem of detection: (1) the instrument signal for the sample must be greater than that observed for the blank before the decision can be made that the radionuclide has been detected; and (2) an estimate must be made of the minimum radionuclide concentration that will yield a sufficiently large observed signal before the correct decision can be made for detection or non-detection.



**Example of overlap of blank and sample measurement distributions.**

In the laboratory, instrument signals must exceed a critical level of  $1.6s$  before the qualitative decision can be made as to whether the radionuclide was detected in a sample. At  $1.6s$  there is about a 95-percent probability that the correct conclusion—not detected—will be made. Given a large number of samples, approximately 5 percent of the samples with measured concentrations greater than or equal to  $1.6s$ , which were concluded as being detected, might not contain the radionuclide. These are referred to as false positives. For purposes of simplicity and consistency with past reporting, the ESER has rounded the  $1.6s$  critical level estimate to  $2s$ .

Once the critical level has been defined, the minimum detectable concentration may be determined. Concentrations that equal  $3s$  represent a measurement at the detection level or minimum detectable concentration. For true concentrations of  $3s$  or greater, there is a 95-percent probability that the radionuclide was detected in the target sample. In a large number of samples, the conclusion—not detected—will be made in 5 percent of the samples with true concentrations at the minimum detectable concentration of  $3s$ . These measurements are known as false negatives. The ESER reports measured radionuclide concentrations greater than or equal to their respective  $3s$  uncertainties as being “detected with confidence.”

Concentrations between 2s and 3s are reported as “questionably detected”. That is, the radionuclide may be present in the sample; however, the detection may not be reliable. Measurements made between 2s and 3s are examined further to determine if they are a part of a pattern (temporal or spatial) that might warrant further investigation or recounting. For example, if a particular radionuclide is typically detected at  $> 3s$  at a specific location, a sample result between 2s and 3s might be considered detected.

If a result is less than or equal to 2s there is little confidence that the radionuclide is present in the sample. Analytical results in this report are presented as the result value  $\pm$  one standard deviation (1s) for reporting consistency with the annual report. To obtain the 2s or 3s values simply multiply the uncertainty term by 2 or 3.

For more information concerning the ESER Program, contact GSS at (208) 525-8250, or visit the Program’s web page (<http://www.gsseser.com>).

## 2. THE INL SITE

The INL Site is a nuclear energy and homeland security research and environmental management facility. It is owned and administered by the U.S. Department of Energy, Idaho Operations Office (DOE-ID) and occupies about 890 mi<sup>2</sup> (2300 km<sup>2</sup>) of the upper Snake River Plain in Southeastern Idaho. The history of the INL Site began during World War II when the U.S. Naval Ordnance Station was located in Pocatello, Idaho. This station, one of two such installations in the U.S., retooled large guns from U.S. Navy warships. The retooled guns were tested on the nearby, uninhabited plain, known as the Naval Proving Ground. In the years following the war, as the nation worked to develop nuclear power, the Atomic Energy Commission (AEC), predecessor to the DOE, became interested in the Naval Proving Ground and made plans for a facility to build, test, and perfect nuclear power reactors.

The Naval Proving Ground became the National Reactor Testing Station (NRTS) in 1949, under the AEC. By the end of 1951, a reactor at the NRTS became the first to produce useful amounts of electricity. Over time the site has operated 52 various types of reactors, associated research centers, and waste handling areas. The NRTS was renamed the Idaho National Engineering Laboratory (INEL) in 1974, and the Idaho National Engineering and Environmental Laboratory (INEEL) in January 1997. With renewed interest in nuclear power the DOE announced in 2003 that Argonne National Laboratory and the INEEL would be the lead laboratories for development of the next generation of power reactors. On February 1, 2005 the INEEL and Argonne National Laboratory-West became the INL. The INL is committed to providing international nuclear leadership for the 21st Century, developing and demonstrating compelling national security technologies, and delivering excellence in science and technology as one of the Department of Energy's multiprogram national laboratories.

The cleanup operation, the ICP, is now a separately managed effort. The ICP is charged with safely and cost-effectively completing the majority of cleanup work from past laboratory missions by 2012.




### 3. AIR SAMPLING

The primary pathway by which radionuclides can move off the INL Site is through the air and for this reason the air pathway is the primary focus of monitoring on and around the INL Site. Samples for particulates and iodine-131 ( $^{131}\text{I}$ ) gas in air were collected weekly for the duration of the quarter at 16 locations using low-volume air samplers. Moisture in the atmosphere was sampled at four locations around the INL Site and analyzed for tritium. Air sampling activities and results for the fourth quarter of 2010 are discussed below. A summary of approximate minimum detectable concentrations (MDCs) for radiological analyses and DOE Derived Concentration Guide (DCG) (DOE 1993) values is provided in Appendix B.

#### LOW-VOLUME AIR SAMPLING

Radioactivity associated with airborne particulates was monitored continuously by 18 low-volume air samplers (two of which are used as replicate samplers) at 16 locations during the fourth quarter of 2010 (Figure 2). Four of these samplers are located on the INL Site, seven are situated off the INL Site near the boundary, and seven have been placed at locations distant to the INL Site. Samplers are divided into INL Site, Boundary, and Distant groups to determine if there is a gradient of radionuclide concentrations, increasing towards the INL Site. Each replicate sampler is relocated every other year to a new location. During 2010, one replicate sampler was operating in Dubois (a Distant location) and one was operating at the Van Buren gate (an INL Site location). An average of 19,017 ft<sup>3</sup> (539 m<sup>3</sup>) of air was sampled at each location, each week, at an average flow rate of 1.89 ft<sup>3</sup>/min (0.05 m<sup>3</sup>/min). Particulates in air were collected on membrane particulate filters (1.2- $\mu\text{m}$  pore size). Gases passing through the filter were collected with an activated charcoal cartridge.



Low-volume air sampler locations.

Filters and charcoal cartridges were changed weekly at each station during the quarter. Each particulate filter was analyzed for gross alpha and gross beta radioactivity using thin-window gas flow proportional counting systems after waiting about four days for naturally-occurring daughter products of radon and thorium to decay.

The weekly particulate filters collected during the quarter for each location were composited and analyzed for gamma-emitting radionuclides. Selected composites were also analyzed by location for  $^{90}\text{Sr}$ ,  $^{238}\text{Pu}$ ,  $^{239/240}\text{Pu}$ , and  $^{241}\text{Am}$  as determined by a rotating quarterly schedule.

Charcoal cartridges were analyzed for gamma-emitting radionuclides, specifically for iodine-131 ( $^{131}\text{I}$ ). Iodine-131 is of particular interest because it is produced in relatively large quantities by nuclear fission, is readily accumulated in human and animal thyroids, and has a half-life of eight days. This means that any elevated level of  $^{131}\text{I}$  in the environment could be from a recent release of fission products.

Gross alpha results are reported in Table C-1. Median gross alpha concentrations in air for INL Site, Boundary, and Distant locations for the fourth quarter of 2010 are shown in Figure 3. Gross alpha data are tested for normality prior to statistical analyses, and generally show no consistent discernable distribution. Box and whisker plots are commonly used when there is no assumed distribution. Each data group in Figure 3 is presented as a box and whisker plot, with a median (small red square), a box enclosing values between the 25<sup>th</sup> and 75<sup>th</sup> percentiles, and whiskers representing the non-outlier range. Outliers and extreme values are identified separately from the box and whiskers. Outliers and extreme values are atypical, infrequent data points that are far from the middle of the data distribution. For this report, outliers are defined as values that are greater than 1.5 times the height of the box, above, or below the box. Extreme values are greater than 2 times the height of the box, above or below the box. Outliers and extreme values may reflect inherent variability, may be due to errors associated with transcription or measurement, or may be related to other anomalies. A careful review of the data collected during the fourth quarter indicates that the outlier values were not due to mistakes in collection, analysis, or reporting procedures, but rather reflect natural variability in the measurements. Thus, rather than dismissing the outliers, they were included in the subsequent statistical analyses.

Because there is no discernable distribution of the data, the nonparametric Kruskal-Wallis test of multiple independent groups was used to test for statistical differences between INL Site, Boundary, and Distant locations. The use of nonparametric tests, such as Kruskal-Wallis, gives less weight to outlier and extreme values thus allowing a more appropriate comparison of data groups. A statistically significant difference exists between data groups if the (p) value is less than 0.05. Values greater than 0.05 translate into a 95 percent confidence that the medians are statistically the same. The p-value for each comparison is shown in Table D-1. For the quarter, there was no statistical difference noted. Figure 3 graphically shows that the gross alpha measurements made at INL Site, Boundary, and Distant locations are similar for the fourth quarter. If the INL Site were a significant source of offsite contamination, concentrations of contaminants could be statistically greater at Boundary locations than at Distant locations.

Comparisons of gross alpha concentrations were made for each month of the quarter (Figures 4 – 6). Again the Kruskal-Wallis test of multiple independent groups was used to determine if statistical differences exist between INL Site, Boundary, and Distant data groups. No statistical differences in gross alpha concentrations between groups were noted during any month of the quarter (Table D-1).




Figure 3. Gross alpha concentrations in air at ESER INL Site, Boundary and Distant locations for the fourth quarter of 2010.




Figure 4. October gross alpha concentrations in air at ESER INL Site, Boundary, and Distant locations. Number of samples (N) = 4 at each location.






Figure 5. November gross alpha concentrations in air at ESER INL Site, Boundary, and Distant locations. Number of samples (N) = 4 at each location, except FAA Tower (N = 2).




Figure 6. December gross alpha concentrations in air at ESER INL Site, Boundary, and Distant locations. Number of samples (N) = 5 at each location, except FAA Tower (N = 3).

As an additional check, comparisons between gross alpha concentrations measured at Boundary and Distant locations were made on a weekly basis. The Mann-Whitney U test was used to compare the Boundary and Distant data because it is the most powerful nonparametric alternative to the t-test for independent samples. INL Site sample results were not included in this analysis because the onsite data, collected at only three locations, are not representative of the entire INL Site and would not aid in determining offsite impacts. In the fourth quarter, there was one week where a statistical difference existed between the two sample groups (Table D-2). During the week of October 6, the Distant group was statistically higher than the Boundary group. Examination of data from this week does not show any unusual pattern, with Jackson being the highest location and Mud Lake and Arco the lowest.

Gross beta results are presented in Table C-1. Gross beta concentrations in air for INL Site, Boundary, and Distant locations for the fourth quarter of 2010 are shown in Figure 7. The data were tested and found to be neither normally nor log-normally distributed. Box and whiskers plots were used for presentation of the data. Outliers and extreme values were retained in subsequent statistical analyses because they are within the range of measurements made in the past five years, and because these values could not be attributed to mistakes in collection, analysis, or reporting procedures. No statistical differences were noted in the quarterly data using the Kruskal-Wallis test (Table D-1).

Monthly median gross beta concentrations in air for each sampling group are shown in Figures 8 – 10. Statistical data are presented in Table D-1. No statistical differences were noted during any month.

Comparison of weekly Boundary and Distant gross beta data sets, using the Mann Whitney U test, showed no statistical differences between Boundary and Distant measurements during any week of the quarter (Table D-1).

Iodine-131 was not reported above the detection level on any charcoal cartridge batches analyzed during the quarter. Weekly  $^{131}\text{I}$  results for each location are listed in Table C-2 of Appendix C. Gamma spectrographic analysis is also done with the  $^{131}\text{I}$  analysis. Cesium-137 was not detected in any of the 26 measured batches of cartridges this quarter.

Weekly filters for the fourth quarter of 2010 were composited by location. All samples were analyzed for gamma-emitting radionuclides, including  $^{137}\text{Cs}$  and  $^{90}\text{Sr}$ . No manmade gamma-emitting radionuclides, actinides ( $^{238}\text{Pu}$ ,  $^{239/240}\text{Pu}$ , and  $^{241}\text{Am}$ ) or  $^{90}\text{Sr}$  were found on any of the composites. Analysis results are shown in Table C-3, Appendix C.

#### **ATMOSPHERIC MOISTURE SAMPLING**

Results were available for 12 atmospheric moisture samples collected during the fourth quarter of 2010 from Atomic City, Blackfoot, Idaho Falls, and Rexburg. Atmospheric moisture is collected by pulling air through a column of absorbent material (molecular sieve material) to absorb water vapor. The water is then extracted from the absorbent material by heat distillation. The resulting water samples are then analyzed for tritium using liquid scintillation.

Eight of the 12 samples exceeded the 3s uncertainty level for tritium, with similar results to those reported previously. All samples were significantly below the DOE DCG for tritium in air of  $1 \times 10^{-7} \mu\text{Ci}/\text{mL}_{\text{air}}$  with a maximum reported value of  $25.75 \times 10^{-13} \mu\text{Ci}/\text{mL}_{\text{air}}$  at Atomic City. All results are shown in Table C-4, Appendix C.




Figure 7. Gross beta concentrations in air at ESER INL Site, Boundary, and Distant locations for the fourth quarter of 2010.




Figure 8. October gross beta concentrations in air at ESER INL Site, Boundary, and Distant locations. Number of samples (N) = 4 at each location.




Figure 9. November gross beta concentrations in air at ESER INL Site, Boundary and Distant locations. Number of samples (N) = 4 at each location, except FAA Tower (N = 2).




Figure 10. December gross beta concentrations in air at ESER INL Site, Boundary, and Distant locations. Number of samples (N) = 5 at each location, except FAA Tower (N = 3).

## 4. PRECIPITATION SAMPLING

### *PRECIPITATION SAMPLING*

Precipitation samples are gathered when sufficient precipitation occurs to allow for the collection of the minimum sample volume of approximately 50 mL. Samples are taken of monthly composites from Idaho Falls and CFA, and weekly from the EFS. Precipitation samples are analyzed for tritium. Storm events in the fourth quarter of 2010 produced sufficient precipitation to yield 14; results were available for 12 of the samples.

Tritium was measured above the 3s values in 6 of the 12 samples for which results were available. Low levels of tritium exist in the environment at all times as a result of cosmic ray reactions with water molecules in the upper atmosphere. The EPA's RadNet program collects precipitation samples from across the United States. From 1980 to 2008, tritium measured in samples from Region 10 (which includes Idaho) averaged 117 pCi/L (EPA 2009). Data for all fourth quarter precipitation samples collected by the ESER Program were just about at this value (averaging 115 pCi/L) and are listed in Table C-5 (Appendix C).





## 5. AGRICULTURAL PRODUCT, WILDLIFE, AND SOIL SAMPLING

Another potential pathway for contaminants to reach humans is through the food chain. The ESER Program samples multiple agricultural products and game animals from around the INL Site and Southeast Idaho. Specifically, milk, grain, potatoes, lettuce, large game animals, and waterfowl are sampled. Milk is sampled throughout the year and large game animals are sampled whenever large game animals are killed onsite from vehicle collisions. Lettuce and grain are sampled during the third quarter, while potatoes are collected during the fourth quarter. Waterfowl are collected in either the third or fourth quarter. See Table A-1, Appendix A, for more details on agricultural product and wildlife sampling. This section discusses results from milk, grain, soil, and large game animal samples collected during the fourth quarter of 2010.

### MILK SAMPLING

Milk samples were collected weekly in Idaho Falls. Monthly samples were collected at six other locations around the INL Site (Figure 11) during the fourth quarter of 2010. In addition, commercially-available organic milk was purchased as a control sample. All samples were analyzed for gamma emitting radionuclides. No Iodine-131 was detected in any sample. Cesium-137 was reported in one sample from Howe at just above the detection level. However,  $^{137}\text{Cs}$  was not confirmed in either a subsequent recount of the sample or in a lab split of the sample. Data for  $^{131}\text{I}$  and  $^{137}\text{Cs}$  in milk samples are listed in Appendix C, Table C-6.

Strontium-90 was detected in four of the seven samples analyzed (including the control sample) at levels within historical measurements, ranging from 0.25 pCi/L (Rupert) to 0.86 pCi/L (Idaho Falls). Tritium was detected in three of the seven samples analyzed, again all within historical measurements (see Table C-7 in Appendix C).




Figure 11. ESER milk sampling locations.

### **LETTUCE SAMPLING**

Lettuce sampling final results were completed during the fourth quarter. Five lettuce samples were collected from prototype self-watering planters and five samples (including a duplicate) were collected from private gardens. Each sample was analyzed for gamma-emitting radionuclides and <sup>90</sup>Sr. No gamma results were measured above the 3s uncertainty value. Strontium-90 was reported in four of the samples. In addition, a commercially available lettuce sample was obtained.

A review of Table C-8, performed in the spring of 2021, identified the <sup>137</sup>Cs result and uncertainty values listed were incorrect. The incorrect values appear to be due to inadvertently listing the 1-sigma and 2-sigma uncertainty values as the <sup>137</sup>Cs result and uncertainty values for the analyte. The result and uncertainty values were updated with the correct values. Cesium-137 was detected in the samples collected from Arco and FAA tower. In addition, the review identified the unit of concentration listed in the column headings were incorrect. Prior to 2010, concentrations were reported in either pCi/g or pCi/kg. In 2010, the concentration unit of pCi/kg was adopted for reporting radionuclide concentrations in soil and biota (vegetation and animals). The reasons for doing this include: 1) the use of one unit (pCi/kg) ensures consistency and comparability in reporting concentrations in various media, 2) the use of one unit (pCi/kg) minimizes mistakes (due to confusion about units) in data entry into the database, and 3) the unit of pCi/kg was selected because it is the unit associated with models that are used for dose calculations and the results tend to be whole numbers (e.g. 14 pCi/kg versus 0.014 pCi/g). The column headings have been updated to the correct units of concentration (pCi/kg and Bq/kg). Strontium-90 was found in two of the samples analyzed. The quantities detected are consistent with those found in previous years, which are attributed to uptake of soil with residual <sup>90</sup>Sr from nuclear weapons testing that took place between 1945 and 1980.

Data for <sup>137</sup>Cs and <sup>90</sup>Sr in all lettuce samples taken during the third quarter are listed in Table C-8 (Appendix C).

### **POTATO SAMPLING**

Ten potato samples were collected from area growers and from an out-of-state location (Oregon). All samples were analyzed for gamma emitting radionuclides and <sup>90</sup>Sr. Strontium-90 was found in the sample from Rupert at just above the detection level. No other manmade radionuclides were detected in 2009 samples.

Data for <sup>137</sup>Cs and <sup>90</sup>Sr in all potato samples taken during the fourth quarter are listed in Table C-9 (Appendix C).

### **LARGE GAME ANIMAL SAMPLING**

Four large game animals (two mule deer, one pronghorn, and one elk) were available for sampling during the fourth quarter. Cesium-137 was detected in the muscle sample collected from the pronghorn at a level just above the detection limit. The reported value of 2.77 pCi/kg was slightly below the background range of concentrations (4 to 12 pCi/kg allowing for decay of <sup>137</sup>Cs) found in a 1999 study on tissue samples from game animals across the western United States. Results are presented in Appendix C, Table C-10.

### **WATERFOWL SAMPLING**

Nine ducks were collected during 2009. Three each were collected from wastewater ponds located at the Advanced Test Reactor Complex (ATR Complex) and near the Materials and Fuels Complex (MFC), and three control samples were collected from the Fort Hall Bottoms area. Each sample was divided into the following three sub-samples: 1) edible tissue (muscle,

gizzard, heart and liver), 2) external portion (feathers, feet and head), and 3) all remaining tissue. All were analyzed for gamma-emitting radionuclides,  $^{90}\text{Sr}$ , plutonium-238 ( $^{238}\text{Pu}$ ), plutonium-239,240 ( $^{239/240}\text{Pu}$ ), and americium-241 ( $^{241}\text{Am}$ ). Concentrations of radionuclides measured in the edible tissues of 2010 waterfowl are shown in Table C-11 (Appendix C).

During the spring of 2021, a review of Table C-11 determined the unit of concentration listed in the column headings were incorrect. Prior to 2010, concentrations were reported in either pCi/g or pCi/kg. In 2010, the concentration unit of pCi/kg was adopted for reporting radionuclide concentrations in soil and biota (vegetation and animals). The reasons for doing this include: 1) the use of one unit (pCi/kg) ensures consistency and comparability in reporting concentrations in various media, 2) the use of one unit (pCi/kg) minimizes mistakes (due to confusion about units) in data entry into the database, and 3) the unit of pCi/kg was selected because it is the unit associated with models that are used for dose calculations and the results tend to be whole numbers (e.g. 14 pCi/kg versus 0.014 pCi/g). The column headings have been updated to the correct units of concentration (pCi/kg and Bq/kg).

Several man-made radionuclides were detected in the samples from the ATR Complex ponds, including  $^{241}\text{Am}$ ,  $^{137}\text{Cs}$ , cobalt-60 ( $^{60}\text{Co}$ ),  $^{238}\text{Pu}$ ,  $^{239/240}\text{Pu}$ ,  $^{90}\text{Sr}$ , and zinc-65 ( $^{65}\text{Zn}$ ). All of these radionuclides except  $^{241}\text{Am}$  were also found in at least one edible tissue sample. Samples from MFC ponds and the control samples contained only  $^{90}\text{Sr}$  and only the control samples contained this nuclide in edible tissue.

Because human-made radionuclides were found in ducks from the ATR Complex and for the most part not at MFC or control locations, it is assumed that radioactive wastewater ponds at the ATR Complex are the source of these radionuclides. The ducks were not taken directly from the two-celled hypalon-lined radioactive wastewater evaporation ponds, but rather from an adjacent sewage lagoon. However, the ducks probably also used the evaporation pond. Concentrations of many of the detected radionuclides from the Advanced Test Reactor Complex were slightly higher to those from 2006 through 2009, but  $^{137}\text{Cs}$  concentrations were significantly lower than in 2005. In addition, concentrations were lower in 2010 than those of a 1994-1998 study (Warren et al. 2001).

Waterfowl hunting is not allowed on the INL Site, but a maximum potential exposure scenario to humans would be someone collecting a contaminated duck directly from the ponds and immediately consuming all muscle, liver, heart, and gizzard tissue. The maximum potential dose from eating 225 g (8 oz) of meat from the most contaminated waterfowl collected in 2010 was estimated to be 0.059 mrem. This dose is lower than dose estimates for some previous periods. The maximum dose estimated for the period from 1993 through 1998 was 0.89 mrem and from 2000 through 2004 was 0.08 mrem. In the late 1970s, when the percolation ponds were still in use, the maximum dose estimated from eating a contaminated duck was estimated to be 54 mrem.

#### **SOIL SAMPLING**

Fourteen soil samples (including one duplicate at Atomic City) were collected at boundary and offsite locations in the third quarter and full results were available in the fourth quarter. A new sampling location was established in 2010 at Frenchman's Cabin, located near the southern boundary of the INL Site. This site has been the location of the hypothetical maximally exposed individual for the past several years.


All samples were analyzed for gamma-emitting radionuclides,  $^{241}\text{Am}$ ,  $^{238}\text{Pu}$ ,  $^{239/240}\text{Pu}$ , and  $^{90}\text{Sr}$  (Table C-12). Cesium-137 was detected in all samples at concentrations consistent with historical measurements and is most likely present from past atmospheric nuclear weapons

testing fallout. Similarly  $^{90}\text{Sr}$ , another fallout radionuclide, was detected in seven of the 14 soil samples at levels within historical measurements.

None of the transuranic radionuclides were detected in any of the samples.

## 6. ENVIRONMENTAL RADIATION

An array of thermoluminescent dosimeters (TLDs) is distributed throughout the Eastern Snake River Plain to monitor for environmental radiation (Figure 12). TLDs are changed out in May and again in November after six months in the field. The results of the TLDs exposed from May through October 2010 are discussed below.



**Figure 12. TLD locations.**

Similar to the low-volume air results the environmental dosimeter locations are also divided into Boundary and Distant groupings. Boundary average exposure rates ranged from a low of 0.29 mR/day at Blue Dome to a high of 0.37 mR/day at Mud Lake. The overall Boundary average was 0.33 mR/day. The Distant group had a high of 0.45 mR/day at Rexburg and a low of 0.29 mR/day at the Dubois location. The overall average Distant value was 0.35 mR/day. There was no statistical difference between Boundary and Distant locations and all values are consistent with past readings. All results are listed in Appendix C, Table C-13.

## 7. QUALITY ASSURANCE

The ESER Quality Assurance Program consists of five ongoing tasks which measure:

1. method uncertainty
2. data completeness
3. data accuracy, using spike, performance evaluation and laboratory control samples
4. data precision, using split samples, duplicate samples and recounts
5. presence of contamination in samples, using blanks.

Data completeness for sample collection and delivery was 100 percent during the fourth quarter, with the following exceptions.

There were four air samples that had volumes below the 7,000 ft<sup>3</sup> or 200 m<sup>3</sup> threshold listed in the air sampling procedure as being a valid sample. All were from the FAA Tower due to a power outage. The power had to be reset from inside the building by personnel of the FAA and then reset outside the building by ESER personnel.

Two milk samples were not collected in the fourth quarter. The cow at Fort Hall was dry during one month and the goat dairy in Blackfoot was not operating in November.

Sample results are compared to criteria described in the Quality Assurance Project Plan for the INL Site Offsite Environmental Surveillance Program (Stoller 2007). The following table summarizes the results of the quality assurance program for the fourth quarter of 2010.

<b>QA Sample Type</b>	<b>Number of Sample Results</b>	<b>Number of Results Meeting Criteria</b>	<b>Percentage Meeting Criteria</b>
Spikes/Laboratory Control Samples	249	246	98.8
Field Duplicates	66	62	93.9
Laboratory Splits	31	31	100
Recounts	181	178	98.3
Blanks	75	70	93.3
Method Uncertainty	1699	1677	98.7

## 8. REFERENCES

- Bartholomay, R.C., Knobel, L.L., and Rousseau, J.P., 2003, *Field Methods and Quality Plan for Quality-of-Water Activities, U.S. Geological Survey, Idaho National Engineering and Environmental Laboratory, Idaho*, DOE/ID-22182, January 2003.
- Currie, L.A., 1984, *Lower Limit of Detection: Definition and Elaboration of a Proposed Position for Radiological Effluent and Environmental Measurements*, NUREG/CR-4007, U.S. Nuclear Regulatory Commission, Washington, D.C., September 1984.
- DOE, 2003, "Environmental Management System," U.S. Department of Energy Order 450.1, January 2003.
- DOE, 1993, "Radiation Protection of the Public and the Environment," U.S. Department of Energy Order 5400.5, January 1993.
- EPA, 2009, RadNet—Tracking Environmental Radiation Nationwide, Web-page: <http://www.epa.gov/narel/radnet/>
- Stoller, 2007, *Quality Assurance Project Plan for the INL Site Offsite Environmental Surveillance Program*, Environmental Surveillance, Education and Research Program, February 2007.



**APPENDIX A**  
***SUMMARY OF SAMPLING SCHEDULE***

**Table A-1. Summary of the ESER Program's Sampling Schedule**

Sample Type Analysis	Collection Frequency	LOCATIONS		
		Distant	Boundary	INL Site
<b>AIR SAMPLING</b>				
<i>LOW-VOLUME AIR</i>				
Gross Alpha, Gross Beta, <sup>131</sup> I	weekly	Blackfoot, Craters of the Moon, Dubois, Idaho Falls, Jackson WY, Rexburg	Arco, Atomic City, FAA Tower, Howe, Montevieu, Mud Lake, Blue Dome	Main Gate, EFS, Van Buren
Gamma Spec	quarterly	Blackfoot, Craters of the Moon, Dubois, Idaho Falls, Jackson WY, Rexburg	Arco, Atomic City, FAA Tower, Howe, Montevieu, Mud Lake, Blue Dome	Main Gate, EFS, Van Buren
<sup>90</sup> Sr, Transuranics	quarterly	Rotating schedule	Rotating schedule	Rotating schedule
<i>ATMOSPHERIC MOISTURE</i>				
Tritium	2 to 13 weeks	Blackfoot, Idaho Falls, Rexburg	Atomic City	None
<i>PRECIPITATION</i>				
Tritium	monthly	Idaho Falls	None	CFA
Tritium	weekly	None	None	EFS
<b>ENVIRONMENTAL RADIATION SAMPLING</b>				
<i>TLDS</i>				
Gamma Radiation	semiannual	Aberdeen, Blackfoot (2), Craters of the Moon, Dubois, Idaho Falls, Jackson WY, Minidoka, Rexburg, Roberts	Arco, Atomic City, Birch Creek, Blue Dome, Howe, Montevieu, Mud Lake	None
<b>SOIL SAMPLING</b>				
<i>SOIL</i>				
Gamma Spec, <sup>90</sup> Sr, Transuranics	biennially	Carey, Crystal Ice Caves (Aberdeen), Blackfoot, St. Anthony	Butte City, Montevieu, Atomic City, FAA Tower, Howe, Mud Lake (2), Birch Creek, Frenchman's Cabin	None

**Table A-1. Summary of the ESER Program's Sampling Schedule (continued)**

Sample Type Analysis	Collection Frequency	LOCATIONS		
		Distant	Boundary	INL Site
<b>FOODSTUFF SAMPLING</b>				
<i>MILK</i>				
Gamma Spec ( <sup>131</sup> I)	weekly	Idaho Falls	None	None
Gamma Spec ( <sup>131</sup> I)	monthly	Blackfoot, Dietrich, Fort Hall, Idaho Falls, Minidoka	Howe, Terreton	None
Tritium, <sup>90</sup> Sr	Semi-annually	Blackfoot, Dietrich, Fort Hall, Idaho Falls, Minidoka	Howe, Terreton	None
<i>POTATOES</i>				
Gamma Spec, <sup>90</sup> Sr	annually	Blackfoot, Idaho Falls, Rupert, Shelley, occasional samples across the U.S.	Arco, Montevieu, Mud Lake, Terreton	None
<i>GRAIN</i>				
Gamma Spec, <sup>90</sup> Sr	annually	American Falls, Blackfoot, Carey Idaho Falls, Minidoka, Roberts	Arco, Montevieu, Mud Lake, Taber, Terreton	None
<i>LETTUCE</i>				
Gamma Spec, <sup>90</sup> Sr	annually	Blackfoot, Carey, Idaho Falls	Arco, Atomic City, FAA Tower, Howe, Montevieu	EFS
<i>BIG GAME</i>				
Gamma Spec	varies	Occasional samples across the U.S.	Public Highways	INL Site roads
<i>WATERFOWL</i>				
Gamma Spec, <sup>90</sup> Sr, Transuranics	annually	Varies among: Heise, Firth, Fort Hall, Mud Lake, Market Lake, and American Falls	None	INL Site wastewater disposal ponds

**APPENDIX B**  
***SUMMARY OF MDCs AND DCGs***

**Table B-1. Summary of Approximate Minimum Detectable Concentrations for Radiological Analyses Performed During Fourth Quarter 2010**

Sample Type	Analysis	Approximate Minimum Detectable Concentration <sup>a</sup> (MDC)	Derived Concentration Guide <sup>b</sup> (DCG)
<b>Air</b> (particulate filter) <sup>e</sup>	Gross alpha <sup>c</sup>	4.74 x 10 <sup>-16</sup> μCi/mL	2 x 10 <sup>-14</sup> μCi/mL
	Gross beta <sup>d</sup>	1.46 x 10 <sup>-15</sup> μCi/mL	3 x 10 <sup>-12</sup> μCi/mL
	Specific gamma ( <sup>137</sup> Cs)	1.69 x 10 <sup>-16</sup> μCi/mL	4 x 10 <sup>-10</sup> μCi/mL
	<sup>238</sup> Pu	1.77 x 10 <sup>-18</sup> μCi/mL	3 x 10 <sup>-14</sup> μCi/mL
	<sup>239/240</sup> Pu	9.93 x 10 <sup>-19</sup> μCi/mL	2 x 10 <sup>-14</sup> μCi/mL
	<sup>241</sup> Am	5.02 x 10 <sup>-18</sup> μCi/mL	2 x 10 <sup>-14</sup> μCi/mL
	<sup>90</sup> Sr	4.00 x 10 <sup>-17</sup> μCi/mL	9 x 10 <sup>-12</sup> μCi/mL
<b>Air</b> (charcoal cartridge) <sup>e</sup>	<sup>131</sup> I	1.58 x 10 <sup>-15</sup> μCi/mL	4 x 10 <sup>-10</sup> μCi/mL
<b>Air</b> (atmospheric moisture)	<sup>3</sup> H	59.3 pCi/L <sub>water</sub>	1 x 10 <sup>-7</sup> μCi/mL <sub>air</sub>
<b>Air</b> (precipitation)	<sup>3</sup> H	102.8 pCi/L	2 x 10 <sup>-3</sup> μCi/mL
<b>Milk</b>	<sup>131</sup> I	0.63 pCi/L	--
	<sup>137</sup> Cs	1.17 pCi/L	--
	<sup>3</sup> H	54.1 pCi/L	--
	<sup>90</sup> Sr	0.34 pCi/L	--
<b>Lettuce</b>	<sup>137</sup> Cs	63 pCi/kg	--
	<sup>90</sup> Sr	32.4 pCi/kg	--
<b>Potatoes</b>	<sup>137</sup> Cs	1.34 pCi/kg	--
	<sup>90</sup> Sr	1.4 pCi/kg	--
<p>a The MDC is an estimate of the concentration of radioactivity in a given sample type that can be identified with a 95 percent level of confidence and precision of plus or minus 100 percent under a specified set of typical laboratory measurement conditions.</p> <p>b DCGs, set by the DOE, represent reference values for radiation exposure. They are based on a radiation dose of 100 mrem/yr for exposure through a particular exposure mode such as direct exposure, inhalation, or ingestion of water.</p> <p>c The DCG for gross alpha is equivalent to the DCGs for <sup>239,240</sup>Pu and <sup>241</sup>Am.</p> <p>d The DCG for gross beta is equivalent to the DCGs for <sup>228</sup>Ra</p> <p>e The approximate MDC is based on an average filtered air volume (pressure corrected) of 445 m<sup>3</sup>/week.</p>			

**APPENDIX C**  
***SAMPLE ANALYSIS RESULTS***

TABLE C-1. Weekly Gross Alpha and Gross Beta Concentrations in Air

Sampling Group and Location	Sampling Date	GROSS ALPHA							GROSS BETA				
		Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)		Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)		Result > 3s	Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)		Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)		Result > 3s		
<b>BOUNDARY</b>													
ARCO	10/6/2010	1.69 ± 0.26	6.25 ± 0.94	Y	43.60 ± 1.02	161.32 ± 3.77	Y						
	10/13/2010	1.47 ± 0.25	5.44 ± 0.92	Y	32.00 ± 0.90	118.40 ± 3.32	Y						
	10/20/2010	2.37 ± 0.28	8.77 ± 1.05	Y	47.10 ± 1.09	174.27 ± 4.03	Y						
	10/27/2010	1.36 ± 0.19	5.03 ± 0.70	Y	24.50 ± 0.69	90.65 ± 2.56	Y						
	11/3/2010	0.98 ± 0.17	3.64 ± 0.61	Y	25.80 ± 0.70	95.46 ± 2.57	Y						
	11/10/2010	1.69 ± 0.20	6.25 ± 0.74	Y	33.00 ± 0.78	122.10 ± 2.88	Y						
	11/17/2010	0.58 ± 0.16	2.15 ± 0.58	Y	23.40 ± 0.66	86.58 ± 2.43	Y						
	11/24/2010	0.44 ± 0.16	1.62 ± 0.57	Y	13.10 ± 0.46	48.47 ± 1.70	Y						
	12/1/2010	1.74 ± 0.21	6.44 ± 0.76	Y	38.40 ± 0.83	142.08 ± 3.06	Y						
	12/8/2010	1.37 ± 0.21	5.07 ± 0.78	Y	28.80 ± 0.79	106.56 ± 2.91	Y						
	12/15/2010	1.47 ± 0.20	5.44 ± 0.73	Y	21.70 ± 0.66	80.29 ± 2.43	Y						
	12/22/2010	1.16 ± 0.17	4.29 ± 0.64	Y	27.00 ± 0.70	99.90 ± 2.60	Y						
	12/29/2010	0.57 ± 0.17	2.11 ± 0.61	Y	27.90 ± 0.71	103.23 ± 2.61	Y						
	ATOMIC CITY	10/6/2010	2.65 ± 0.27	9.81 ± 1.00	Y	43.40 ± 0.95	160.58 ± 3.50	Y					
10/13/2010		2.13 ± 0.25	7.88 ± 0.92	Y	33.80 ± 0.83	125.06 ± 3.07	Y						
10/20/2010		2.34 ± 0.29	8.66 ± 1.06	Y	54.00 ± 1.16	199.80 ± 4.29	Y						
10/27/2010		1.19 ± 0.16	4.40 ± 0.60	Y	26.40 ± 0.64	97.68 ± 2.35	Y						
11/3/2010		1.13 ± 0.15	4.18 ± 0.55	Y	26.00 ± 0.60	96.20 ± 2.20	Y						
11/10/2010		1.91 ± 0.19	7.07 ± 0.70	Y	38.70 ± 0.74	143.19 ± 2.73	Y						
11/17/2010		0.56 ± 0.13	2.08 ± 0.48	Y	16.60 ± 0.50	61.42 ± 1.85	Y						
11/24/2010		0.59 ± 0.12	2.20 ± 0.45	Y	11.40 ± 0.35	42.18 ± 1.28	Y						
12/1/2010		0.92 ± 0.13	3.39 ± 0.50	Y	30.60 ± 0.63	113.22 ± 2.32	Y						
12/8/2010		0.80 ± 0.13	2.96 ± 0.47	Y	12.60 ± 0.43	46.62 ± 1.60	Y						
12/15/2010		0.67 ± 0.12	2.47 ± 0.44	Y	9.02 ± 0.38	33.37 ± 1.40	Y						
12/22/2010		1.01 ± 0.13	3.74 ± 0.47	Y	23.50 ± 0.52	86.95 ± 1.91	Y						
12/29/2010		0.33 ± 0.13	1.22 ± 0.50	Y	19.20 ± 0.56	71.04 ± 2.07	Y						
BLUE DOME		10/6/2010	2.12 ± 0.26	7.84 ± 0.97	Y	36.60 ± 0.92	135.42 ± 3.40	Y					
	10/13/2010	0.95 ± 0.21	3.52 ± 0.76	Y	29.20 ± 0.80	108.04 ± 2.97	Y						
	10/20/2010	2.22 ± 0.27	8.21 ± 0.99	Y	42.80 ± 1.02	158.36 ± 3.77	Y						
	10/27/2010	0.78 ± 0.15	2.89 ± 0.54	Y	25.70 ± 0.65	95.09 ± 2.40	Y						
	11/3/2010	0.89 ± 0.23	3.30 ± 0.84	Y	24.00 ± 0.94	88.80 ± 3.49	Y						
	11/10/2010	1.04 ± 0.15	3.85 ± 0.56	Y	32.60 ± 0.70	120.62 ± 2.59	Y						
	11/17/2010	0.39 ± 0.14	1.45 ± 0.53	Y	17.00 ± 0.58	62.90 ± 2.16	Y						
	11/24/2010	0.70 ± 0.14	2.59 ± 0.52	Y	13.00 ± 0.39	48.10 ± 1.45	Y						
	12/1/2010	0.89 ± 0.13	3.31 ± 0.49	Y	29.80 ± 0.62	110.26 ± 2.31	Y						
	12/8/2010	0.97 ± 0.14	3.59 ± 0.51	Y	13.60 ± 0.45	50.32 ± 1.67	Y						
	12/15/2010	0.46 ± 0.11	1.68 ± 0.40	Y	16.60 ± 0.48	61.42 ± 1.76	Y						
	12/22/2010	0.57 ± 0.11	2.12 ± 0.39	Y	16.30 ± 0.46	60.31 ± 1.69	Y						
	12/29/2010	0.18 ± 0.12	0.66 ± 0.46	Y	18.50 ± 0.55	68.45 ± 2.02	Y						
	FAA TOWER	10/6/2010	2.56 ± 0.27	9.47 ± 1.01	Y	43.60 ± 0.97	161.32 ± 3.59	Y					
10/13/2010		1.24 ± 0.24	4.59 ± 0.90	Y	30.20 ± 0.90	111.74 ± 3.31	Y						
10/20/2010		2.42 ± 0.27	8.95 ± 0.99	Y	52.30 ± 1.06	193.51 ± 3.92	Y						
10/27/2010		0.91 ± 0.18	3.38 ± 0.67	Y	28.30 ± 0.79	104.71 ± 2.93	Y						
11/3/2010		1.13 ± 0.17	4.18 ± 0.64	Y	24.30 ± 0.67	89.91 ± 2.48	Y						
11/10/2010		1.27 ± 0.17	4.70 ± 0.64	Y	37.00 ± 0.78	136.90 ± 2.90	Y						
11/17/2010		±	±	±	±	±	±						
11/24/2010		±	±	±	±	±	±						
12/1/2010		±	±	±	±	±	±						
12/8/2010		±	±	±	±	±	±						
a	12/15/2010	0.40 ± 0.10	1.46 ± 0.37	Y	9.99 ± 0.39	36.96 ± 1.43	Y						
	12/22/2010	0.76 ± 0.13	2.82 ± 0.47	Y	17.40 ± 0.51	64.38 ± 1.90	Y						
	12/29/2010	0.05 ± 0.11	0.17 ± 0.41	Y	17.10 ± 0.52	63.27 ± 1.91	Y						
HOWE	10/6/2010	2.05 ± 0.24	7.59 ± 0.89	Y	40.40 ± 0.89	149.48 ± 3.30	Y						
	10/13/2010	1.40 ± 0.20	5.18 ± 0.74	Y	32.10 ± 0.74	118.77 ± 2.75	Y						

TABLE C-1. Weekly Gross Alpha and Gross Beta Concentrations in Air

Sampling Group and Location	Sampling Date	GROSS ALPHA					GROSS BETA				
		Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)		Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)		Result > 3s	Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)		Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)		Result > 3s
	10/20/2010	2.57	± 0.29	9.51	± 1.05	Y	49.10	± 1.08	181.67	± 4.00	Y
	10/27/2010	0.93	± 0.14	3.46	± 0.53	Y	29.90	± 0.64	110.63	± 2.38	Y
	11/3/2010	1.21	± 0.19	4.48	± 0.68	Y	26.00	± 0.72	96.20	± 2.68	Y
	11/10/2010	1.34	± 0.16	4.96	± 0.58	Y	34.30	± 0.68	126.91	± 2.50	Y
	11/17/2010	0.69	± 0.16	2.56	± 0.58	Y	23.70	± 0.64	87.69	± 2.38	Y
	11/24/2010	0.51	± 0.13	1.87	± 0.47	Y	13.50	± 0.39	49.95	± 1.45	Y
	12/1/2010	0.78	± 0.14	2.89	± 0.51	Y	34.70	± 0.70	128.39	± 2.60	Y
	12/8/2010	1.06	± 0.15	3.92	± 0.54	Y	22.50	± 0.55	83.25	± 2.05	Y
	12/15/2010	0.81	± 0.13	2.99	± 0.48	Y	18.60	± 0.50	68.82	± 1.86	Y
	12/22/2010	0.59	± 0.11	2.19	± 0.39	Y	19.10	± 0.48	70.67	± 1.78	Y
	12/29/2010	0.27	± 0.14	1.00	± 0.51	Y	25.30	± 0.64	93.61	± 2.38	Y
MONTEVIEW	10/6/2010	2.27	± 0.29	8.40	± 1.08	Y	41.90	± 1.05	155.03	± 3.89	Y
	10/13/2010	1.78	± 0.25	6.59	± 0.92	Y	31.70	± 0.85	117.29	± 3.14	Y
	10/20/2010	1.93	± 0.24	7.14	± 0.87	Y	34.90	± 0.88	129.13	± 3.24	Y
	10/27/2010	1.88	± 0.20	6.96	± 0.74	Y	27.80	± 0.68	102.86	± 2.51	Y
	11/3/2010	1.51	± 0.21	5.59	± 0.77	Y	24.60	± 0.74	91.02	± 2.73	Y
	11/10/2010	1.19	± 0.16	4.40	± 0.59	Y	33.00	± 0.70	122.10	± 2.60	Y
	11/17/2010	0.84	± 0.19	3.09	± 0.71	Y	21.40	± 0.71	79.18	± 2.61	Y
	11/24/2010	0.53	± 0.14	1.97	± 0.53	Y	12.70	± 0.41	46.99	± 1.53	Y
	12/1/2010	0.95	± 0.15	3.50	± 0.54	Y	30.20	± 0.67	111.74	± 2.49	Y
	12/8/2010	1.17	± 0.16	4.33	± 0.60	Y	25.20	± 0.62	93.24	± 2.28	Y
	12/15/2010	1.23	± 0.18	4.55	± 0.66	Y	17.20	± 0.58	63.64	± 2.14	Y
	12/22/2010	0.66	± 0.12	2.44	± 0.45	Y	18.70	± 0.52	69.19	± 1.94	Y
	12/29/2010	1.17	± 0.18	4.33	± 0.66	Y	26.50	± 0.64	98.05	± 2.37	Y
MUD LAKE	10/6/2010	1.71	± 0.26	6.33	± 0.96	Y	44.50	± 1.04	164.65	± 3.85	Y
	10/13/2010	1.08	± 0.21	4.00	± 0.77	Y	30.10	± 0.80	111.37	± 2.96	Y
	10/20/2010	2.50	± 0.29	9.25	± 1.07	Y	56.00	± 1.17	207.20	± 4.33	Y
	10/27/2010	1.16	± 0.17	4.29	± 0.63	Y	33.50	± 0.74	123.95	± 2.72	Y
	11/3/2010	1.74	± 0.20	6.44	± 0.74	Y	30.70	± 0.73	113.59	± 2.72	Y
	11/10/2010	1.47	± 0.18	5.44	± 0.65	Y	42.60	± 0.80	157.62	± 2.96	Y
	11/17/2010	0.98	± 0.17	3.61	± 0.63	Y	20.20	± 0.60	74.74	± 2.22	Y
	11/24/2010	0.52	± 0.14	1.92	± 0.50	Y	13.30	± 0.41	49.21	± 1.50	Y
	12/1/2010	1.34	± 0.17	4.96	± 0.61	Y	36.10	± 0.72	133.57	± 2.65	Y
	12/8/2010	1.14	± 0.16	4.22	± 0.60	Y	25.60	± 0.62	94.72	± 2.29	Y
	12/15/2010	1.24	± 0.16	4.59	± 0.59	Y	24.00	± 0.59	88.80	± 2.18	Y
	12/22/2010	0.57	± 0.11	2.09	± 0.42	Y	20.30	± 0.53	75.11	± 1.96	Y
	12/29/2010	0.70	± 0.16	2.58	± 0.58	Y	33.00	± 0.70	122.10	± 2.58	Y
DISTANT											
BLACKFOOT CMS	10/6/2010	2.66	± 0.29	9.84	± 1.08	Y	47.70	± 1.05	176.49	± 3.89	Y
	10/13/2010	1.47	± 0.23	5.44	± 0.86	Y	29.10	± 0.81	107.67	± 3.01	Y
	10/20/2010	2.41	± 0.27	8.92	± 1.00	Y	52.10	± 1.08	192.77	± 4.00	Y
	10/27/2010	1.58	± 0.20	5.85	± 0.73	Y	27.40	± 0.71	101.38	± 2.62	Y
	11/3/2010	0.93	± 0.15	3.43	± 0.56	Y	25.10	± 0.64	92.87	± 2.38	Y
	11/10/2010	1.61	± 0.19	5.96	± 0.69	Y	37.40	± 0.78	138.38	± 2.87	Y
	11/17/2010	0.56	± 0.15	2.09	± 0.56	Y	16.20	± 0.56	59.94	± 2.09	Y
	11/24/2010	0.42	± 0.13	1.55	± 0.49	Y	12.00	± 0.40	44.40	± 1.46	Y
	12/1/2010	1.11	± 0.17	4.11	± 0.63	Y	33.70	± 0.77	124.69	± 2.85	Y
	12/8/2010	0.91	± 0.17	3.38	± 0.64	Y	28.00	± 0.73	103.60	± 2.69	Y
	12/15/2010	0.57	± 0.14	2.12	± 0.50	Y	19.60	± 0.58	72.52	± 2.14	Y
	12/22/2010	1.08	± 0.15	4.00	± 0.56	Y	18.50	± 0.55	68.45	± 2.02	Y
	12/29/2010	0.44	± 0.14	1.62	± 0.52	Y	17.00	± 0.54	62.90	± 1.99	Y
CRATERS OF THE MOON	10/6/2010	1.99	± 0.26	7.36	± 0.97	Y	37.60	± 0.95	139.12	± 3.51	Y
	10/13/2010	1.42	± 0.24	5.25	± 0.89	Y	29.20	± 0.85	108.04	± 3.15	Y
	10/20/2010	1.90	± 0.29	7.03	± 1.07	Y	47.10	± 1.19	174.27	± 4.40	Y
	10/27/2010	1.21	± 0.19	4.48	± 0.70	Y	21.20	± 0.69	78.44	± 2.54	Y
	11/3/2010	0.83	± 0.15	3.06	± 0.56	Y	21.20	± 0.62	78.44	± 2.29	Y



TABLE C-1. Weekly Gross Alpha and Gross Beta Concentrations in Air

Sampling Group and Location	Sampling Date	GROSS ALPHA			GROSS BETA			
		Result ± 1s Uncertainty (x 10 <sup>15</sup> µCi/mL)	Result ± 1s Uncertainty (x 10 <sup>11</sup> Bq/mL)	Result > 3s	Result ± 1s Uncertainty (x 10 <sup>15</sup> µCi/mL)	Result ± 1s Uncertainty (x 10 <sup>11</sup> Bq/mL)	Result > 3s	
	11/10/2010	1.11 ± 0.17	4.11 ± 0.63	Y	30.80 ± 0.76	113.96 ± 2.80	Y	
	11/17/2010	0.50 ± 0.15	1.84 ± 0.56	Y	20.20 ± 0.62	74.74 ± 2.28	Y	
	11/24/2010	0.45 ± 0.15	1.67 ± 0.55	Y	10.50 ± 0.41	38.85 ± 1.51	Y	
	12/1/2010	0.83 ± 0.15	3.06 ± 0.54	Y	28.20 ± 0.68	104.34 ± 2.52	Y	
	12/8/2010	0.73 ± 0.16	2.72 ± 0.58	Y	19.10 ± 0.61	70.67 ± 2.26	Y	
	12/15/2010	0.61 ± 0.14	2.26 ± 0.52	Y	15.30 ± 0.54	56.61 ± 2.00	Y	
	12/22/2010	0.48 ± 0.12	1.79 ± 0.44	Y	16.80 ± 0.55	62.16 ± 2.04	Y	
	12/29/2010	0.31 ± 0.14	1.14 ± 0.52	Y	15.40 ± 0.54	56.98 ± 2.00	Y	
	DUBOIS	10/6/2010	2.41 ± 0.27	8.92 ± 1.01	Y	39.20 ± 0.94	145.04 ± 3.49	Y
		10/13/2010	1.01 ± 0.21	3.74 ± 0.76	Y	31.80 ± 0.81	117.66 ± 3.01	Y
		10/20/2010	2.10 ± 0.26	7.77 ± 0.97	Y	48.80 ± 1.06	180.56 ± 3.92	Y
		10/27/2010	1.48 ± 0.19	5.48 ± 0.70	Y	26.30 ± 0.68	97.31 ± 2.53	Y
	11/3/2010	1.04 ± 0.16	3.85 ± 0.60	Y	23.60 ± 0.65	87.32 ± 2.41	Y	
	11/10/2010	1.37 ± 0.17	5.07 ± 0.62	Y	30.90 ± 0.69	114.33 ± 2.56	Y	
	11/17/2010	0.62 ± 0.16	2.30 ± 0.59	Y	19.20 ± 0.61	71.04 ± 2.27	Y	
	11/24/2010	0.68 ± 0.16	2.53 ± 0.58	Y	12.30 ± 0.42	45.51 ± 1.57	Y	
	12/1/2010	0.88 ± 0.15	3.27 ± 0.56	Y	30.40 ± 0.71	112.48 ± 2.61	Y	
	12/8/2010	0.58 ± 0.14	2.14 ± 0.52	Y	22.00 ± 0.61	81.40 ± 2.27	Y	
	12/15/2010	0.60 ± 0.15	2.21 ± 0.54	Y	22.10 ± 0.64	81.77 ± 2.36	Y	
	12/22/2010	0.68 ± 0.12	2.50 ± 0.46	Y	18.60 ± 0.53	68.82 ± 1.96	Y	
	12/29/2010	0.47 ± 0.15	1.72 ± 0.57	Y	20.60 ± 0.61	76.22 ± 2.25	Y	
QA-2 (DUBOIS)	10/6/2010	2.36 ± 0.27	8.73 ± 1.00	Y	39.80 ± 0.95	147.26 ± 3.50	Y	
	10/13/2010	1.43 ± 0.25	5.29 ± 0.91	Y	30.50 ± 0.88	112.85 ± 3.26	Y	
	10/20/2010	2.56 ± 0.27	9.47 ± 1.01	Y	47.80 ± 1.03	176.86 ± 3.81	Y	
	10/27/2010	0.92 ± 0.16	3.40 ± 0.58	Y	27.50 ± 0.68	101.75 ± 2.52	Y	
	11/3/2010	1.25 ± 0.16	4.63 ± 0.61	Y	24.20 ± 0.62	89.54 ± 2.29	Y	
	11/10/2010	2.01 ± 0.22	7.44 ± 0.81	Y	33.50 ± 0.80	123.95 ± 2.97	Y	
	11/17/2010	0.64 ± 0.15	2.38 ± 0.57	Y	19.30 ± 0.59	71.41 ± 2.19	Y	
	11/24/2010	0.64 ± 0.15	2.38 ± 0.57	Y	13.10 ± 0.43	48.47 ± 1.59	Y	
	12/1/2010	0.99 ± 0.15	3.64 ± 0.56	Y	27.00 ± 0.66	99.90 ± 2.44	Y	
	12/8/2010	0.80 ± 0.16	2.97 ± 0.58	Y	21.20 ± 0.63	78.44 ± 2.32	Y	
	12/15/2010	1.11 ± 0.17	4.11 ± 0.63	Y	21.10 ± 0.61	78.07 ± 2.26	Y	
	12/22/2010	0.40 ± 0.11	1.47 ± 0.39	Y	16.40 ± 0.51	60.68 ± 1.89	Y	
	12/29/2010	0.52 ± 0.15	1.93 ± 0.56	Y	20.70 ± 0.60	76.59 ± 2.21	Y	
IDAHO FALLS	10/6/2010	2.72 ± 0.33	10.06 ± 1.22	Y	45.20 ± 1.15	167.24 ± 4.26	Y	
	10/13/2010	1.63 ± 0.26	6.03 ± 0.97	Y	31.20 ± 0.91	115.44 ± 3.36	Y	
	10/20/2010	3.05 ± 0.32	11.29 ± 1.17	Y	51.30 ± 1.14	189.81 ± 4.22	Y	
	10/27/2010	1.42 ± 0.21	5.25 ± 0.76	Y	29.60 ± 0.79	109.52 ± 2.92	Y	
	11/3/2010	1.27 ± 0.19	4.70 ± 0.70	Y	27.00 ± 0.74	99.90 ± 2.75	Y	
	11/10/2010	1.45 ± 0.19	5.37 ± 0.70	Y	35.20 ± 0.80	130.24 ± 2.97	Y	
	11/17/2010	0.49 ± 0.17	1.82 ± 0.63		18.60 ± 0.67	68.82 ± 2.48	Y	
	11/24/2010	0.32 ± 0.15	1.18 ± 0.56		11.30 ± 0.44	41.81 ± 1.62	Y	
	12/1/2010	0.98 ± 0.18	3.61 ± 0.65	Y	36.60 ± 0.85	135.42 ± 3.13	Y	
	12/8/2010	0.59 ± 0.16	2.18 ± 0.60	Y	25.30 ± 0.73	93.61 ± 2.69	Y	
	12/15/2010	0.72 ± 0.15	2.65 ± 0.57	Y	18.00 ± 0.60	66.60 ± 2.21	Y	
	12/22/2010	0.51 ± 0.12	1.89 ± 0.46	Y	19.40 ± 0.58	71.78 ± 2.14	Y	
	12/29/2010	0.42 ± 0.16	1.55 ± 0.58		19.30 ± 0.62	71.41 ± 2.28	Y	
JACKSON	10/6/2010	3.45 ± 0.36	12.77 ± 1.34	Y	47.50 ± 1.19	175.75 ± 4.40	Y	
	10/13/2010	1.45 ± 0.27	5.37 ± 1.01	Y	29.40 ± 0.95	108.78 ± 3.52	Y	
	10/20/2010	2.13 ± 0.28	7.88 ± 1.04	Y	40.70 ± 1.07	150.59 ± 3.96	Y	
	10/27/2010	1.00 ± 0.17	3.70 ± 0.64	Y	26.00 ± 0.72	96.20 ± 2.65	Y	
	11/3/2010	1.03 ± 0.17	3.81 ± 0.64	Y	23.40 ± 0.69	86.58 ± 2.54	Y	
	11/10/2010	1.51 ± 0.20	5.59 ± 0.75	Y	35.60 ± 0.85	131.72 ± 3.13	Y	
	11/17/2010	0.03 ± 0.13	0.11 ± 0.48		12.20 ± 0.56	45.14 ± 2.08	Y	
	11/23/2010	0.95 ± 0.21	3.51 ± 0.78	Y	12.80 ± 0.52	47.36 ± 1.91	Y	
	12/1/2010	1.20 ± 0.17	4.44 ± 0.61	Y	26.50 ± 0.66	98.05 ± 2.45	Y	

TABLE C-1. Weekly Gross Alpha and Gross Beta Concentrations in Air

Sampling Group and Location	Sampling Date	GROSS ALPHA			GROSS BETA		
		Result ± 1s Uncertainty (x 10 <sup>15</sup> µCi/mL)	Result ± 1s Uncertainty (x 10 <sup>11</sup> Bq/mL)	Result > 3s	Result ± 1s Uncertainty (x 10 <sup>15</sup> µCi/mL)	Result ± 1s Uncertainty (x 10 <sup>11</sup> Bq/mL)	Result > 3s
	12/8/2010	1.10 ± 0.19	4.07 ± 0.70	Y	24.70 ± 0.71	91.39 ± 2.63	Y
	12/15/2010	0.43 ± 0.14	1.58 ± 0.50	Y	12.00 ± 0.53	44.40 ± 1.95	Y
	12/22/2010	1.27 ± 0.18	4.70 ± 0.66	Y	22.30 ± 0.66	82.51 ± 2.45	Y
	12/29/2010	1.08 ± 0.20	4.00 ± 0.74	Y	25.30 ± 0.71	93.61 ± 2.62	Y
REXBURG CMS	10/6/2010	2.67 ± 0.30	9.88 ± 1.11	Y	45.90 ± 1.06	169.83 ± 3.92	Y
	10/13/2010	1.81 ± 0.25	6.70 ± 0.94	Y	30.70 ± 0.86	113.59 ± 3.17	Y
	10/20/2010	2.65 ± 0.28	9.81 ± 1.05	Y	47.80 ± 1.05	176.86 ± 3.89	Y
	10/27/2010	1.71 ± 0.20	6.33 ± 0.74	Y	29.00 ± 0.71	107.30 ± 2.64	Y
	11/3/2010	1.01 ± 0.17	3.74 ± 0.61	Y	23.90 ± 0.67	88.43 ± 2.47	Y
	11/10/2010	1.50 ± 0.18	5.55 ± 0.67	Y	33.20 ± 0.74	122.84 ± 2.72	Y
	11/17/2010	0.56 ± 0.15	2.08 ± 0.54	Y	15.90 ± 0.55	58.83 ± 2.03	Y
	11/24/2010	0.70 ± 0.15	2.59 ± 0.54	Y	10.50 ± 0.38	38.85 ± 1.39	Y
	12/1/2010	1.13 ± 0.16	4.18 ± 0.59	Y	30.30 ± 0.69	112.11 ± 2.54	Y
	12/8/2010	1.03 ± 0.18	3.81 ± 0.67	Y	25.70 ± 0.71	95.09 ± 2.63	Y
	12/15/2010	1.01 ± 0.16	3.74 ± 0.60	Y	20.00 ± 0.60	74.00 ± 2.21	Y
	12/22/2010	0.58 ± 0.12	2.16 ± 0.46	Y	18.80 ± 0.56	69.56 ± 2.06	Y
	12/29/2010	0.74 ± 0.16	2.73 ± 0.59	Y	18.50 ± 0.56	68.45 ± 2.08	Y
<b>INL SITE</b>							
EFS	10/6/2010	2.16 ± 0.25	7.99 ± 0.92	Y	40.10 ± 0.90	148.37 ± 3.33	Y
	10/13/2010	1.13 ± 0.21	4.18 ± 0.76	Y	29.20 ± 0.77	108.04 ± 2.86	Y
	10/20/2010	1.94 ± 0.24	7.18 ± 0.90	Y	49.10 ± 1.02	181.67 ± 3.77	Y
	10/27/2010	1.12 ± 0.19	4.14 ± 0.68	Y	29.40 ± 0.77	108.78 ± 2.85	Y
	11/3/2010	0.90 ± 0.14	3.31 ± 0.52	Y	25.80 ± 0.61	95.46 ± 2.24	Y
	11/10/2010	1.33 ± 0.17	4.92 ± 0.63	Y	35.80 ± 0.74	132.46 ± 2.74	Y
	11/17/2010	0.56 ± 0.13	2.07 ± 0.48	Y	18.00 ± 0.51	66.60 ± 1.90	Y
	11/24/2010	0.48 ± 0.12	1.77 ± 0.45	Y	12.00 ± 0.36	44.40 ± 1.34	Y
	12/1/2010	1.02 ± 0.14	3.77 ± 0.51	Y	32.20 ± 0.63	119.14 ± 2.34	Y
	12/8/2010	1.09 ± 0.17	4.03 ± 0.61	Y	12.20 ± 0.50	45.14 ± 1.85	Y
	12/15/2010	0.93 ± 0.14	3.44 ± 0.50	Y	15.00 ± 0.46	55.50 ± 1.71	Y
	12/22/2010	1.30 ± 0.15	4.81 ± 0.55	Y	25.20 ± 0.57	93.24 ± 2.09	Y
	12/29/2010	0.53 ± 0.13	1.94 ± 0.48	Y	25.50 ± 0.57	94.35 ± 2.12	Y
MAIN GATE	10/6/2010	1.99 ± 0.28	7.36 ± 1.04	Y	43.50 ± 1.06	160.95 ± 3.92	Y
	10/13/2010	1.11 ± 0.23	4.11 ± 0.85	Y	33.90 ± 0.90	125.43 ± 3.32	Y
	10/20/2010	2.55 ± 0.31	9.44 ± 1.14	Y	52.70 ± 1.20	194.99 ± 4.44	Y
	10/27/2010	0.99 ± 0.18	3.67 ± 0.65	Y	27.40 ± 0.75	101.38 ± 2.76	Y
	11/3/2010	1.35 ± 0.19	5.00 ± 0.72	Y	27.40 ± 0.74	101.38 ± 2.75	Y
	11/10/2010	1.43 ± 0.20	5.29 ± 0.73	Y	38.40 ± 0.86	142.08 ± 3.19	Y
	11/17/2010	0.65 ± 0.17	2.40 ± 0.62	Y	19.20 ± 0.64	71.04 ± 2.35	Y
	11/24/2010	0.41 ± 0.13	1.53 ± 0.50	Y	12.90 ± 0.41	47.73 ± 1.52	Y
	12/1/2010	0.87 ± 0.15	3.20 ± 0.54	Y	35.00 ± 0.72	129.50 ± 2.67	Y
	12/8/2010	1.18 ± 0.17	4.37 ± 0.63	Y	16.20 ± 0.55	59.94 ± 2.03	Y
	12/15/2010	0.67 ± 0.14	2.49 ± 0.50	Y	12.40 ± 0.48	45.88 ± 1.77	Y
	12/22/2010	1.01 ± 0.15	3.74 ± 0.56	Y	21.80 ± 0.59	80.66 ± 2.20	Y
	12/29/2010	0.35 ± 0.24	1.28 ± 0.90	Y	24.10 ± 0.94	89.17 ± 3.46	Y
VAN BUREN GATE	10/6/2010	2.39 ± 0.27	8.84 ± 0.98	Y	42.90 ± 0.95	158.73 ± 3.52	Y
	10/13/2010	1.33 ± 0.23	4.92 ± 0.84	Y	31.00 ± 0.83	114.70 ± 3.08	Y
	10/20/2010	2.30 ± 0.28	8.51 ± 1.03	Y	51.30 ± 1.12	189.81 ± 4.14	Y
	10/27/2010	1.25 ± 0.16	4.63 ± 0.61	Y	27.70 ± 0.65	102.49 ± 2.40	Y
	11/3/2010	0.64 ± 0.14	2.38 ± 0.51	Y	27.40 ± 0.68	101.38 ± 2.51	Y
	11/10/2010	1.43 ± 0.24	5.29 ± 0.90	Y	55.30 ± 1.21	204.61 ± 4.48	Y
	11/17/2010	0.40 ± 0.15	1.46 ± 0.54		16.10 ± 0.58	59.57 ± 2.16	Y
	11/24/2010	0.34 ± 0.12	1.27 ± 0.44		9.44 ± 0.34	34.93 ± 1.27	Y
	12/1/2010	1.00 ± 0.15	3.69 ± 0.54	Y	30.20 ± 0.66	111.74 ± 2.45	Y
	12/8/2010	1.01 ± 0.16	3.74 ± 0.57	Y	19.00 ± 0.56	70.30 ± 2.06	Y
	12/15/2010	0.68 ± 0.13	2.52 ± 0.48	Y	21.40 ± 0.56	79.18 ± 2.06	Y
	12/22/2010	0.89 ± 0.14	3.30 ± 0.51	Y	22.10 ± 0.58	81.77 ± 2.13	Y

TABLE C-1. Weekly Gross Alpha and Gross Beta Concentrations in Air

Sampling Group and Location	Sampling Date	GROSS ALPHA				GROSS BETA			
		Result ± 1s Uncertainty (x 10 <sup>15</sup> µCi/mL)		Result ± 1s Uncertainty (x 10 <sup>11</sup> Bq/mL)		Result ± 1s Uncertainty (x 10 <sup>15</sup> µCi/mL)		Result ± 1s Uncertainty (x 10 <sup>11</sup> Bq/mL)	
	12/29/2010	0.35 ± 0.13	1.30 ± 0.48		22.70 ± 0.58	83.99 ± 2.13	Y		
QA-1 (VAN BUREN)	10/6/2010	2.30 ± 0.28	8.51 ± 1.02	Y	41.80 ± 0.99	154.66 ± 3.66	Y		
	10/13/2010	1.51 ± 0.22	5.59 ± 0.81	Y	34.50 ± 0.81	127.65 ± 3.01	Y		
	10/20/2010	2.74 ± 0.28	10.14 ± 1.04	Y	54.40 ± 1.08	201.28 ± 4.00	Y		
	10/27/2010	1.36 ± 0.18	5.03 ± 0.65	Y	29.00 ± 0.69	107.30 ± 2.54	Y		
	11/3/2010	0.84 ± 0.14	3.11 ± 0.53	Y	25.90 ± 0.64	95.83 ± 2.35	Y		
	11/10/2010	1.83 ± 0.26	6.77 ± 0.98	Y	52.30 ± 1.19	193.51 ± 4.40	Y		
	11/17/2010	0.71 ± 0.17	2.63 ± 0.61	Y	16.50 ± 0.59	61.05 ± 2.18	Y		
	11/24/2010	0.19 ± 0.12	0.70 ± 0.44		9.34 ± 0.36	34.56 ± 1.33	Y		
	12/1/2010	0.91 ± 0.14	3.38 ± 0.50	Y	30.80 ± 0.63	113.96 ± 2.34	Y		
	12/8/2010	1.06 ± 0.15	3.92 ± 0.54	Y	19.30 ± 0.53	71.41 ± 1.94	Y		
	12/15/2010	0.82 ± 0.13	3.04 ± 0.48	Y	22.80 ± 0.54	84.36 ± 2.01	Y		
	12/22/2010	0.90 ± 0.13	3.33 ± 0.48	Y	22.00 ± 0.55	81.40 ± 2.02	Y		
	12/29/2010	0.40 ± 0.13	1.47 ± 0.47	Y	25.40 ± 0.59	93.98 ± 2.18	Y		
a. Invalid Sample Result									

TABLE C-2. Weekly Iodine-131 Activity in Air.

Sampling Group and Location	Sampling Date	Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)			Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)			Result > 3s	
<b>BOUNDARY</b>									
ARCO	10/06/2010	-0.70	±	1.14	-2.59	±	4.23		
	10/13/2010	-0.44	±	1.17	-1.64	±	4.31		
	10/20/2010	-1.04	±	1.31	-3.84	±	4.86		
	10/27/2010	-1.28	±	1.21	-4.74	±	4.49		
	11/03/2010	1.27	±	1.25	4.69	±	4.61		
	11/10/2010	0.89	±	1.23	3.31	±	4.55		
	11/17/2010	-1.66	±	1.11	-6.16	±	4.13		
	11/24/2010	2.65	±	1.31	9.79	±	4.84		
	12/01/2010	-1.87	±	1.19	-6.94	±	4.39		
	12/08/2010	0.56	±	1.32	2.08	±	4.89		
	12/15/2010	0.00	±	1.24	0.01	±	4.60		
	12/22/2010	-0.35	±	1.19	-1.30	±	4.39		
	12/29/2010	-1.50	±	1.14	-5.56	±	4.20		
ATOMIC CITY	10/06/2010	-0.62	±	1.01	-2.29	±	3.74		
	10/13/2010	-0.38	±	1.00	-1.41	±	3.69		
	10/20/2010	-1.06	±	1.34	-3.92	±	4.96		
	10/27/2010	-1.05	±	1.00	-3.89	±	3.69		
	11/03/2010	0.96	±	0.95	3.56	±	3.50		
	11/10/2010	0.72	±	0.99	2.67	±	3.67		
	11/17/2010	-1.33	±	0.89	-4.91	±	3.29		
	11/24/2010	1.80	±	0.89	6.65	±	3.28		
	12/01/2010	-1.37	±	0.87	-5.06	±	3.21		
	12/08/2010	0.35	±	0.83	1.31	±	3.08		
	12/15/2010	0.00	±	0.87	0.01	±	3.22		
	12/22/2010	-0.23	±	0.77	-0.84	±	2.84		
	12/29/2010	-1.31	±	0.99	-4.84	±	3.65		
BLUE DOME	10/06/2010	-0.20	±	1.01	-0.74	±	3.74		
	10/13/2010	0.92	±	0.97	3.41	±	3.58		
	10/20/2010	-0.23	±	1.15	-0.85	±	4.27		
	10/27/2010	0.51	±	0.99	1.90	±	3.68		
	11/03/2010	-0.90	±	1.96	-3.33	±	7.25		
	11/10/2010	-0.71	±	0.93	-2.62	±	3.46		
	11/17/2010	0.00	±	1.04	-0.02	±	3.84		
	11/24/2010	0.00	±	0.90	0.01	±	3.32		
	12/01/2010	-0.49	±	0.84	-1.81	±	3.11		
	12/08/2010	0.38	±	0.80	1.42	±	2.97		
	12/15/2010	0.89	±	0.81	3.28	±	2.99		
	12/22/2010	1.06	±	0.77	3.91	±	2.84		
	12/29/2010	-1.38	±	0.95	-5.10	±	3.50		
FAA TOWER	10/06/2010	-0.19	±	0.98	-0.72	±	3.63		
	10/13/2010	1.07	±	1.12	3.96	±	4.16		
	10/20/2010	-0.22	±	1.09	-0.80	±	4.02		
	10/27/2010	0.67	±	1.30	2.49	±	4.82		
	11/03/2010	-0.51	±	1.12	-1.90	±	4.13		
	11/10/2010	-0.79	±	1.03	-2.90	±	3.83		
	a	11/17/2010	-0.04	±	8.35	-0.14	±	30.89	
	a	11/24/2010		±			±		
	a	12/01/2010		±			±		
	a	12/08/2010		±			±		
		12/15/2010	0.87	±	0.79	3.20	±	2.92	

**TABLE C-2. Weekly Iodine-131 Activity in Air.**

Sampling Group and Location	Sampling Date	Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)			Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)			Result > 3s
<b>BOUNDARY</b>								
	12/22/2010	1.23	±	0.89	4.55	±	3.30	
	12/29/2010	-1.32	±	0.91	-4.90	±	3.36	
HOWE	10/06/2010	-0.18	±	0.90	-0.66	±	3.33	
	10/13/2010	0.77	±	0.81	2.84	±	2.98	
	10/20/2010	-0.23	±	1.17	-0.86	±	4.34	
	10/27/2010	0.45	±	0.87	1.67	±	3.23	
	11/03/2010	-0.56	±	1.21	-2.05	±	4.47	
	11/10/2010	-0.64	±	0.84	-2.36	±	3.11	
	11/17/2010	0.00	±	0.99	-0.02	±	3.65	
	11/24/2010	0.00	±	0.87	0.00	±	3.21	
	12/01/2010	-0.54	±	0.93	-2.00	±	3.43	
	12/08/2010	0.39	±	0.81	1.44	±	3.01	
	12/15/2010	0.90	±	0.83	3.35	±	3.06	
	12/22/2010	1.04	±	0.75	3.84	±	2.79	
	12/29/2010	-1.47	±	1.01	-5.45	±	3.74	
MONTEVIEW	10/06/2010	-0.23	±	1.15	-0.85	±	4.27	
	10/13/2010	0.96	±	1.00	3.54	±	3.71	
	10/20/2010	-0.20	±	1.03	-0.76	±	3.82	
	10/27/2010	0.52	±	1.01	1.92	±	3.72	
	11/03/2010	-0.60	±	1.30	-2.21	±	4.80	
	11/10/2010	-0.71	±	0.94	-2.63	±	3.47	
	11/17/2010	-0.01	±	1.22	-0.02	±	4.53	
	11/24/2010	0.00	±	1.00	0.01	±	3.70	
	12/01/2010	-0.56	±	0.95	-2.06	±	3.53	
	12/08/2010	0.43	±	0.91	1.60	±	3.35	
	12/15/2010	1.19	±	1.09	4.42	±	4.04	
	12/22/2010	1.22	±	0.88	4.50	±	3.26	
	12/29/2010	-1.41	±	0.97	-5.21	±	3.58	
MUD LAKE	10/06/2010	-0.22	±	1.09	-0.80	±	4.02	
	10/13/2010	0.90	±	0.95	3.33	±	3.50	
	10/20/2010	-0.24	±	1.21	-0.89	±	4.49	
	10/27/2010	0.52	±	1.01	1.93	±	3.74	
	11/03/2010	-0.51	±	1.10	-1.87	±	4.07	
	11/10/2010	-0.73	±	0.96	-2.70	±	3.55	
	11/17/2010	0.00	±	0.97	-0.02	±	3.60	
	11/24/2010	0.00	±	0.93	0.01	±	3.45	
	12/01/2010	-0.54	±	0.92	-1.99	±	3.42	
	12/08/2010	0.43	±	0.90	1.59	±	3.34	
	12/15/2010	0.98	±	0.90	3.64	±	3.32	
	12/22/2010	1.17	±	0.85	4.33	±	3.14	
	12/29/2010	-1.39	±	0.95	-5.14	±	3.53	
<b>DISTANT</b>								
BLACKFOOT CMS	10/06/2010	-0.69	±	1.13	-2.55	±	4.18	
	10/13/2010	-0.40	±	1.06	-1.49	±	3.91	
	10/20/2010	-0.95	±	1.20	-3.52	±	4.45	
	10/27/2010	-1.24	±	1.18	-4.58	±	4.35	
	11/03/2010	1.13	±	1.11	4.18	±	4.12	
	11/10/2010	0.81	±	1.12	3.00	±	4.13	
	11/17/2010	-1.62	±	1.09	-5.99	±	4.02	
	11/24/2010	2.19	±	1.08	8.11	±	4.01	

**TABLE C-2. Weekly Iodine-131 Activity in Air.**

Sampling Group and Location	Sampling Date	Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)			Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)			Result > 3s
<b>BOUNDARY</b>								
	12/01/2010	-1.83	±	1.16	-6.79	±	4.30	
	12/08/2010	0.50	±	1.18	1.86	±	4.37	
	12/15/2010	0.00	±	1.08	0.01	±	4.00	
	12/22/2010	-0.30	±	1.01	-1.10	±	3.73	
	12/29/2010	-1.32	±	1.00	-4.89	±	3.69	
<b>CRATERS</b>								
	10/06/2010	-0.68	±	1.11	-2.52	±	4.12	
	10/13/2010	-0.43	±	1.13	-1.59	±	4.19	
	10/20/2010	-1.21	±	1.53	-4.47	±	5.66	
	10/27/2010	-1.39	±	1.32	-5.15	±	4.88	
	11/03/2010	1.19	±	1.17	4.41	±	4.34	
	11/10/2010	0.90	±	1.24	3.33	±	4.58	
	11/17/2010	-1.65	±	1.10	-6.09	±	4.08	
	11/24/2010	2.50	±	1.23	9.24	±	4.57	
	12/01/2010	-1.69	±	1.07	-6.26	±	3.97	
	12/08/2010	0.48	±	1.13	1.78	±	4.19	
	12/15/2010	0.00	±	1.13	0.01	±	4.17	
	12/22/2010	-0.32	±	1.09	-1.19	±	4.03	
	12/29/2010	-1.42	±	1.07	-5.24	±	3.96	
<b>DUBOIS</b>								
	10/06/2010	-0.20	±	1.01	-0.74	±	3.72	
	10/13/2010	0.89	±	0.94	3.30	±	3.47	
	10/20/2010	-0.23	±	1.14	-0.83	±	4.20	
	10/27/2010	0.55	±	1.07	2.04	±	3.94	
	11/03/2010	-0.50	±	1.08	-1.84	±	4.00	
	11/10/2010	-0.72	±	0.95	-2.68	±	3.53	
	11/17/2010	0.00	±	1.05	-0.02	±	3.88	
	11/24/2010	0.00	±	1.06	0.01	±	3.91	
	12/01/2010	-0.60	±	1.03	-2.22	±	3.81	
	12/08/2010	0.47	±	0.99	1.74	±	3.66	
	12/15/2010	1.19	±	1.09	4.42	±	4.03	
	12/22/2010	1.24	±	0.90	4.59	±	3.33	
	12/29/2010	-1.54	±	1.05	-5.68	±	3.90	
<b>QA-2</b>								
	10/06/2010	-0.20	±	1.00	-0.74	±	3.71	
	10/13/2010	1.04	±	1.09	3.84	±	4.04	
	10/20/2010	-0.22	±	1.10	-0.81	±	4.08	
	10/27/2010	0.53	±	1.03	1.97	±	3.81	
	11/03/2010	-0.45	±	0.97	-1.65	±	3.60	
	11/10/2010	-0.88	±	1.16	-3.25	±	4.28	
	11/17/2010	0.00	±	0.99	-0.02	±	3.65	
	11/24/2010	0.00	±	1.04	0.01	±	3.85	
	12/01/2010	-0.58	±	1.00	-2.15	±	3.69	
	12/08/2010	0.50	±	1.04	1.84	±	3.85	
	12/15/2010	1.15	±	1.05	4.26	±	3.89	
	12/22/2010	1.27	±	0.93	4.72	±	3.42	
	12/29/2010	-1.48	±	1.01	-5.47	±	3.75	
<b>IDAHO FALLS</b>								
	10/06/2010	-0.25	±	1.26	-0.93	±	4.67	
	10/13/2010	1.08	±	1.13	3.98	±	4.18	
	10/20/2010	-0.25	±	1.25	-0.91	±	4.61	
	10/27/2010	0.65	±	1.25	2.40	±	4.64	
	11/03/2010	-0.57	±	1.23	-2.09	±	4.56	
	11/10/2010	-0.85	±	1.12	-3.15	±	4.14	

**TABLE C-2. Weekly Iodine-131 Activity in Air.**

Sampling Group and Location	Sampling Date	Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)			Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)			Result > 3s
<b>BOUNDARY</b>								
	11/17/2010	-0.01	±	1.22	-0.02	±	4.52	
	11/24/2010	0.00	±	1.18	0.01	±	4.38	
	12/01/2010	-0.72	±	1.23	-2.65	±	4.55	
	12/08/2010	0.57	±	1.19	2.10	±	4.41	
	12/15/2010	1.23	±	1.12	4.54	±	4.15	
	12/22/2010	1.40	±	1.02	5.19	±	3.77	
	12/29/2010	-1.63	±	1.12	-6.02	±	4.13	
<b>JACKSON</b>								
	10/06/2010	-0.85	±	1.39	-3.13	±	5.13	
	10/13/2010	-0.51	±	1.34	-1.88	±	4.95	
	10/20/2010	-1.11	±	1.40	-4.10	±	5.19	
	10/27/2010	-1.31	±	1.24	-4.85	±	4.60	
	11/03/2010	1.32	±	1.30	4.90	±	4.82	
	11/10/2010	0.98	±	1.35	3.63	±	4.99	
	11/17/2010	-1.86	±	1.25	-6.90	±	4.62	
	11/23/2010	3.99	±	2.73	14.78	±	10.10	
	12/01/2010	-1.68	±	1.06	-6.20	±	3.93	
	12/08/2010	0.53	±	1.24	1.95	±	4.60	
	12/15/2010	0.00	±	1.24	0.01	±	4.59	
	12/22/2010	-0.36	±	1.22	-1.33	±	4.52	
	12/29/2010	-1.61	±	1.21	-5.95	±	4.49	
<b>REXBURG CMS</b>								
	10/06/2010	-0.22	±	1.11	-0.81	±	4.09	
	10/13/2010	0.99	±	1.04	3.65	±	3.83	
	10/20/2010	-0.23	±	1.14	-0.84	±	4.22	
	10/27/2010	0.55	±	1.07	2.04	±	3.96	
	11/03/2010	-0.51	±	1.12	-1.90	±	4.14	
	11/10/2010	-0.76	±	1.00	-2.82	±	3.72	
	11/17/2010	0.00	±	0.97	-0.02	±	3.61	
	11/24/2010	0.00	±	0.96	0.01	±	3.56	
	12/01/2010	-0.57	±	0.99	-2.12	±	3.64	
	12/08/2010	0.54	±	1.14	2.01	±	4.21	
	12/15/2010	1.14	±	1.04	4.23	±	3.86	
	12/22/2010	1.34	±	0.98	4.98	±	3.61	
	12/29/2010	-1.44	±	0.99	-5.33	±	3.66	
<b>INL SITE</b>								
<b>EFS</b>								
	10/06/2010	-0.60	±	0.98	-2.22	±	3.64	
	10/13/2010	-0.37	±	0.97	-1.37	±	3.61	
	10/20/2010	-0.91	±	1.15	-3.35	±	4.24	
	10/27/2010	-1.36	±	1.29	-5.04	±	4.78	
	11/03/2010	1.00	±	0.98	3.70	±	3.64	
	11/10/2010	0.78	±	1.07	2.87	±	3.94	
	11/17/2010	-1.32	±	0.88	-4.87	±	3.26	
	11/24/2010	1.88	±	0.93	6.95	±	3.43	
	12/01/2010	-1.33	±	0.84	-4.93	±	3.12	
	12/08/2010	0.45	±	1.06	1.67	±	3.94	
	12/15/2010	0.00	±	0.88	0.01	±	3.27	
	12/22/2010	-0.25	±	0.85	-0.93	±	3.16	
	12/29/2010	-1.12	±	0.85	-4.14	±	3.13	
<b>MAIN GATE</b>								
	10/06/2010	-0.75	±	1.22	-2.76	±	4.52	
	10/13/2010	-0.43	±	1.13	-1.60	±	4.20	
	10/20/2010	-1.13	±	1.43	-4.18	±	5.29	

TABLE C-2. Weekly Iodine-131 Activity in Air.

Sampling Group and Location	Sampling Date	Result ± 1s Uncertainty (x 10 <sup>-15</sup> µCi/mL)			Result ± 1s Uncertainty (x 10 <sup>-11</sup> Bq/mL)			Result > 3s
<b>BOUNDARY</b>								
	10/27/2010	-1.35	±	1.28	-5.00	±	4.75	
	11/03/2010	1.36	±	1.34	5.03	±	4.95	
	11/10/2010	0.96	±	1.32	3.54	±	4.87	
	11/17/2010	-1.78	±	1.19	-6.59	±	4.42	
	11/24/2010	2.22	±	1.10	8.23	±	4.06	
	12/01/2010	-1.59	±	1.01	-5.88	±	3.73	
	12/08/2010	0.45	±	1.05	1.65	±	3.88	
	12/15/2010	0.00	±	1.05	0.01	±	3.87	
	12/22/2010	-0.31	±	1.04	-1.13	±	3.84	
	12/29/2010	-2.60	±	1.96	-9.61	±	7.26	
<b>VAN BUREN GATE</b>								
	10/06/2010	-0.63	±	1.03	-2.33	±	3.80	
	10/13/2010	-0.40	±	1.06	-1.49	±	3.92	
	10/20/2010	-1.02	±	1.29	-3.78	±	4.78	
	10/27/2010	-1.05	±	1.00	-3.90	±	3.70	
	11/03/2010	1.17	±	1.15	4.31	±	4.24	
	11/10/2010	1.32	±	1.81	4.87	±	6.70	
	11/17/2010	-1.71	±	1.15	-6.34	±	4.25	
	11/24/2010	2.03	±	1.00	7.50	±	3.71	
	12/01/2010	-1.52	±	0.96	-5.63	±	3.57	
	12/08/2010	0.42	±	0.98	1.55	±	3.64	
	12/15/2010	0.00	±	0.96	0.01	±	3.55	
	12/22/2010	-0.29	±	0.97	-1.06	±	3.60	
	12/29/2010	-1.23	±	0.93	-4.55	±	3.43	
<b>QA-1</b>								
	10/06/2010	-0.68	±	1.12	-2.52	±	4.13	
	10/13/2010	-0.36	±	0.95	-1.34	±	3.53	
	10/20/2010	-0.93	±	1.17	-3.44	±	4.35	
	10/27/2010	-1.12	±	1.07	-4.16	±	3.95	
	11/03/2010	1.08	±	1.06	4.00	±	3.93	
	11/10/2010	1.33	±	1.83	4.93	±	6.78	
	11/17/2010	-1.72	±	1.15	-6.37	±	4.27	
	11/24/2010	2.20	±	1.09	8.15	±	4.02	
	12/01/2010	-1.38	±	0.88	-5.11	±	3.24	
	12/08/2010	0.37	±	0.88	1.38	±	3.25	
	12/15/2010	0.00	±	0.87	0.01	±	3.23	
	12/22/2010	-0.26	±	0.89	-0.97	±	3.29	
	12/29/2010	-1.17	±	0.89	-4.35	±	3.28	
<b>a. Invalid Sample Result</b>								



TABLE C-3. Quarterly Cesium-137 and Strontium-90 Concentrations in Composite Air Filters.

Sampling Group and Location	Sampling Date	Analyte	Result ± 1s Uncertainty (x 10 <sup>-18</sup> µCi/mL)		Result ± 1s Uncertainty (x 10 <sup>-13</sup> Bq/mL)		Result > 3s
<b>BOUNDARY</b>							
ARCO	12/29/2010	AMERICIUM-241	1.38	± 0.98	5.11	± 3.62	
		CESIUM-137	-68.20	± 154.00	-252.34	± 569.80	
		PLUTONIUM-238	-0.25	± 0.90	-0.92	± 3.32	
ATOMIC CITY	12/29/2010	PLUTONIUM-239/240	0.75	± 0.56	2.76	± 2.06	
		CESIUM-137	-39.50	± 120.00	-146.15	± 444.00	
		STRONTIUM-90	-27.30	± 18.70	-101.01	± 69.19	
BLUE DOME	12/29/2010	AMERICIUM-241	7.94	± 4.03	29.38	± 14.91	
		CESIUM-137	-168.00	± 128.00	-621.60	± 473.60	
		PLUTONIUM-238	0.00	± 0.11	0.00	± 0.41	
FAA TOWER HOWE	12/29/2010	PLUTONIUM-239/240	0.43	± 0.30	1.57	± 1.11	
		CESIUM-137	-68.90	± 172.00	-254.93	± 636.40	
		AMERICIUM-241	0.00	± 0.10	0.00	± 0.37	
MONTEVIEW	12/29/2010	CESIUM-137	-115.00	± 105.00	-425.50	± 388.50	
		PLUTONIUM-238	-0.30	± 0.79	-1.10	± 2.92	
		PLUTONIUM-239/240	0.00	± 0.10	0.00	± 0.37	
MUD LAKE	12/29/2010	CESIUM-137	120.00	± 129.00	444.00	± 477.30	
		STRONTIUM-90	-4.85	± 10.60	-17.95	± 39.22	
<b>DISTANT</b>							
BLACKFOOT	12/29/2010	CESIUM-137	-187.00	± 132.00	-691.90	± 488.40	
		STRONTIUM-90	-15.90	± 11.50	-58.83	± 42.55	
CRATERS	12/29/2010	AMERICIUM-241	1.40	± 1.40	5.18	± 5.18	
		CESIUM-137	-95.60	± 84.60	-353.72	± 313.02	
		PLUTONIUM-238	2.31	± 1.18	8.55	± 4.37	
DUBOIS	12/29/2010	PLUTONIUM-239/240	0.51	± 0.73	1.90	± 2.69	
		CESIUM-137	-258.00	± 140.00	-954.60	± 518.00	
		CESIUM-137	-51.40	± 139.00	-190.18	± 514.30	
IDAHO FALLS	12/29/2010	CESIUM-137	111.00	± 92.80	410.70	± 343.36	
JACKSON	12/29/2010	AMERICIUM-241	1.80	± 1.80	6.66	± 6.66	
		CESIUM-137	52.10	± 142.00	192.77	± 525.40	
		PLUTONIUM-238	-0.29	± 0.66	-1.08	± 2.43	
REXBURG CMS	12/29/2010	PLUTONIUM-239/240	-0.29	± 0.66	-1.08	± 2.43	
		CESIUM-137	-142.00	± 80.30	-525.40	± 297.11	
		STRONTIUM-90	-10.20	± 10.80	-37.74	± 39.96	
<b>INL SITE</b>							
EFS	12/29/2010	CESIUM-137	9.27	± 103.00	34.30	± 381.10	
MAIN GATE	12/29/2010	CESIUM-137	-236.00	± 151.00	-873.20	± 558.70	
VAN BUREN GATE	12/29/2010	CESIUM-137	-44.70	± 80.60	-165.39	± 298.22	
		STRONTIUM-90	-15.90	± 9.35	-58.83	± 34.60	
VAN BUREN GATE (QA-1)	12/29/2010	CESIUM-137	0.60	± 75.80	0.00	± 0.00	
		STRONTIUM-90	-7.02	± 9.10	-25.97	± 33.67	

TABLE C-4. Tritium Concentrations in Atmospheric Moisture.

Sampling Group and Location	Start Date	Sampling Date	Result ± 1s Uncertainty			Result ± 1s Uncertainty			Collection Medium	Result > 3s
			(x 10 <sup>-13</sup> μCi/mL <sub>air</sub> )			(x 10 <sup>-9</sup> Bq/mL <sub>air</sub> )				
<b>BOUNDARY</b>										
ATOMIC CITY	09/15/2010	10/13/2010	15.75	±	2.30	58.27	±	8.52	Molecular Sieve	Y
ATOMIC CITY	10/13/2010	11/10/2010	3.73	±	1.81	13.80	±	6.70	Molecular Sieve	
ATOMIC CITY	11/10/2010	12/15/2010	5.49	±	1.35	20.31	±	4.99	Molecular Sieve	Y
<b>DISTANT</b>										
BLACKFOOT	09/22/2010	10/13/2010	5.51	±	1.59	20.38	±	5.90	Molecular Sieve	Y
BLACKFOOT	10/13/2010	11/10/2010	11.21	±	1.11	41.47	±	4.10	Molecular Sieve	Y
BLACKFOOT	11/10/2010	12/08/2010	10.29	±	1.18	38.06	±	4.35	Molecular Sieve	Y
IDAHO FALLS	09/22/2010	10/13/2010	8.00	±	1.84	29.58	±	6.82	Molecular Sieve	Y
IDAHO FALLS	10/13/2010	11/03/2010	5.71	±	1.75	21.13	±	6.46	Molecular Sieve	Y
IDAHO FALLS	11/03/2010	11/24/2010	4.98	±	1.68	18.41	±	6.22	Molecular Sieve	
REXBURG	09/29/2010	10/13/2010	8.14	±	3.02	30.12	±	11.18	Molecular Sieve	
REXBURG	10/13/2010	11/03/2010	4.31	±	2.30	15.94	±	8.50	Molecular Sieve	
REXBURG	11/03/2010	12/08/2010	6.97	±	1.92	25.80	±	7.11	Molecular Sieve	Y

**TABLE C-5. Monthly and Weekly Tritium Concentrations in Precipitation.**

Location	Start Date	End Date	Result $\pm$ 1s Uncertainty			Result $\pm$ 1s Uncertainty			Result > 3s
			(pCi/L)			(Bq/L)			
IDAHO FALLS	9/1/2010	10/1/2010	87.20	$\pm$	32.40	3.23	$\pm$	1.20	
	10/1/2010	11/1/2010	137.00	$\pm$	33.60	5.07	$\pm$	1.24	Y
	11/1/2010	12/1/2010	78.00	$\pm$	33.30	2.89	$\pm$	1.23	
CFA	9/2/2010	10/1/2010	106.00	$\pm$	32.50	3.92	$\pm$	1.20	Y
	10/1/2010	11/1/2010	101.00	$\pm$	33.30	3.74	$\pm$	1.23	Y
	11/1/2010	12/1/2010	73.10	$\pm$	35.50	2.70	$\pm$	1.31	
EFS	9/29/2010	10/6/2010	160.00	$\pm$	32.60	5.92	$\pm$	1.21	Y
	10/6/2010	10/13/2010	203.00	$\pm$	33.00	7.51	$\pm$	1.22	Y
	10/20/2010	10/27/2010	94.60	$\pm$	33.60	3.50	$\pm$	1.24	
	11/3/2010	11/10/2010	90.70	$\pm$	35.60	3.36	$\pm$	1.32	
	11/17/2010	11/24/2010	146.00	$\pm$	36.00	5.40	$\pm$	1.33	Y
	11/24/2010	12/1/2010	106.00	$\pm$	35.70	3.92	$\pm$	1.32	

Table C-6. Weekly and Monthly Iodine-131 and Cesium-137 Concentrations in Milk

Location	Sampling Date	Iodine-131				Cesium-137			
		Result ± 1s Uncertainty (pCi/L)		Result ± 1s Uncertainty (Bq/L)		Result ± 1s Uncertainty (pCi/L)		Result ± 1s Uncertainty (Bq/L)	
					Result > 3s				Result > 3s
BLACKFOOT									
	10/12/10	-0.27 ± 1.59	-0.010 ± 0.059			-0.28 ± 0.93	-0.010 ± 0.034		
	11/02/10	1.27 ± 1.22	0.047 ± 0.045			-1.56 ± 0.94	-0.058 ± 0.035		
	12/07/10	0.81 ± 1.81	0.030 ± 0.067			-0.30 ± 0.91	-0.011 ± 0.034		
CONTROL									
	10/05/10	-1.14 ± 1.84	-0.042 ± 0.068			2.56 ± 1.35	0.095 ± 0.050		
	11/02/10	-0.08 ± 1.27	-0.003 ± 0.047			0.30 ± 0.89	0.011 ± 0.033		
	12/09/10	0.81 ± 1.33	0.030 ± 0.049			-0.29 ± 0.90	-0.011 ± 0.033		
DIETRICH									
	10/05/10	-0.64 ± 1.49	-0.024 ± 0.055			3.12 ± 1.35	0.116 ± 0.050		
	11/02/10	2.27 ± 1.49	0.084 ± 0.055			2.54 ± 1.38	0.094 ± 0.051		
	12/07/10	0.18 ± 1.17	0.007 ± 0.043			1.19 ± 0.86	0.044 ± 0.032		
FORT HALL									
	11/02/10	0.83 ± 1.53	0.031 ± 0.057			1.69 ± 1.36	0.063 ± 0.050		
	12/07/10	-0.40 ± 0.98	-0.015 ± 0.036			0.66 ± 0.78	0.024 ± 0.029		
HOWE									
	10/05/10	0.00 ± 0.00	0.000 ± 0.000			4.67 ± 1.33	0.173 ± 0.049		
	11/02/10	0.00 ± 0.00	0.000 ± 0.000			0.06 ± 0.85	0.002 ± 0.032		
	12/07/10	0.00 ± 0.00	0.000 ± 0.000			-1.02 ± 0.91	-0.038 ± 0.034		
Duplicate	12/07/10	0.00 ± 0.00	0.000 ± 0.000			0.43 ± 1.38	0.016 ± 0.051		
IDAHO FALLS									
	10/05/10	-0.17 ± 0.92	-0.006 ± 0.034			0.62 ± 0.79	0.023 ± 0.029		
	10/12/10	-2.43 ± 1.53	-0.090 ± 0.057			0.35 ± 1.35	0.013 ± 0.050		
	10/19/10	3.09 ± 1.49	0.114 ± 0.055			-1.19 ± 1.35	-0.044 ± 0.050		
	10/26/10	0.50 ± 0.92	0.019 ± 0.034			-0.34 ± 0.77	-0.013 ± 0.028		
	11/02/10	1.96 ± 1.88	0.073 ± 0.070			0.97 ± 1.41	0.036 ± 0.052		
	11/09/10	0.16 ± 1.46	0.006 ± 0.054			1.82 ± 1.28	0.067 ± 0.047		
	11/16/10	1.96 ± 0.93	0.073 ± 0.034			0.92 ± 0.81	0.034 ± 0.030		
	11/23/10	1.14 ± 0.91	0.042 ± 0.034			0.76 ± 0.75	0.028 ± 0.028		
	11/30/10	-0.59 ± 0.93	-0.022 ± 0.034			-0.07 ± 0.78	-0.003 ± 0.029		
	12/07/10	-0.28 ± 0.90	-0.010 ± 0.033			2.14 ± 0.78	0.079 ± 0.029		
	12/14/10	0.75 ± 0.90	0.028 ± 0.033			-1.57 ± 0.79	-0.058 ± 0.029		
	12/21/10	0.28 ± 0.82	0.010 ± 0.030			0.31 ± 0.73	0.012 ± 0.027		
	12/28/10	0.97 ± 0.91	0.036 ± 0.034			2.05 ± 0.78	0.076 ± 0.029		
RUPERT									
	10/05/10	0.00 ± 1.75	0.000 ± 0.065			-4.47 ± 1.43	-0.166 ± 0.053		
Duplicate	10/05/10	-0.38 ± 0.96	-0.014 ± 0.035			0.86 ± 0.77	0.032 ± 0.028		
	11/02/10	0.72 ± 0.97	0.027 ± 0.036			0.61 ± 0.77	0.023 ± 0.028		
	12/07/10	0.69 ± 1.73	0.025 ± 0.064			0.68 ± 1.43	0.025 ± 0.053		
TERRETON									
	10/05/10	1.96 ± 1.88	0.073 ± 0.070			0.97 ± 1.41	0.036 ± 0.052		
	11/02/10	-1.37 ± 1.10	-0.051 ± 0.041			-0.41 ± 0.79	-0.015 ± 0.029		
	12/07/10	-0.39 ± 1.04	-0.014 ± 0.039			1.95 ± 0.81	0.072 ± 0.030		

Table C-7. Strontium-90 and Tritium Concentrations in Milk

Strontium-90								
Location	Sampling Date	Result ± 1s Uncertainty			Result ± 1s Uncertainty			Result > 3s
		(pCi/L)			(Bq/L)			
CONTROL	11/02/10	0.42	±	0.08	0.016	±	0.003	Y
DIETRICH	11/02/10	0.63	±	0.09	0.023	±	0.004	Y
FORT HALL	11/03/10	0.24	±	0.10	0.009	±	0.004	
HOWE	11/02/10	0.06	±	0.10	0.002	±	0.004	
IDAHO FALLS	11/02/10	0.86	±	0.10	0.032	±	0.004	Y
RUPERT	11/02/10	0.25	±	0.08	0.009	±	0.003	Y
TERRETON	11/02/10	0.26	±	0.11	0.010	±	0.004	

Tritium								
Location	Sampling Date	Concentration ± 1s			Concentration ± 1s			Result > 3s
		(pCi/L)			(Bq/L)			
BLACKFOOT	11/02/10	112.32	±	34.09	4.160	±	1.262	Y
CONTROL	11/02/10	88.07	±	32.21	3.262	±	1.193	
DIETRICH	11/02/10	64.17	±	32.94	2.377	±	1.220	
FORT HALL	11/02/10	113.21	±	33.93	4.193	±	1.257	Y
HOWE	11/02/10	35.61	±	33.28	1.319	±	1.233	
IDAHO FALLS	11/02/10	75.63	±	33.78	2.801	±	1.251	
RUPERT	11/02/10	219.07	±	34.23	8.114	±	1.268	Y
TERRETON	11/02/10	13.26	±	33.87	0.491	±	1.254	

Table C-8. Cesium-137 and Strontium-90 Concentrations in Lettuce

		Cesium-137						
Location	Sampling Date	Result ± 1s Uncertainty <sup>a</sup>			Result ± 1s Uncertainty <sup>a</sup>			Result > 3s
		pCi/kg <sup>b</sup>			(x 10 <sup>-2</sup> Bq/kg) <sup>b</sup>			
ARCO	07/28/2010	251.00	±	71.50	929.63	±	264.81	Y
ATOMIC CITY	07/28/2010	114.00	±	71.20	422.22	±	263.70	
BLACKFOOT	08/20/2010	22.80	±	34.10	84.44	±	126.30	
CAREY	08/03/2010	-28.50	±	86.30	-105.56	±	319.63	
CAREY (DUPLICATE)	08/03/2010	112.00	±	58.00	414.81	±	214.81	
CONTROL	09/13/2010	30.40	±	55.70	112.59	±	206.30	
EFS	07/28/2010	-32.20	±	111.00	-119.26	±	411.11	
FAA TOWER	09/01/2010	215.00	±	60.60	796.30	±	224.44	Y
HOWE	08/04/2010	85.30	±	52.90	315.93	±	195.93	
IDAHO FALLS	08/09/2010	20.90	±	69.40	77.41	±	257.04	
MONTEVIEW	07/28/2010	218.00	±	75.40	807.41	±	279.26	

		Strontium-90						
Location	Sampling Date	Result ± 1s Uncertainty			Result ± 1s Uncertainty			Result > 3s
		pCi/kg <sup>b</sup>			(x 10 <sup>-2</sup> Bq/kg) <sup>b</sup>			
ARCO	07/28/2010	13.70	±	11.50	50.74	±	42.59	
ATOMIC CITY	07/28/2010	41.10	±	11.80	152.22	±	43.70	Y
BLACKFOOT	08/20/2010	-8.70	±	5.90	-32.22	±	21.85	
CAREY	08/03/2010	-14.30	±	8.70	-52.96	±	32.22	
CAREY (DUPLICATE)	08/03/2010	1.40	±	9.90	5.19	±	36.67	
CONTROL	09/13/2010	23.20	±	8.15	85.93	±	30.19	
EFS	07/28/2010	12.00	±	13.10	44.44	±	48.52	
FAA TOWER	09/01/2010	73.50	±	12.50	272.22	±	46.30	Y
HOWE	08/04/2010	-15.10	±	7.60	-55.93	±	28.15	
IDAHO FALLS	08/09/2010	5.65	±	8.15	20.93	±	30.19	
MONTEVIEW	07/28/2010	30.40	±	13.90	112.59	±	51.48	

<sup>a</sup> A review of the table, performed in the spring of 2021, revealed that the <sup>137</sup>Cs result and uncertainty values listed were incorrect. The <sup>137</sup>Cs result and uncertainty values were updated to the correct values. For further discussion, see Lettuce Sampling Section 5.

<sup>b</sup> During the spring of 2021, a review of the table determined the unit of concentration listed in the column headings were incorrect. The column headings have been updated to the correct units of concentration (pCi/kg and Bq/kg). For further discussion, see Lettuce Sampling in Section 5.

Table C-9. Cesium-137 and Strontium-90 Concentrations in Potatoes

Location	Sampling Date	Cesium-137						
		Result ± 1s Uncertainty pCi/kg			Result ± 1s Uncertainty Bq/kg			Result > 3s
ARCO	9/29/2010	-0.55	±	0.46	-0.02	±	0.02	
BLACKFOOT	9/29/2010	0.07	±	0.51	0.00	±	0.02	
CONTROL	9/21/2010	-0.47	±	0.69	-0.02	±	0.03	
IDAHO FALLS	10/5/2010	0.87	±	0.65	0.03	±	0.02	
MONTEVIEW	9/29/2010	0.95	±	0.51	0.04	±	0.02	
MONTEVIEW (DUPLICATE)	9/29/2010	0.66	±	0.46	0.02	±	0.02	
MUK LAKE	9/29/2010	0.40	±	0.48	0.01	±	0.02	
RUPERT	10/5/2010	-0.27	±	0.49	-0.01	±	0.02	
SHELLEY	9/28/2010	-0.18	±	0.71	-0.01	±	0.03	
TERRETON	10/5/2010	0.54	±	0.69	0.02	±	0.03	
Strontium-90								Result > 3s
Result ± 1s Uncertainty pCi/kg			Result ± 1s Uncertainty Bq/kg					
ARCO	9/29/2010	0.68	±	0.39	0.03	±	0.01	
BLACKFOOT	9/29/2010	0.49	±	0.36	0.02	±	0.01	
CONTROL	9/21/2010	0.45	±	0.46	0.02	±	0.02	
IDAHO FALLS	10/5/2010	-1.35	±	0.47	-0.05	±	0.02	
MONTEVIEW	9/29/2010	0.71	±	0.45	0.03	±	0.02	
MONTEVIEW (DUPLICATE)	9/29/2010	1.21	±	0.47	0.04	±	0.02	
MUK LAKE	9/29/2010	0.79	±	0.43	0.03	±	0.02	
RUPERT	10/5/2010	1.57	±	0.51	0.06	±	0.02	Y
SHELLEY	9/28/2010	0.39	±	0.47	0.01	±	0.02	
TERRETON	10/5/2010	-0.44	±	0.46	-0.02	±	0.02	

Table C-10. Cesium-137 and Iodine-131 Concentrations in Large Game Animals

Species	Collection		Analyte	Result $\pm$ 1s Uncertainty			Result $\pm$ 1s Uncertainty			Result > 3s
	Date	Tissue		(pCi/kg wet weight)			$(\times 10^{-2}$ Bq/kg wet weight)			
PRONGHORN	10/5/2010	Muscle	<sup>131</sup> I	-2.50	$\pm$	9.19	-9.24	$\pm$	34.01	Y
			<sup>137</sup> Cs	2.77	$\pm$	0.88	10.24	$\pm$	3.24	
	Thyroid	<sup>131</sup> I	-5.56	$\pm$	142.00	-20.57	$\pm$	525.40		
		<sup>137</sup> Cs	149.00	$\pm$	150.00	551.30	$\pm$	555.00		
MULE DEER	10/11/2010	Muscle	<sup>131</sup> I	0.16	$\pm$	2.11	0.58	$\pm$	7.82	
			<sup>137</sup> Cs	2.13	$\pm$	1.70	7.90	$\pm$	6.29	
	Thyroid	<sup>131</sup> I	382.00	$\pm$	487.00	1413.40	$\pm$	1801.90		
		<sup>137</sup> Cs	-123.00	$\pm$	461.00	-455.10	$\pm$	1705.70		
MULE DEER	10/12/2010	Muscle	<sup>131</sup> I	2.86	$\pm$	2.71	10.60	$\pm$	10.03	
			<sup>137</sup> Cs	3.21	$\pm$	1.15	11.87	$\pm$	4.24	
ELK	11/15/2010	Liver	<sup>131</sup> I	50.07	$\pm$	33.38	185.26	$\pm$	123.51	
			<sup>137</sup> Cs	4.16	$\pm$	1.40	15.39	$\pm$	5.18	
	Muscle	<sup>131</sup> I	47.95	$\pm$	25.29	177.40	$\pm$	93.58		
		<sup>137</sup> Cs	1.43	$\pm$	1.06	5.28	$\pm$	3.92		
	Thyroid	<sup>131</sup> I	35.90	$\pm$	51.60	132.83	$\pm$	190.92		
		<sup>137</sup> Cs	32.20	$\pm$	52.80	119.14	$\pm$	195.36		



**Table C-11. Gamma-emitting Radionuclides.  
Strontium-90, and Actinides in Waterfowl**

Location	Sampling		Result ± Uncertainty(1s)			Result ± Uncertainty(1s)			
Species	Date	Analyte	pCi/kg <sup>a</sup>			(x 10 <sup>-2</sup> ) Bq/kg <sup>a</sup>			Result > 3s
ATR Complex Mallard	9/17/2010	AMERICIUM-241	-0.06	±	0.18	-0.22	±	0.67	
		CESIUM-137	2737.93	±	59.02	10140.49	±	218.58	Y
		CHROMIUM-51	-302.61	±	503.05	-1120.79	±	1863.13	
		COBALT-60	67.47	±	4.08	249.87	±	15.11	Y
		PLUTONIUM-238	0.45	±	0.14	1.68	±	0.53	Y
		PLUTONIUM-239/240	9.15	±	0.87	33.89	±	3.22	Y
		STRONTIUM-90	318.00	±	5.40	1177.78	±	20.00	Y
		ZINC-65	329.47	±	18.54	1220.25	±	68.65	Y
ATR Complex Mallard	9/17/2010	AMERICIUM-241	-0.04	±	0.04	-0.13	±	0.13	
		CESIUM-137	492.95	±	12.91	1825.74	±	47.80	Y
		CHROMIUM-51	368.22	±	362.80	1363.78	±	1343.70	
		COBALT-60	40.18	±	3.03	148.83	±	11.21	Y
		PLUTONIUM-238	0.26	±	0.16	0.96	±	0.59	
		PLUTONIUM-239/240	3.48	±	0.33	12.89	±	1.21	Y
		STRONTIUM-90	1050.00	±	9.00	3888.89	±	33.33	Y
		ZINC-65	287.42	±	15.35	1064.51	±	56.84	Y
ATR Complex American Coot	9/17/2010	AMERICIUM-241	0.21	±	0.26	0.79	±	0.97	
		CESIUM-137	650.27	±	18.21	2408.42	±	67.46	Y
		CHROMIUM-51	-866.12	±	593.81	-3207.85	±	2199.28	
		COBALT-60	105.65	±	5.78	391.28	±	21.42	Y
		PLUTONIUM-238	0.38	±	0.13	1.42	±	0.49	
		PLUTONIUM-239/240	0.13	±	0.10	0.47	±	0.35	
		STRONTIUM-90	9.08	±	3.46	33.63	±	12.81	
		ZINC-65	521.86	±	27.87	1932.81	±	103.22	Y
MFC Ruddy Duck	9/19/2010	AMERICIUM-241	0.51	±	0.23	1.90	±	0.87	
		CESIUM-137	6.69	±	3.83	24.79	±	14.18	
		CHROMIUM-51	80.66	±	483.98	298.75	±	1792.52	
		COBALT-60	1.49	±	3.00	5.53	±	11.11	
		PLUTONIUM-238	0.00	±	0.00	0.00	±	0.00	
		PLUTONIUM-239/240	0.13	±	0.10	0.47	±	0.35	
		STRONTIUM-90	3.69	±	3.24	13.67	±	12.00	
		ZINC-65	-36.56	±	10.86	-135.42	±	40.21	
MFC American Coot	9/19/2010	AMERICIUM-241	0.13	±	0.14	0.47	±	0.52	
		CESIUM-137	3.30	±	4.50	12.22	±	16.65	
		CHROMIUM-51	-429.23	±	589.75	-1589.74	±	2184.25	
		COBALT-60	6.36	±	5.29	23.55	±	19.61	
		PLUTONIUM-238	-0.11	±	0.13	-0.40	±	0.50	
		PLUTONIUM-239/240	0.33	±	0.15	1.21	±	0.57	
		STRONTIUM-90	2.19	±	2.87	8.11	±	10.63	
		ZINC-65	-0.39	±	11.97	-1.44	±	44.34	
MFC Green-winged Teal	9/19/2010	AMERICIUM-241	0.10	±	0.10	0.37	±	0.37	
		CESIUM-137	5.43	±	7.40	20.12	±	27.42	
		CHROMIUM-51	-706.88	±	971.23	-2618.07	±	3597.14	
		COBALT-60	10.47	±	8.72	38.78	±	32.29	
		PLUTONIUM-238	0.61	±	0.29	2.24	±	1.06	
		PLUTONIUM-239/240	0.43	±	0.26	1.60	±	0.96	
		STRONTIUM-90	10.20	±	5.75	37.78	±	21.30	
		ZINC-65	-0.64	±	19.72	-2.36	±	73.02	

<sup>a</sup> During the spring of 2021, a review of the table determined the activity concentration values reported for the media were correct, however, the (x10<sup>-5</sup>) Bq/g unit of concentration listed in the column headings were incorrect. The column headings have been updated to the correct units of concentration (pCi/kg and (x10<sup>-2</sup>) Bq/kg). For further discussion see Waterfowl sampling Section 5.

**Table C-11. Gamma-emitting Radionuclides.  
Strontium-90, and Actinides in Waterfowl**

Location	Sampling		Result ± Uncertainty(1s)			Result ± Uncertainty(1s)			
Species	Date	Analyte	pCi/kg <sup>a</sup>			(x 10 <sup>-2</sup> ) Bq/kg <sup>a</sup>			Result > 3s
<b>CONTROL</b>	10/30/2010								
Mallard		AMERICIUM-241	0.44	±	0.25	1.64	±	0.91	
		CESIUM-137	3.88	±	1.61	14.38	±	5.95	
		CHROMIUM-51	3.32	±	87.11	12.29	±	322.61	
		COBALT-60	1.02	±	125.92	3.79	±	466.36	
		PLUTONIUM-238	-0.01	±	0.02	-0.05	±	0.09	
		PLUTONIUM-239/240	-0.07	±	0.05	-0.26	±	0.20	
		STRONTIUM-90	-0.97	±	0.90	-3.58	±	3.33	
		ZINC-65	-8.21	±	4.36	-30.40	±	16.13	
<b>CONTROL</b>	11/13/2010								
Common		AMERICIUM-241	0.34	±	0.12	1.26	±	0.45	
Goldeneye		CESIUM-137	1.55	±	1.95	5.74	±	7.24	
		CHROMIUM-51	-41.24	±	68.74	-152.76	±	254.60	
		COBALT-60	1.02	±	1.46	3.78	±	5.40	
		PLUTONIUM-238	0.12	±	0.07	0.46	±	0.27	
		PLUTONIUM-239/240	0.16	±	0.10	0.61	±	0.37	
		STRONTIUM-90	0.84	±	1.06	3.09	±	3.93	
		ZINC-65	-13.97	±	5.14	-51.72	±	19.03	
<b>CONTROL</b>	12/11/2010								
Northern		AMERICIUM-241	0.04	±	0.12	0.15	±	0.44	
Shoveler		CESIUM-137	3.13	±	3.32	11.58	±	12.30	
		CHROMIUM-51	-26.47	±	59.55	-98.02	±	220.54	
		COBALT-60	-2.97	±	240.31	-11.00	±	890.04	
		PLUTONIUM-238	0.22	±	0.16	0.83	±	0.59	
		PLUTONIUM-239/240	0.11	±	0.19	0.41	±	0.71	
		STRONTIUM-90	16.30	±	2.56	60.37	±	9.48	Y
		ZINC-65	-10.67	±	7.86	-39.53	±	29.13	

<sup>a</sup> During the spring of 2021, a review of the table determined the activity concentration values reported for the media were correct, however, the (x10<sup>-5</sup>) Bq/g unit of concentration listed in the column headings were incorrect. The column headings have been updated to the correct units of concentration (pCi/kg and (x10<sup>-2</sup>) Bq/kg). For further discussion see Waterfowl sampling Section 5.

**Table C-12. Cesium-137,  
Strontium-90, and Actinides in Soil**

Location	Sampling Date	Americium-241						Result > 3s
		Concentration ± 1s (pCi/Kg)			Concentration ± 1s (Bq/Kg)			
<b>BOUNDARY</b>								
ATOMIC CITY	07/27/10	-9.04	±	5.24	-0.33	±	0.19	
ATOMIC CITY (DUP)	07/27/10	-11.40	±	29.00	-0.42	±	1.07	
BUTTE CITY	07/27/10	18.40	±	27.70	0.68	±	1.03	
FAA TOWER	07/19/10	23.90	±	27.50	0.89	±	1.02	
FRENCHMAN'S CABIN	07/27/10	17.50	±	14.20	0.65	±	0.53	
HOWE	07/19/10	9.96	±	11.50	0.37	±	0.43	
MONTEVIEW	07/19/10	1.76	±	10.70	0.07	±	0.40	
MUD LAKE #1	07/19/10	19.70	±	17.90	0.73	±	0.66	
MUD LAKE #2	07/19/10	1.73	±	10.60	0.06	±	0.39	
RENO RANCH	07/19/10	-7.77	±	12.20	-0.29	±	0.45	
<b>DISTANT</b>								
BLACKFOOT SOIL	08/02/10	4.18	±	10.80	0.15	±	0.40	
CAREY SOIL	07/27/10	-7.16	±	12.30	-0.27	±	0.46	
CRYSTAL ICE CAVES	08/02/10	-3.09	±	3.11	-0.11	±	0.12	
ST. ANTHONY SOIL	07/19/10	-5.60	±	14.20	-0.21	±	0.53	

Location	Sampling Date	Cesium-137						Result > 3s
		Concentration ± 1s (pCi/Kg)			Concentration ± 1s (Bq/Kg)			
<b>BOUNDARY</b>								
ATOMIC CITY	07/27/10	329.00	±	8.59	12.19	±	0.32	Y
ATOMIC CITY (DUP)	07/27/10	412.00	±	9.24	15.26	±	0.34	Y
BUTTE CITY	07/27/10	496.00	±	10.80	18.37	±	0.40	Y
FAA TOWER	07/19/10	519.00	±	11.40	19.22	±	0.42	Y
FRENCHMAN'S CABIN	07/27/10	359.00	±	8.01	13.30	±	0.30	Y
HOWE	07/19/10	266.00	±	7.13	9.85	±	0.26	Y
MONTEVIEW	07/19/10	264.00	±	6.16	9.78	±	0.23	Y
MUD LAKE #1	07/19/10	343.00	±	7.78	12.70	±	0.29	Y
MUD LAKE #2	07/19/10	238.00	±	5.78	8.81	±	0.21	Y
RENO RANCH	07/19/10	573.00	±	12.50	21.22	±	0.46	Y
<b>DISTANT</b>								
BLACKFOOT SOIL	08/02/10	369.00	±	9.37	13.67	±	0.35	Y
CAREY SOIL	07/27/10	186.00	±	4.95	6.89	±	0.18	Y
CRYSTAL ICE CAVES	08/02/10	460.00	±	10.30	17.04	±	0.38	Y
ST. ANTHONY SOIL	07/19/10	710.00	±	15.10	26.30	±	0.56	Y

Location	Sampling Date	Plutonium-238						Result > 3s
		Concentration ± 1s (pCi/Kg)			Concentration ± 1s (Bq/Kg)			
<b>BOUNDARY</b>								
ATOMIC CITY	07/27/10	4.52	±	27.60	0.17	±	1.02	
ATOMIC CITY (DUP)	07/27/10	7.10	±	4.96	0.26	±	0.18	
BUTTE CITY	07/27/10	16.60	±	11.80	0.61	±	0.44	
FAA TOWER	07/19/10	-1.66	±	5.81	-0.06	±	0.22	
FRENCHMAN'S CABIN	07/27/10	110.00	±	48.60	4.07	±	1.80	
HOWE	07/19/10	-7.26	±	6.85	-0.27	±	0.25	
MONTEVIEW	07/19/10	-2.69	±	5.30	-0.10	±	0.20	
MUD LAKE #1	07/19/10	1.51	±	9.23	0.06	±	0.34	
MUD LAKE #2	07/19/10	44.00	±	32.00	1.63	±	1.19	
RENO RANCH	07/19/10	-1.58	±	9.91	-0.06	±	0.37	
<b>DISTANT</b>								
BLACKFOOT SOIL	08/02/10	-37.30	±	11.30	-1.38	±	0.42	
CAREY SOIL	07/27/10	4.28	±	7.81	0.16	±	0.29	
CRYSTAL ICE CAVES	08/02/10	-16.20	±	10.70	-0.60	±	0.40	
ST. ANTHONY SOIL	07/19/10	65.70	±	34.50	2.43	±	1.28	

**Table C-12. Cesium-137,  
Strontium-90, and Actinides in Soil**

Location	Sampling Date	Plutonium-239/240						Result > 3s
		Concentration ± 1s (pCi/Kg)			Concentration ± 1s (Bq/Kg)			
<b>BOUNDARY</b>								
ATOMIC CITY	07/27/10	-12.60	±	21.70	-0.47	±	0.80	
ATOMIC CITY (DUP)	07/27/10	16.20	±	6.80	0.60	±	0.25	
BUTTE CITY	07/27/10	18.90	±	19.50	0.70	±	0.72	
FAA TOWER	07/19/10	12.40	±	8.62	0.46	±	0.32	
FRENCHMAN'S CABIN	07/27/10	16.70	±	27.80	0.62	±	1.03	
HOWE	07/19/10	7.83	±	6.09	0.29	±	0.23	
MONTEVIEW	07/19/10	5.49	±	6.70	0.20	±	0.25	
MUD LAKE #1	07/19/10	20.90	±	11.70	0.77	±	0.43	
MUD LAKE #2	07/19/10	2.18	±	19.50	0.08	±	0.72	
RENO RANCH	07/19/10	5.73	±	5.91	0.21	±	0.22	
<b>DISTANT</b>								
BLACKFOOT SOIL	08/02/10	9.36	±	15.00	0.35	±	0.56	
CAREY SOIL	07/27/10	8.67	±	6.06	0.32	±	0.22	
CRYSTAL ICE CAVES	08/02/10	16.00	±	12.50	0.59	±	0.46	
ST. ANTHONY SOIL	07/19/10	-5.89	±	25.60	-0.22	±	0.95	
<b>Strontium-90</b>								
Location	Sampling Date	Strontium-90						Result > 3s
		Concentration ± 1s (pCi/Kg)			Concentration ± 1s (Bq/Kg)			
<b>BOUNDARY</b>								
ATOMIC CITY	07/27/10	45.30	±	11.40	1.68	±	0.42	Y
ATOMIC CITY (DUP)	07/27/10	46.30	±	10.30	1.71	±	0.38	Y
BUTTE CITY	07/27/10	44.80	±	12.80	1.66	±	0.47	Y
FAA TOWER	07/19/10	9.68	±	14.00	0.36	±	0.52	
FRENCHMAN'S CABIN	07/27/10	43.20	±	11.00	1.60	±	0.41	Y
HOWE	07/19/10	20.40	±	11.40	0.76	±	0.42	
MONTEVIEW	07/19/10	84.40	±	12.80	3.13	±	0.47	Y
MUD LAKE #1	07/19/10	41.70	±	13.10	1.54	±	0.49	Y
MUD LAKE #2	07/19/10	-28.90	±	12.80	-1.07	±	0.47	
RENO RANCH	07/19/10	23.00	±	13.90	0.85	±	0.51	
<b>DISTANT</b>								
BLACKFOOT SOIL	08/02/10	12.60	±	12.10	0.47	±	0.45	
CAREY SOIL	07/27/10	5.15	±	10.70	0.19	±	0.40	
CRYSTAL ICE CAVES	08/02/10	36.10	±	13.80	1.34	±	0.51	
ST. ANTHONY SOIL	07/19/10	43.20	±	11.00	1.60	±	0.41	Y

Table C-13. Environmental Radiation Measurements

Location	Start Date	End Date	Radiation Measurement ± 2s Uncertainty mR	Exposure mR/day
<b>BOUNDARY</b>				
ARCO	5/5/2010	11/3/2010	66.3 ± 13.0	0.36
ATOMIC CITY	5/5/2010	11/3/2010	65.1 ± 12.8	0.36
BIRCH CREEK	5/5/2010	11/3/2010	54.5 ± 10.7	0.30
BLUE DOME	5/5/2010	11/3/2010	52.8 ± 10.3	0.29
HOWE	5/5/2010	11/3/2010	59.2 ± 11.6	0.33
MONTEVIEW	5/5/2010	11/3/2010	60.7 ± 11.9	0.33
MUD LAKE	5/5/2010	11/3/2010	67.8 ± 13.3	0.37
<b>Boundary Average</b>				<b>0.33</b>
<b>DISTANT</b>				
ABERDEEN	5/4/2010	11/2/2010	67.2 ± 13.2	0.37
BLACKFOOT	5/5/2010	11/3/2010	60.3 ± 11.8	0.33
BLACKFOOT CMS	5/5/2010	11/3/2010	55.8 ± 10.9	0.31
CRATERS	5/5/2010	11/3/2010	65.1 ± 12.8	0.36
DUBOIS	5/5/2010	11/3/2010	52.8 ± 10.3	0.29
IDAHO FALLS	5/5/2010	11/3/2010	63.7 ± 12.5	0.35
MINIDOKA	5/6/2010	11/2/2010	58.2 ± 11.4	0.32
REXBURG	5/5/2010	11/3/2010	82.3 ± 16.1	0.45
ROBERTS	5/4/2010	11/2/2010	70.9 ± 13.9	0.39
<b>Distant Average</b>				<b>0.35</b>
<b>OUT-OF-STATE</b>				
JACKSON	5/6/2010	10/29/2010	50.4 ± 9.9	0.29

**APPENDIX D**  
***STATISTICAL ANALYSIS RESULTS***

**Table D-1. Results of the Kruskal-Wallace statistical test between INL Site, Boundary and Distant sample groups by month.**

<b>Parameter</b>	<b>P<sup>a</sup></b>
<b>Gross Alpha</b>	
Quarter	0.71
October	0.36
November	0.42
December	0.56
<b>Gross Beta</b>	
Quarter	0.62
October	0.95
November	0.47
December	0.95

a. A 'p' value greater than 0.05 signifies no statistical difference between data groups.

**Table D-2. Statistical difference in weekly gross alpha and gross beta concentrations measured at Boundary and Distant locations.**

Parameter	Mann-Whitney U test	
	Week	P <sup>a</sup>
<b>Gross Alpha</b>		
	October 6	0.05
	October 13	0.62
	October 20	0.89
	October 27	0.12
	November 3	0.20
	November 10	0.67
	November 17	0.13
	November 24	0.47
	December 1	1.00
	December 8	0.05
	December 15	0.32
	December 22	0.78
	December 29	0.39
<b>Gross Beta</b>		
	October 6	0.20
	October 13	0.17
	October 20	0.62
	October 27	0.57
	November 3	0.09
	November 10	0.48
	November 17	0.07
	November 24	0.36
	December 1	0.34
	December 8	0.63
	December 15	0.67
	December 22	0.57
	December 29	0.15

a. A 'p' value greater than 0.05 signifies no statistical difference between data groups.